

Identification and antimicrobial activity of lactic acid bacteria from the digestive tract of eels (*Monopterus albus*)

RIDWAN¹, ENDAH RETNANINGRUM^{2,*}, MIFTAHUL ILMI², BUDI SETIADI DARYONO³

¹Graduate Program in Biology, Faculty of Biology, Universitas Gadjah Mada. Jl. Teknik Selatan, Sekip Utara, Sleman 55281, Yogyakarta, Indonesia

²Microbiology Laboratory, Faculty of Biology, Universitas Gadjah Mada. Jl. Teknik Selatan, Sekip Utara, Sleman 55281, Yogyakarta, Indonesia.
Tel./fax.: +62-274-580839, *email: endahr@ugm.ac.id

³Molecular Genetics and Breeding Laboratory, Faculty of Biology, Universitas Gadjah Mada. Jl. Teknik Selatan, Sekip Utara, Sleman 55281, Yogyakarta, Indonesia

Manuscript received: 4 January 2019. Revision accepted: 12 May 2019.

Abstract. Ridwan, Retnaningrum E, Ilmi M, Daryono BS. 2019. Identification and antimicrobial activity of lactic acid bacteria from the digestive tract of eels (*Monopterus albus*). *Bioteknologi* 16: 5-10. This study aims to identify the species of lactic acid bacteria from the digestive tract of eels (*Monopterus albus*) which have inhibitory activity against pathogenic bacteria. The eel used in this study was obtained from the eel cultivation site in Yogyakarta. The isolation results obtained thirteen isolates; the isolated bacteria were then phenotypically characterized to select lactic acid bacteria. The results of phenotypic characterization obtained seven isolates that were in accordance with the character of lactic acid bacteria. Seven lactic acid bacterial isolates were then screened to select the three best isolates in inhibiting the growth of pathogenic bacteria (*Aeromonas hydrophila*, *Staphylococcus aureus*, and *Vibrio harveyi*). The results showed that seven isolates of lactic acid bacteria had antimicrobial activity, which were able to inhibit the growth of pathogenic bacteria. Three of the best isolates were selected to be used for molecular identification and monitoring of antimicrobial activity. The results showed that the three selected isolates produced an antimicrobial extract which was able to inhibit the growth of pathogenic bacteria. Furthermore, the results of the molecular identification showed that the three selected isolates were *Lactococcus lactis* species.

Keywords: 16s rRNA gene, antimicrobial, eel, lactic acid bacteria, *Monopterus albus*

INTRODUCTION

Lactic acid bacteria have various important roles for host health; for example, stimulating the host's immune system (Gatesoupe 2008). In addition, lactic acid bacteria have antimicrobial activity due to the production of lactic acid, acetic acid, bacteriocin, bacteriocin-like compounds and bioactive molecules such as ethanol, formic acid, fatty acids, hydrogen peroxide, diacetyl, reuterin, reutericyclin (Awen et al. 2012). The antimicrobial activity of acid bacteria in inhibiting the growth of pathogenic bacteria attracted the attention of researchers to isolate these bacteria in various sources.

A source that can be used to isolate lactic acid bacteria is the digestive tract of fish. Previous researchers have used the digestive tract of fish to isolate lactic acid bacteria. Lactic acid bacteria isolated from the digestive tract of fish generally have antimicrobial activity, which is stable to high temperatures, pH and stable to organic solvents such as chloroform. Loh et al. (2017) reported that *Lactococcus lactis* subsp. *lactis* CF4MRS isolated from the digestive tract of fish can inhibit bacterial growth of *Escherichia coli*, *Edwardsiella tarda*, *Aeromonas hydrophila*, *Pseudomonas fluorescens*, *Pseudomonas aeruginosa*, *Klebsiella pneumonia* and *Serratia marcescens*.

One group of fish that can be used as a source for isolating lactic acid bacteria are eels (*Monopterus albus*). Eels are aquaculture animals that are very tolerant of low dissolved oxygen content and can live in temperature

which are close to freezing (Roy 2013). The ability of eel (*Monopterus albus*) to live on low oxygen content and near-freezing temperatures is likely due to the presence of superior lactic acid bacteria compared to other aquaculture animals. So, further research regarding the antimicrobial characterization of lactic acid bacteria from the eel digestive tract (*Monopterus albus*) is needed. In addition, molecular identification is needed to find out the species of lactic acid bacteria which have the best antimicrobial activity.

MATERIALS AND METHODS

Procedures

Isolation of lactic acid bacteria

Isolation of lactic acid bacteria in this study used eel (*Monopterus albus*) obtained from the Pingit Market, Yogyakarta, Indonesia. Three eels were taken to the Microbiology Laboratory, Faculty of Biology, University Gadjah Mada, Yogyakarta. The eels were then differentiated and the intestines were taken and then mashed using mortar. After that, isolation was carried out using a dilution technique with the pour plate method. The sample was diluted using sterile distilled water, and then a 400 μ L 10^{-2} to 10^{-6} diluent was put into a petri dish and then added to MRSA liquid. Petri dishes that had the sample added and MRSA media, were then incubated at room temperature for 2 x 24 hours. The growing bacterial

colonies were selected based on the formation of clear zones around the colonies for purification. The purification results were propagated on the sloping MRSA media and stored in the refrigerator as stock for further research (Bajpai et al. 2016).

Screening of inhibitory activity of lactic acid bacteria

The screening of lactic acid bacteria used the Well diffusion method. This involved making a 5 mm well in the NA media (agar nutrient) in a petri dish which had been previously tested by bacterial culture (*Aeromonas hydrophila*, *Vibrio harveyi*, *Staphylococcus aureus*) using the pour plate method. Next, 50 µL of lactic acid bacteria culture was added into the well. Then the well was incubated for 24-48 hours at room temperature. The clear zone formed after incubation was measured as the activity of inhibiting lactic acid bacteria (Edalatian et al. 2015).

Antimicrobial production

Lactic acid bacteria culture was grown in 20 mL MRSB media, then incubated in the shaker incubator for 24 hours at 37°C. After 24 hours incubation, the culture was centrifuged at 10,000 rpm at 10°C for 15 minutes. Then the supernatant was taken as a crude antimicrobial extract and stored in the refrigerator (Hyronimus et al. 1998, Bizani and Brandelli 2002).

Cell-Free Supernatant (CFS) test

Test of CFC's activity used the Well diffusion method. First this involved making a 5 mm well on NA media (agar nutrient) in a petri dish which had previously been added to test bacteria (*Aeromonas hydrophila*, *Vibrio harveyi*, *Staphylococcus aureus*) using the pour plate method. Followed by adding 50 µL of antimicrobial crude extract, which was filtered using a bacterial filter measuring 0.22 µL pore. Then the well was incubated for 24-48 hours at room temperature. The clear zone formed was measured and calculated using the formula:

$$\text{Inhibition Score (IS)} = \frac{\text{Ø clear zone of inhibition (mm)}(\text{mm})}{\text{Ø size of well (mm)}}$$

Antimicrobial inhibitory activity of CFC (Cell-Free Supernatant) is said to be low ($1 < \text{IS} < 3$), moderate ($3 \leq \text{IS} < 5$), strong ($5 \leq \text{IS} < 7$), or very strong ($7 \leq \text{IS} < 9$) (Tremonte et al. 2017).

Molecular identification

Bacterial DNA extraction

DNA extraction was carried out by following the procedure contained in the Geneaid Presto™ Mini gDNA Bacteria Kit. This was to prepare bacterial culture on MRSB media for 48 hours. Bacterial culture of 1.5 mL was then transferred to the microcentrifugation tube. Then it was centrifuged for 1 minute at a speed of 14-16,000 x g and then the supernatant was removed. 200 µL gram (+) buffer was inserted into the centrifuge tube and lysozyme (4 mg/mL) was added. Then it was vortexed until lysozyme was completely dissolved. A gram (+) of buffer solution

that has been added to lysozyme was put into the sample then the pellet was resuspended using a vortex or pipette. The samples were then incubated at 37 ° C for 20 minutes, and it was reversed several times during incubation. Next, 20 µL proteinase K was added and vortexed until homogeneous. It was incubated at 60°C for 10 minutes, and it was reversed every 3 minutes during incubation. Next the sample was added with 200 µL GB buffer before being vortexed. Afterwards, it was incubated for 10 minutes at 70°C. Next the sample was added with 200 µL ethanol absolute and then was vortexed for 10 seconds. It was then put into GD column and centrifuged at a speed of 14-16,000 g for 2 minutes. The previously used tube was then replaced with a new one. After that, 400 µL of W1 buffer was added and was centrifuged at a speed of 14-16,000 g for 30 seconds. Then 600 µL of W1 buffer was added and centrifuged at a speed of 14-16000 g for 30 seconds. Next the GD column was transferred into a sterile Eppendorf tube and added with 100 µL of pre-heated elution. Lastly, it was centrifuged at a speed of 14-16000 g for 3 minutes. The supernatant which was still in the Eppendorf tube was DNA extract (Geneaid Biotech Ltd., 2017).

Amplification of the 16S rRNA gene Lactic Acid Bacteria

The amplification reaction was carried out in a PCR tube. The amplification used TaKara Eq Taq as much as 0.25 µL, 10 x Ex Taq Buffer (containing Mg²⁺ + 2.5 mM) as much as 5 µL, dNTP mixture (2.5 mM) as much as 4 µL, primer 27f (5'-AGAGTTTAGTCCTGGCT CAG-3') and 1492r (5'-GGTTACCTTGTTA CACTT-3') with 0.5 µL each, 1 µg of the genome extract and 10 µL of distilled water added. Amplification was carried out to obtain a single band. The amplification process was initiated with pre-denaturation at 96°C for 4 minutes, followed by 30 cycles at 94°C for 1 minute, and annealing at 51.5°C for 1.30 minutes. After extension at 68°C for 8 minutes, PCR was completed by elongation for 10 minutes at 68 °C. PCR products were then taken and stored at 4 ° C and examined using 1% agarose (Lawalata et al. 2011).

Agarose electrophoresis

1% agarose was heated to boiling with TAE buffer solvent (tris acetate). After it was cooled and poured it into the electrophoresis tray, it was left to be solidified. Then the comb was lifted. Furthermore, into the electrophoresis tool, 1x TAE buffer was poured. The tray containing the solid agarose was then placed on electrophoresis until it was submerged by a buffer. The loading dye was prepared on top of the parafilm then was mixed with 4 µL PCR products and put into agarose well using a micropipette. In addition, a 3 µL marker was prepared. The electrophoresis device was conducted at a voltage of 100 volts for 30 minutes. Agarose results from electrophoresis were immersed in an ethidium bromide dye solution for 10 minutes, then they were visualized in the transilluminator. The presence of DNA bands with a size of ± 1500 bp indicated that the 16S rRNA gene has been amplified, then the results were documented using a camera.

Data analysis

The results of 16S rRNA gene sequencing were analyzed on the website of www.ncbi.nlm.nih.gov. The BLAST program was used to determine the similarity of nucleotide sequences so that identification can be done based on the similarity of nucleotide sequences. Then Phylogeny Tree Construction was carried out to determine the level of closeness between species using the application of MEGAX (Kumar et al. 2018).

RESULTS AND DISCUSSION

Isolation of lactic acid bacteria

Thirteen isolates from this study were phenotypically characterized. Phenotypic characterization being carried out was colony morphology, cell morphology, catalase test and gram staining properties. The results of the phenotypic characteristics of the thirteen bacterial isolates are listed in Table 1.

The results of phenotypic characterization listed in Table 1 show that there are seven bacterial isolates that are suitable for the characterization of lactic acid bacteria. These seven isolates were then screened to find three isolates that had the best antimicrobial activity in inhibiting the growth of test bacteria (*Aeromonas hydrophila*, *Staphylococcus aureus*, and *Vibrio harveyi*).

Screening of inhibitory activity of lactic acid bacteria

Screening the ability of lactic acid bacteria is carried out on seven bacterial isolates. This screening aims to select lactic acid bacteria which have the best activity in inhibiting the growth of pathogenic bacteria (*Aeromonas hydrophila*, *Staphylococcus aureus* and *Vibrio harveyi*). The inhibition ability is best seen based on the diameter of the clear zone formed around the well (Figure 1). The results of screening for inhibitory activity of lactic acid bacteria are listed in Table 2.

Table 1. Characteristics of lactic acid bacteria strain isolated from eels (*Monopterus albus*).

Character	Bacterial Isolates												
	B1A	B1B	B1C	B1D	B1E	B1F	B1G	B1H	B1I	B1J	B1K	B1L	B1M
Cell shape	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci	Cocci
Gram strain	+	+	+	+	+	+	+	-	-	+	-	-	-
Catalase	-	-	-	-	-	-	-	+	+	+	+	+	+
Color													
White	-	-	+	+	-	+	+	+	+	+	+	+	+
Cream	+	+	-	-	+	-	-	-	-	-	-	-	-
Size													
Small	+	+	+	+	-	+	-	+	+	-	-	+	+
Pintpoint	-	-	-	-	+	-	+	-	-	+	+	-	-
Elevation													
Convex	+	-	+	+	+	+	+	+	+	+	+	+	+
Flat	-	+	-	-	-	-	-	-	-	-	-	-	-




Figure 1. Zone of inhibition of lactic acid bacteria at the screening stage i.e.: A. Inhibition zone of *Aeromonas hydrophila*, B. Inhibition zone of *Vibrio harveyi*, C. Zone of inhibition of *Staphylococcus aureus*.

Table 2. Screening results of seven isolates of lactic acid bacteria isolated from the eel digestive tract (*Monopterus albus*).

Strain code	Ø Well (mm)	Ø Mean inhibitory zone (mm)		
		<i>Vibrio harveyi</i>	<i>Aeromonas hydrophila</i>	<i>Staphylococcus aureus</i>
B1A	5	23.33 ± 1.15 ^{ab}	11.67 ± 0.58 ^a	16.67 ± 0.58 ^a
B1B	5	21.00 ± 1.00 ^{cd}	11.33 ± 1.15 ^a	14.67 ± 1.15 ^{ab}
B1C	5	21.00 ± 3.46 ^{cd}	12.33 ± 2.31 ^a	9.00 ± 1.73 ^c
B1D	5	25.33 ± 2.89 ^a	12.33 ± 2.52 ^a	13.00 ± 2.65 ^b
B1E	5	22.33 ± 3.21 ^{bc}	12.00 ± 3.00 ^a	11.33 ± 1.15 ^{bc}
B1F	5	21.00 ± 1.73 ^{cd}	11.33 ± 1.15 ^a	11.67 ± 2.08 ^{bc}
B1G	5	18.67 ± 0.58 ^d	11.67 ± 0.58 ^a	13.67 ± 4.04 ^{ab}
K-	5	-	-	-

Note: Ø = Diameter. Different letters in the same column show the level of real difference – Duncan test $\alpha < 0.05$ (mean ± std)

Table 2 shows that antimicrobial extract of the three isolates had inhibitory activity against the test bacteria (*Aeromonas hydrophila*, *Staphylococcus aureus* and *Vibrio harveyi*). The inhibition activities of seven isolates of lactic acid bacteria were then analyzed statistically using SPSS 17.0 software. The results of statistical analysis showed that the inhibitory activity of lactic acid bacteria had a significant effect on inhibiting the growth of pathogenic bacteria. Ringo (2005) states that lactic acid bacteria isolated from aquatic animals can inhibit the growth of pathogenic bacteria such as *Aeromonas hydrophila* and *Vibrio harveyi*. Buntin et al. (2008) reported that lactic acid bacteria isolated from the digestive tract of fish were able to inhibit the growth of *Staphylococcus aureus* bacteria. Based on the results of the screening, three isolates were selected for further testing. Selected isolates in this study were B1A, B1D and B1E.

Cell-Free Supernatant (CFS) test


The CFC test was conducted to determine the activity of antimicrobial crude extract in inhibiting the growth of test bacteria. The antimicrobial crude extract was obtained from 24-hour-old culture of lactic acid bacteria. Antimicrobial extracts were filtered using a bacterial filter measuring 0.22 µL pore to separate cells from

supernatants. The crude extract filtered was tested for its ability to inhibit the growth of test bacteria. The results of the antimicrobial crude extract activity test are listed in Table 3.

Based on Table 3, the ability of antimicrobial extract from the three bacterial isolates is known now. The results showed that the antimicrobial extract of the three isolates had inhibitory activity against the test bacteria. The three isolates had an Inhibitor Score (IS) <3 or a weak category against *Aeromonas hydrophila* and *Vibrio harveyi*. Furthermore, isolates B1A and B1D have Inhibitor Score (IS) <3 or weak categories while isolates B1E have Inhibitor Score (IS) > 3 or moderate categories against *Staphylococcus aureus* bacteria.

Molecular identification

Molecular identification aims to determine the isolate species of lactic acid bacteria. Molecular identification used the 16S rRNA gene. Amplification of the 16S rRNA gene used 27F primers (5'-AGAGTTTA GTCCTGGCTCAG-3') and 1492R (5'-GGTT ACCTTGTTACGACTT-3'). The amplification results are listed in Figure 2.

**Figure 2.** Results of amplification of the 16S rRNA gene.**Table 3.** Test results for antimicrobial crude extract activity.

Test bacteria	Strain code	Ø Well (mm)	Ø Inhibition zona (mm)			Mean Ø Inhibition zona (mm)	Inhibition Score (IS)
			U1	U2	U3		
<i>Vibrio harveyi</i>	B1A	5	13	15	14	14.00 ± 1.00	2.80 ^a
	B1D	5	14	13	13	13.33 ± 0.58	2.67 ^a
	B1E	5	15	15	14	14.67 ± 0.58	2.93 ^a
	K-	5	-	-	-	-	-
<i>Aeromonas hydrophila</i>	B1A	5	13	14	14	13.67 ± 0.58	2.73 ^a
	B1D	5	14	15	14	14.33 ± 0.58	2.86 ^a
	B1E	5	14	15	15	14.67 ± 0.58	2.93 ^a
	K-	5	-	-	-	-	-
<i>Staphylococcus aureus</i>	B1A	5	15	15	14	14.67 ± 0.58	2.93 ^a
	B1D	5	15	14	15	14.33 ± 0.58	2.86 ^a
	B1E	5	14	16	16	15.78 ± 1.15	3.16 ^b
	K-	5	-	-	-	-	-

Note: Ø = diameter. ^a = weak, ^b = moderate, ^c = strong




Figure 3. Tree of phylogeny isolates of lactic acid bacteria from eel (*Monopterus albus*) (bootstrap = 1000)

The results of 16S rRNA gene amplification were then sequenced using the same primer as the amplification process, 27F and 1492R primers. The sequencing results were then analyzed using BLAST on the site of www.ncbi.nlm.nih.gov with a database of ribosomal 16S rRNA genes. The BLAST analysis results showed that the three isolates had 99% similarity with *Lactococcus lactis* species. Then the phylogenetic tree construction was carried out to determine the evolutionary relationship between bacterial species from isolation and BLAST results. The phylogenetic tree was reconstructed using MEGAX software. The results of the phylogenetic tree reconstruction are shown in Figure 3.

The phylogeny tree in Figure 3 shows that isolates B1A, B1D, and B1E isolated from the eel digestive tract were in one cluster with *Lactococcus lactis*. The results showed that the isolates of lactic acid bacteria isolated from the eel digestive tract (*Monopterus albus*) are a species of *Lactococcus lactis*. Marrifield et al. (2014) reported that lactic acid bacteria include indigenous bacteria in the digestive tract of fish. Lactic acid bacteria that occupy the digestive tract of fish include *Lactococcus lactis*, *Streptococcus* spp., *Weissella cibaria*, and *Lactobacillus* spp. Kato et al. (2016) reported that *Lactococcus* spp. from the digestive tract of fish have antimicrobial activity and can inhibit the growth of *Staphylococcus aureus* (inhibitory zone 6 ± 0.17), *Escherichia coli* (inhibition zone 7 ± 0.10) and *Proteus* spp. (3.5 ± 0.10).

Discussion

Lactic acid bacteria have many important roles; for example, it can stimulate the host's immune system (Verschuere et al. 2000). It can also produce antimicrobial compounds that can inhibit the growth of harmful pathogenic bacteria (Muthukumar and Kandepan 2015). Previous studies by Balcazar et al. (2008) reported that lactic acid bacteria isolated from the digestive tract of fish were able to inhibit the growth of pathogenic bacteria such as *Aeromonas hydrophila*, *Aeromonas salmonicida*, *Yersinia ruckeri* dan *Vibrio anguillarum*.

This study shows that isolated lactic acid bacteria from the eel digestive tract could potentially inhibit the growth of pathogenic bacteria. Lactic acid bacteria that have the potential to inhibit the growth of pathogenic bacteria are

then identified molecularly. The purpose of this study was to find lactic acid bacteria that can inhibit the growth of *Vibrio harveyi*, *Staphylococcus aureus*, and *Aeromonas hydrophila*. *Staphylococcus aureus* is one of the food pathogenic bacteria found throughout the world. Then *Vibrio harveyi* is one of the causes of gastroenteric disease and tissue invasion, especially in hosts who experience a decline in the immune system (Auerbach et al. 2016). *Aeromonas hydrophila* is one of the causes of infectious diseases in fish (Percival and Williams 2014). This study found that lactic acid bacteria were able to inhibit the growth of *Vibrio harveyi*, *Staphylococcus aureus*, and *Aeromonas hydrophila*. These lactic acid bacteria are included in the species *Lactococcus lactis*.

ACKNOWLEDGEMENTS

We thank all of those who have given criticism and suggestions for improving this manuscript. Thanks to the Microbiology Laboratory and the Falitma Laboratory, Faculty of Biology, Gadjah Mada University, Indonesia for being willing to facilitate research.

REFERENCES

- Auerbach PS, Cushing TA, Harris NS. 2016. Auerbach's Wilderness Medicine, 7th ed. Elsevier, New York.
- Aween M, Hassan Z, Belal M. 2012. Evaluation on antibacterial activity of *Lactobacillus acidophilus* strains isolated from honey. *Amer J Appl Sci* 9: 807-817.
- Bajpai VK, Han JH, Nam GJ, Majumder R, Park C, Lim J, Paek WK, Rather IA, Park YH. 2016. Characterization and pharmacological potential of *Lactobacillus sakei* 111 isolated from freshwater fish *Zacco koreanus*. *DARU J Pharamaceut Sci* 24: 8. DOI: 10.1186/s40199-016-0147-8
- Balcazar JL, Vendrell D, de Blas I, Girones O. 2008. Characterization of probiotic properties of lactic acid bacteria isolated from intestinal microbiota of fish. *Aquaculture* 278 (1-4): 188-191.
- Bizani D, Brandelli A. 2002. Characterization of a bacteriocin produced by a newly isolated *Bacillus* sp. *Atrain* 8A. *J Appl Microbiol* 93: 512-519.
- Buntin N, Chanthachum S, Hongpattarakere T. 2008. Screening of lactic acid bacteria from gastrointestinal tracts of marine fish for their potential use as probiotics. *Songklanakarinn J Sci Technol* 30 (1): 141-148.

- Edalatian MR, Habibi NMB, Mortazavi A. 2015. Identification of antimicrobial producing *Enterococci* isolated from Iranian raw milk cheeses using polyphasic approach. *Appl Food Biotechnol* 2(1): 19-24.
- Gatesoupe FJ. 2008. Updating the importance of lactic acid bacteria in fish farming: natural occurrence and probiotic treatments. *J Mol Microbiol Biotechnol* 14 (1-3): 107-14.
- Geneaid Biotech Ltd. 2017. Presto™ Mini gDNA Bacteria Kit, Geneaid Instruction Manual, For Research Use Only, Ver. 02.10.
- Hyronymous BC, Marrec L, Urdaci MC. 1998. Coagulatin, a bacteriocin-like inhibitory substance produced by *Bacillus coagulans*14. *J Appl Microbiol* 35: 42-50.
- Kato CD, Kahuma CE, Namulawa VT, Nasser Kasozi N. 2016. Antibacterial activity of *Lactobacillus* spp. and *Lactococcus* spp. isolated from various parts of pebbly fish, *Alestes baremoze*. *British Microbiol Res J* 17 (2): 1-7.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35 (6): 1547-1549.
- Lawalata HJ, Sembiring L, Sembiring ES. 2011. Molecular identification of lactic acid bacteria producing antimicrobial agents from Bakasang, An Indonesian traditional fermented fish product. *Indonesian J Biotechnol* 16 (2): 93-99.
- Loh JY, Lim YY, Ting ASY. 2017. Bacteriocin-like substances produced by *Lactococcus lactis* subsp. *lactis* CF4MRS isolated from fish intestine: Antimicrobial activities and inhibitory properties. *Intl Food Res J* 24 (1): 394-400.
- Merrifield DL, Balcazar J, Daniels C, Zhou Z, Carnevali O, Sun Y, Hoseinifar SH, Ringo E. 2014. Indigenous lactic acid bacteria in fish and crustaceans. In: *Aquaculture Nutrition: Gut Health, Probiotics and Prebiotics*, 1st ed, Wiley-Blackwell, New York.
- Muthukumar P, Kandeepan C. 2015. Isolation, identification, and characterization of probiotic organisms from intestine of freshwater fishes. *Intl J Curr Microbiol Appl Sci* 4 (3): 2319-7706.
- Ringo E, Schillinger U, Holzapfel W. 2005. Antimicrobial activity of lactic acid bacteria isolated from aquatic animals and the use of lactic acid bacteria in aquaculture. In: Holzapfel WH, Naughton PJ. (eds.). *Microbial Ecology in Growing Animals*. Elsevier, New York.
- Roy R. 2013. *Budidaya Belut*. PT Media Agro Pustaka, Jakarta. [Indonesian]
- Tremonte P, Pannella G, Succi M, Tipaldi L, ISturchio M, Coppola R, Luongo D, Sorrentino E. 2017. Antimicrobial activity of *Lactobacillus plantarum* strains isolated from different environments: a preliminary study. *Intl Food Res J* 24 (2): 852-859.
- Verschuere L, Rombaut G, Sorgeloos P, Verstraete W. 2000. Probiotic bacteria as biological control agents in aquaculture. *Microbiol Mol Biol Rev* 64 (4): 655-671.