

Plant diversity in natural forest of Guilan Rural Heritage Museum in Iran

ROYA ABEDI¹, HASSAN POURBABAEI²

Department of Forestry, Faculty of Natural Resources, Somehsara, University of Guilan, Islamic Republic of Iran. P.O. Box 1144, Tel.: +98-182-3220895, Fax.: +98-182-3223600, email: ¹Abedi.Roya@yahoo.com, ²H_pourbabaei@guilan.ac.ir

Manuscript received: 31 August 2010. Revision accepted: 29 October 2010.

ABSTRACT

Abedi R, Pourbabaei H (2010) Plant diversity in natural forest of Guilan Rural Heritage Museum in Iran. *Biodiversitas* 11: 182-186. The aim of this study was to determine plant species diversity in Guilan Rural Heritage Museum in Iran. Eighty nine sampling plots were sampled based on systematic random method. Data analysis was carried out using diversity indices of richness, diversity (Shannon-Wiener, Simpson, Mc Arthur's (N_1) and Hill's (N_2) and Smith and Wilson's evenness index (E_{var}). Results indicated that Rosaceae and Labiatae families have the highest number of species. *Quercus castaneifolia* and *Ruscus hyrcanus* were the most dominant woody plants for class of tree and shrub, respectively. *Carex divolsa* and *Viola odorata* were dominant herbaceous species. Herbaceous layer had the highest richness, evenness and diversity. Mc Arthur's N_1 index had the highest value among diversity indices.

Key words: diversity, richness, evenness, Rural Heritage Museum, Guilan.

INTRODUCTION

Deforestation is one of the primary causes of biodiversity loss. Forests represent about 30% of terrestrial habitats, and support an exceptional number of species. Forests also provide economically important products and services. Small, isolated forest fragments are typically less able to provide these goods and services, or support a full complement of native species (Mayer and Tikka 2006). Natural forests decline in both extent and quality worldwide; there is an increasing recognition of the biodiversity conservation value of production landscapes (Le Brocque et al. 2009). Maintenance of biodiversity has been recognized as an important component of sustainable development and protection of native forests is a major means of biodiversity conservation (Muller et al. 2006). Efficiency of management and maintain of endangered species of a region could be evaluated when we have entire consciousness about biodiversity (Asri 2008).

Plant biodiversity consists of diversity into plant population structure, distribution, composition and abundance patterns. It is used as an index for comparison of forest ecosystems conditions (Pourbabaei 2001). Biological diversity has an indispensable value to society in that it (i) serves as a reservoir of genetic material that enhance productivity and stress tolerance of domesticated species and a source of new medicine, energy and industrial feed stock, (ii) provides ecological services such as amelioration of climate, water purification, soil stabilization and flood control and (iii) provides animals and natural landscape which have an overall benefit on human health and well-being through various forms of outdoor recreation (Brockway 1998).

Biological diversity is the richness and evenness of species amongst and within living organisms and ecological complexes (Polyakov et al. 2008). Biodiversity is mostly studied in species level. There are different indices to measure biodiversity. The most commonly considered facet of biodiversity is species richness. Evenness is another important factor of biodiversity. (Kharkwal et al. 2004). Evenness has been considered as a fundamental fact in habitats with more than one species (Hashemi 2010). Nowadays, numerous efforts to incorporate biodiversity into forests management and planning are encouraging (Brockway 1998).

Many studies have been carried out on plant biodiversity indices in Iran and around the world. Gholami et al. (2007) compared plant diversity, richness and evenness indices around protected area of the Bazangan Lake in Khorasan province, northeast of Iran. They indicated the highest value in Shannon-Wiener index. Ravanbakhsh et al. (2007) studied under-storey and over-storey plant biodiversity in Gisoom reserved forest in Guilan province, north of Iran and they showed that under-storey vegetation was disturbed and affected by human impacts. Abasi et al. (2009) investigated the effects of conservation on woody species diversity in protected regions of Oshtorankoo in Lorestan province, west of Iran. They expressed that trees and shrubs living in the protected regions species have significantly higher diversity, richness, evenness and better living condition than they living in non-protected region. Comparison of species richness and Hill's diversity indices showed that total species richness was higher in natural stands. Also, more fertile sites have significantly higher values of Hill's diversity index in mature stands of spruce plantation and natural stands in Southeaster New Brunswick, Canada

(Roberts 2002). Measurement of Shannon-Wiener and evenness indices on *Pinus massoniana* communities in Conservation project of plant biodiversity in Yangtze Three Gorges reservoir area, China showed that biodiversity of shrubs layer was the highest, followed by grass layer and the middle, while tree layer was the lowest (Tian et al. 2007).

Main objective of this study was to quantitatively analyze the biodiversity of vegetation cover in tree, shrub, herbaceous and regeneration layers in Guilan Rural Heritage Museum, Iran.

MATERIALS AND METHODS

Study area

The study was carried out at the Guilan Rural Heritage Museum with approximately 260 ha in extent that is located in Saravan Forest Park in the north of Iran (37°6 to 37°8 N latitude and 49°37 to 49°39 E longitude). The altitude ranges from 60 to 120 m asl. The climate is humid and very humid with cool winter according to Emberger climate classification. Mean annual temperature is 16.33°C and annual precipitation is 1366.64 mm. Maximum and minimum temperature is 27.8°C in August and 4.1°C in February, respectively (data obtained from 1985 to 2005, <http://www.weather.ir>) (Figure 1). This area is located at about 15 km far away from Rasht, the capital city of Guilan province, Iran.

Field sampling

Sampling procedure was the systematic random method. In this method, the sampling network size was 100×200 m. The distances between sampling strips were 200 m and the distances between circular plots on strips were 100 m. Then, start point was randomly selected and the sampling network was systematically located on the map (Poorbabaei et al. 2008; Pourmajidian et al. 2009; Poorbabaei and Poorrostam 2009; Shafiei et al. 2010). Totally, 89 sampling plots were taken. Data was collected to the class of tree (< 10 cm dbh) and shrub layers (number

of individual) in 1000 m² circular plots (Zobeiry 2005; Pourmajidian et al. 2009; Shafiei et al. 2010). In the center of these plots, the cover percentage of herbaceous species, including herbs, ferns and mosses, was estimated using Domin criterion by minimal area method with 32 m² areas. Number of regeneration in two classes include, sapling (< 1.30 m height and 10 cm dbh) and seedling (<1.30 m height) were sampled in 100 m² circular plots. Plant specimens were collected and stored in the Herbarium of Department of Forestry in Faculty of Natural Resources at University of Guilan.

Data analysis

Data analysis was carried out using diversity indices of Shannon-Wiener, Simpson, Mc Arthur's N₁, Hill's N₂, and Smith and Wilson's evenness (E_{var}). The diversity indices were calculated separately to different life forms: trees, shrubs, saplings, seedlings and herbs. Indices were used as following (Pitkanen 1998; Krebs 2001; Nagendra 2002; Nangendo et al. 2002; Small and McCarthy 2005; Lamb et al. 2009; Hashemi 2010):

Shannon-Wiener' H':
$$H' = -\sum_{i=1}^S p_i \ln p_i$$

Simpson' 1-D:
$$1 - D = 1 - \sum_{i=1}^S p_i^2$$

Mc Arthur's N₁:
$$N_1 = 2^{H'}$$

Hill's N₂:
$$N_2 = \frac{1}{\sum_{i=1}^S p_i^2}$$

S: the total number of species in the sample.

P_i: the proportion of individuals in the ith species (P_i= n_i/N, n_i is the number of individuals in the ith species and N is the total number of individuals)

Smith and Wilson's evenness (E_{var}) (Krebs 2001; Gosselin 2006):




Figure 1. Location of the study area: Guilan Rural Heritage Museum, Guilan Province, Islamic Republic of Iran.

$$E_{var} = 1 - \left[\frac{2}{\pi \arctan \left\{ \frac{\sum_{i=1}^S (\log_e(n_i) - \sum_{j=1}^S \log_e(n_j) / S)^2 / S}{S} \right\}} \right]$$

Arctan: measured as an angle in radians

n_i : basal area for over-storey species and is coverage for the under-storey of the i th species in the sampling plot

n_j : basal area for over-storey species and is coverage for the under-storey of the j th species in the sampling plot

S: total number of species in the entire sample

Numbers of species per plot was taken as a measure of species richness (S) (Timilsina et al. 2007). Data were calculated in Ecological Methodology software (Krebs 2001).

Species importance value (SIV) was calculated for all species by summing relative frequency, relative density and relative dominance values for woody species and summed relative frequency and relative dominance for herbaceous species. SIV was used to identify dominance species in the study area. The following formulas were used for each calculation (Maingi and Marsh 2006; Adam et al. 2007):

$$\text{Relative frequency} = \frac{\text{Number of plots that contain a species}}{\text{Number of all plots}} \times 100$$

$$\text{Relative density} = \frac{\text{Number of a species in all plots}}{\text{Total Number of species in all plots}} \times 100$$

$$\text{Relative dominance} = \frac{\text{Total basal area of a species in all plots}}{\text{Total basal area of all species in all plots}} \times 100$$

RESULTS AND DISCUSSION

Results

Totally, 75 plant species including 73 native species belong to 43 families and 72 genera were recorded throughout the study area (Table 1). Rosaceae and Labiatae families had the highest number of species (Figure 2). *Quercus castaneifolia* (227.24%) and *Ruscus hyrcanus* (126.60%) had the highest value of SIV and were the most dominant woody species to tree and shrub layers, respectively. *Carex divulsa* (76.47%) and *Viola odorata* (75.80%) were the most dominant herbaceous species. Mean richness of species (S) was higher in herbaceous layer and followed by regeneration, tree and shrub layers (Figure 3). Herbaceous layer had the highest amount of Smith and Wilson's evenness index (E_{var}) and followed by shrubs, regeneration and tree layers (Figure 4). Diversity indices of Simpson (1-D), Hill's N_2 , Shannon-Wiener (H) and Mc Arthur's N_1 in tree, shrub, herbaceous and regeneration layers were showed in Figures 5. Mc Arthur's N_1 was higher diversity index and followed by Hill's N_2 , Shannon-Wiener and Simpson in all layers of vegetation cover in the study area.

Table 1. List of plant species in Guilan Rural Heritage Museum.

Scientific Name	Family
<i>Acalypha australis</i> L.	Euphorbiaceae
<i>Acer insigne</i> Boiss.	Aceraceae
<i>Albizia julibrissin</i> Durazz.	Mimosaceae
<i>Alisma plantago-aquatica</i> L.	Alismataceae
<i>Alnus subcordata</i> C. A. Mey.	Betulaceae
<i>Artemisia annua</i> L.	Asteraceae
<i>Asplenium adiantum-nigrum</i> L.	Aspleniaceae
<i>Athyrium filix-femina</i> (L.) Roth.	Athyriaceae
<i>Azolla filiculoides</i> Lam.	Salviniaceae
<i>Brachythecium plumosum</i> (Hedw.) Schimp.	Brachytheciaceae
<i>Carex divulsa</i> Stokes.	Cyperaceae
<i>Carpinus betulus</i> L.	Betulaceae
<i>Chelidonium majus</i> L.	Papaveraceae
<i>Convolvulus betonicifolius</i> Mill.	Convolvulaceae
<i>Cirsium arvense</i> (L.) Scop.	Asteraceae
<i>Crataegus ambigua</i> M. B.	Rosaceae
<i>Cynodon dactylon</i> (L.) Pers.	Poaceae
<i>Cyperus rotundus</i> L.	Cyperaceae
<i>Danae racemosa</i> (L.) Moench.	Liliaceae
<i>Diospyros lotus</i> L.	Ebenaceae
<i>Epipactis latifolia</i> All.	Orchidaceae
<i>Erigeron canadensis</i> L.	Asteraceae
<i>Euphorbia amygdaloides</i> L.	Euphorbiaceae
<i>Ficus carica</i> L.	Moraceae
<i>Fragaria vesca</i> L.	Rosaceae
<i>Geum heterocarpum</i> Boiss.	Rosaceae
<i>Gleditsia caspica</i> Dest.	Caesalpinjiaceae
<i>Hedera helix</i> L.	Araliaceae
<i>Hedera pastuchovii</i> Woron. Ex Grossh.	Araliaceae
<i>Hypericum androsaemum</i> L.	Hypericaceae
<i>Hypericum perforatum</i> L.	Hypericaceae
<i>Ilex aquifolium</i> L.	Aquifoliaceae
<i>Juncus bufonius</i> L.	Juncaceae
<i>Juncus glaucus</i> Ehrh.	Juncaceae
<i>Lamium album</i> L.	Labiatae
<i>Lycopus europaeus</i> L.	Labiatae
<i>Mentha pulegium</i> L.	Labiatae
<i>Mespilus germanica</i> L.	Rosaceae
<i>Morus alba</i> L.	Moraceae
<i>Nepeta involucrate</i> (Bunge) Bornm.	Labiatae
<i>Oplismenus undulatifolius</i> (Ard.) P.	Poaceae
<i>Oxalis corniculata</i> L.	Oxalidaceae
<i>Palamocladium</i> sp.	Brachytheciaceae
<i>Parrotia persica</i> C. A. Mey.	Hamamelidaceae
<i>Periploca graeca</i> L.	Asclepiadaceae
<i>Phyla nodiflora</i> (L.) Greene.	Verbenaceae
<i>Phyllitis scolopendrium</i> (L.) Scop.	Aspleniaceae
<i>Plagiomnium cuspidatum</i>	Mniaceae
<i>Polygonum aviculare</i> L.	Polygonaceae
<i>Polypodium vulgare</i> L.	Polypodiaceae
<i>Populus caspica</i> Bornm.	Salicaceae
<i>Potentilla reptans</i> L.	Rosaceae
<i>Primula heterochroma</i> Starf.	Primulaceae
<i>Prunella vulgaris</i> L.	Labiatae
<i>Prunus domestica</i> L.	Rosaceae
<i>Pteridium aquilinum</i> L. Kuhn in Decken.	Hypolepidaceae
<i>Pteris cretica</i> L.	Pteridaceae
<i>Pterocarya fraxinifolia</i> (Lam.) Spach.	Juglandaceae
<i>Pyrus communis</i> L.	Rosaceae
<i>Quercus castaneifolia</i> C. A. Mey.	Fagaceae
<i>Rubus persicus</i> Boiss.	Rosaceae
<i>Ruscus hyrcanus</i> Woron.	Liliaceae
<i>Salix alba</i> L.	Salicaceae
<i>Salix aegyptiaca</i> L.	Salicaceae
<i>Sambucus ebulus</i> L.	Caprifoliaceae
<i>Scutellaria albida</i> L.	Labiatae
<i>Setaria glauca</i> (L.) P. Beauv.	Poaceae
<i>Smilax excelsa</i> L.	Liliaceae
<i>Solanum dulcamara</i> L.	Solanaceae
<i>Solidago virga-aurea</i> L.	Asteraceae
<i>Ulmus carpiniifolia</i> G. Suckow.	Ulmaceae
<i>Urtica dioica</i> L.	Urticaceae
<i>Viburnum lantana</i> L.	Caprifoliaceae
<i>Viola odorata</i> L.	Violaceae
<i>Zelkova carpinifolia</i> (Pall.) Diels	Ulmaceae




Figure 2. Number of species in each family of plants in Guilan Rural Heritage Museum, Iran.




Figure 3. Mean and standard error of richness in tree, shrub, Regeneration and herbaceous layers




Figure 4. Mean and standard error of evenness index in tree, shrub, regeneration and herbaceous layers




Figure 5. Mean diversity indices, richness, evenness and their standard errors in tree layer

Discussion

The assessment of biodiversity in forest has become an important issue for studying ecosystems and their conservation (Aubert et al. 2003). Biodiversity measurement is recognized as guidance for conservation plans in local scale. Species biodiversity is used greatly in vegetation studies, and environmental evaluation is one of the main criteria to determine ecosystems condition. So that, many research considered that high species diversity is equal to stability in ecological systems (Mirdavoodi and Zahedi Pour 2005).

The presence of 75 plant species in 260 ha area indicates considerable plant diversity in the study area. Our results showed that herb layer had the highest diversity indices (richness, diversity and evenness). The light penetration was high due to forest disturbance and decreasing canopy coverage, it led to increasing herbaceous species. Also, response of under-storey species to physiographical condition will be used as index of disturbances and changes in environmental and edaphical condition in sites (Mirzaei et al. 2008).

Many studies have emphasized the effects of slopes, aspect and elevation on plant diversity. It seems that high plant diversity in our study area is due to topographic and physiographic condition. This study area is flat in most parts (average of slope in most parts is 0-30 % and in some parts is 30-60%). High plant diversity is also due to fertility and humidity of sites. Steep slopes cause negative effects on site qualities by drainage of available water, soil erosion and decrease of soil nutrients (Sohrabi et al. 2007). In the other hand, low degree of slope and humidity were the most important factors of increasing diversity of herbaceous species (Gholami et al. 2007).

Studies showed that the highest values of richness and evenness occurred in the middle altitudes because of favorable temperature (Mirzaei et al. 2008). Thus, altitude ranging from 60 to 120 m asl. in our study area can be an appropriate condition for plant diversity. Northern slopes have more diversity due to must humidity (Kooch et al. 2009). Our study area was flat in many parts but have northern slopes in some parts. This can be evaluated as a reason of acceptable plant diversity.

Study on the effective environmental condition such as slope, aspects, altitudes and specially edaphic, climatic and anthropogenic factors on plant species diversity should be considered. Mc Arthur's N_1 is sensitive to the number of species (Pourbabaei and Ahani 2004). It was the highest value in herbaceous layer due to high number of herbaceous species in the study area. Simpson index is sensitive to the frequency of species (Pourbabaei and Ahani 2004). The highest Simpson index is herb layer, followed by shrub, regeneration and tree layers (shrubs have the lowest number of species but dispersed distributed in the study area). Roberts (2002) showed that fertile sites have the highest value of Hill's N_2 index and this index is significantly higher in natural stands. In our study, value of this index for herb layer was the highest and followed by shrubs, trees and regeneration layers. Comparison of this study area to Gisoom reserved forest in the west of Guilan province (both of them are coastal plain forests in north of Iran) showed that species richness was the same but biodiversity in Gisoom has better condition than in our study area. It seems that conservation strategies are important factor in plant diversity conservation in Gisoom (Ravanbakhsh et al. 2007).

CONCLUSION

Attempts to establish a diverse ecosystem with aesthetic values and help for plant conservation are necessary in Guilan Rural heritage Museum. Diversity is one of the main factors of sustainable forest management. Identifying plants species of a region and their biodiversity is very effective way to identify disturbance factors and develop recovery plans. It is also essential to maintain a high proportion of native woody species, create protection programs and preserve the area against human and livestock disturbances.

REFERENCES

- Abasi S, Hosseini SM, Pilehvar B, Zare H (2009) Effects of conservation on woody species diversity in Oshorankoo region, Lorestan. *Iranian J Forest* 1 (1): 1-10.
- Adam JH, Mahmud AM, Muslim NE (2007) Cluster analysis on floristic and forest structure of hilly lowland forest in Lak Kawi, Sabah of Malaysia. *Intl J Bot* 3 (4): 351-358.
- Asri Y (2008) Plant diversity in Mouthef Refuge, Iran. *Rostaniha* 9 (1): 25-37.
- Aubert M, Alard D, Bureau F (2003) Diversity of plant Assemblages in managed temperate forests: a case study in Normandy (France). *Forest Ecol Manag* 175: 321-337.
- Brockway DG (1998) Forest plant diversity at local and landscape scales in the Cascade Mountains of southwestern Washington. *Forest Ecol Manag* 109: 323-341.
- Gholami A, Ejtahadi H, Ghassemzadeh F, Ghorashi-Al-Hosseini J (2007) Study of Plant Biodiversity around Protected Area of the Bazangan Lake. *Iranian J Biol* 19 (4): 398-407.
- Gosselin F (2006) An assessment of the dependence of evenness indices on species richness. *J Theoretical Biol* 242: 591-597.
- Hashemi SA (2010) Evaluation plant species diversity and physiographical factors in natural broad leaf forest. *Amer J Environ Sci* 6 (1): 20-25.
- Kharkwal G, Mehrotra P, Rawat YS, Pangtey YPS (2004) Comparative study of herb layer diversity in pine forest stands at different altitudes of central Himalaya. *Appl Ecol Environ Res* 2 (2): 15-24.
- Kooch Y, Jalilvand H, Bahmnyar MA, Pormajidian MR (2009) Comparison of wood species diversity indices with respect to characteristics of natural lowland forest stands in Chalous. *Iranian J Biol* 22 (1): 183-192.
- Krebs CJ (2001) *Ecological methodology*. 2nd ed. Addison Wesley Longman. London.
- Lamb EG, Bayne E, Holloway G, Schieck J, Boutin S, Herbers J, Haughland DL (2009) Indices for monitoring biodiversity change: Are some more effective than others?. *Ecol indicators* 9: 432-444.
- Le Brocque AF, Goodhew KA, Zammit CA (2009) Over storey tree density and under storey re growth effects on plant composition, stand structure and floristic richness in grazed temperate woodlands in eastern Australia. *Agric, Ecosyst Environ* 129: 17-27.
- Maingi JK, Marsh SE (2006) Composition, structure, and regeneration patterns in a gallery forest along the Tana River near Bura, Kenya. *Forest Ecol Manag* 236: 211-228.
- Mayer AL, Tikka PM (2006) Biodiversity conservation incentive programs for privately owned forests. *Environ Sci Pol* 9: 614-625.
- Mirdavoodi HR, Zahedi Pour H (2005) Determination of suitable species diversity model for Meghan playa plant association and effect of some ecological factors on diversity change. *Pajuhesh & Sazandegi* 68: 56-65.
- Mirzaei J, Akbarinia M, Hosseini SM, Sohrabi H, Hosseinzade J (2008) Biodiversity of herbaceous species in related to physiographic factors in forest ecosystems in central Zagros. *Iranian J Biol* 20 (4): 375-382.
- Muller SW, Rusterholz HP, Baur B (2006) Effects of forestry practices on relict plant species on limestone cliffs in the northern Swiss Jura Mountains. *For Ecol Manag* 237: 227-236.
- Nagendra H (2002) Opposite trends in response for the Shannon and Simpson indices of landscape diversity. *Appl Geogr* 22: 175-186.
- Nangendo G, Stein A, Gelens M, de Gier A, Albricht R (2002) Quantifying differences in biodiversity between a tropical forest area and a grassland area subject to traditional burning. *For Ecol Manag* 164: 109-120.
- Pitkanen S (1998) The use of diversity indices to assess the diversity of vegetation in managed boreal forests. *For Ecol Manag* 112: 121-137.
- Polyakov M, Majumdar I, Teeter L (2008) Spatial and temporal analysis of the anthropogenic effects on local diversity of forest trees. *For Ecol Manag* 255: 1379-1387
- Poorbabaei H, Heydari M, Najafifar A (2008) The relationship between plant diversity and physiographic factors in ghalarang protected area, Ilam, western Iran; Proceedings of the Global Conference on Global Warming (GCGW) 6-10 July 2008. Istanbul. [Turkey].
- Poorbabaei H, Poor-rostam A (2009) The effect of shelterwood silvicultural method on the plant species diversity in a beech (*Fagus orientalis* Lipsky) forest in the north of Iran. *J For Sci* 55 (8): 387-394.
- Pourbabaei H (2001) Distribution of Box tree (*Buxus hyrcana* L.) sites and woody species diversity in Guilan province forests; proceeding of 7th scientific and research conference of Guilan University, Rasht, 5-6 March 2001. [Iran].
- Pourbabaei H, Ahani H (2004) Biodiversity of woody species in *Acer platanoides* sites in the shafaroud forests, Gilan (Iran). *Rostaniha* 5: 147-158.
- Pourmajidian MR, Malakshah NE, Fallah A, Parsakhoo A (2009) Evaluating the shelterwood harvesting system after 25 years in a beech (*Fagus orientalis* Lipsky) forest in Iran. *J Forest Sci* 55 (6): 270-278.
- Ravanbakhsh M, Ejtahadi H, Pourbabaei H, Ghoreschi-Al-Hoseini J (2007) Investigation on plants species diversity of Gisoom Taleh Reserve forest, Gilan province, Iran. *Iranian J Biol* 20 (3): 218-229.
- Roberts MR (2002) Effects of forest plantation management on herbaceous-layer composition and diversity. *Canadian J Bot* 80: 378-389.
- Shafiei AB, Akbarinia M, Jalali Gh, Hosseini M (2010) Forest fire effects in beech dominated mountain forest of Iran. *Forest Ecol Manag* 259: 2191-2196.
- Small CJ, McCarthy BC (2005) Relationship of understory diversity to soil nitrogen, topographic variation, and stand age in an eastern oak forest, USA. *Forest Ecol Manag* 217: 229-243.
- Sohrabi H, Akbarinia M, Hosseini SM (2007) Plants species diversity in ecosystem units in the Dehsorkh forests, Javanroud, Iran. *J Environ Stud* 33 (41): 61-68.
- Tian Z, Chenb W, Zhaob C, Chenc Y, Zheng B (2007) Plant biodiversity and its conservation strategy in the inundation and resettlement districts of the Yangtze Three Gorges, China. *Acta Oecologica Sinica* 27: 3110-3118.
- Timilsina N, Ross MS, Heinen JT (2007) A community analysis of sal (*Shorea robusta*) forests in the western Terai of Nepal. *Forest Ecol Manag* 241: 223-234.
- Zobeiry M (2005) *Forest inventory (measurement of tree and forest)*. 3rd ed. University of Tehran Press. Iran.
- www.weather.ir [Accessed 2008]

BIODIVERSITAS

Volume 11, Number 4, October 2010

Pages: 182-186

ISSN: 1412-033X (printed edition)

ISSN: 2085-4722 (electronic)

DOI: 10.13057/biodiv/d110403