

Asian Journal of Natural Product Biochemistry

| Asian J Nat Prod Biochem | vol. 19 | no. 2 | December 2021 |
| ISSN 2775-4189 | E-ISSN 2775-4197 |



Honey photo by Ingo Hoffmann

Asian Journal of Natural Product Biochemistry

| Asian J Nat Prod Biochem | vol. 19 | no. 2 | December 2021 |

Comparative wound healing potential of <i>Mitracarpus hirtus</i> ointment and honey in diabetic albino rats by collagen assessment BAWA INALEGWU, JACOB AONDONGUSHA JATO, OVYE AKYENGO, JOHN AKIGHIR	39-44
Anticancer mechanism of artonin E and related prenylated flavonoids from the medicinal plant <i>Artocarpus elasticus</i> CHRISTIAN BAILLY	45-57
Phytochemical analysis and in-vitro antimicrobial screening of the leaf extract of <i>Senna occidentalis</i> (Fabaceae) A.A. TAMASI, M.O. SHOGE, T.T. ADEGBOYEGA, E.C. CHUKWUMA	58-65
Detection of terpenoids and steroids in <i>Lindsaea obtusa</i> with thin layer chromatography YUSFIA URWATUL WUTSQA, SURATMAN, SITI LUSI ARUM SARI	66-69
Physicochemical composition of some new craft beers consumed in Maroua town from Far North region of Cameroon M. DIGUIR, A. LAYA, H. WANGSO, J.P. BAYANG, B.B. KOUBALA	70-80
Comparative determination of total antioxidant effects of ethanol extract of <i>Phyllanthus amarus</i> leaves OJOCHENEMI EJEH YAKUBU, MICHAEL SUNDAY ABU, JOHN AKIGHIR, JOB ITANYI ONUCHE, ALI ARABI	81-85



Asian Journal of Natural Product Biochemistry

| Asian J Nat Prod Biochem | vol. 19 | no. 2 | December 2021 |

ONLINE

<http://smujo.id/jnpb>

p-ISSN

2775-4189

e-ISSN

2775-4197

PUBLISHER

Society for Indonesian Biodiversity

CO-PUBLISHER

Universitas Sebelas Maret, Surakarta, Indonesia

OFFICE ADDRESS

Jl. Ir. Sutami 36A, Surakarta 57126, Central Java, Indonesia. Tel./fax. +62-271-663375, email: editors@smujo.id

PERIOD OF ISSUANCE

June, December

EDITOR-IN-CHIEF

Khalid Ahmad Khalid – National Research Center, Cairo, Egypt

ASSOCIATE EDITOR

Ahmad Dwi Setyawan – Universitas Sebelas Maret, Surakarta, Indonesia

EDITORIAL BOARD

Andri Frediansyah – Unit for Natural Product Technology, National Research and Innovation Agency, Indonesia

Asish Kumar Parida – CSIR-Central Salt & Marine Chemicals Research Institute, Bhavnagar, India

C.J. Sugiharjo – Universitas Gadjah Mada, Yogyakarta, Indonesia

Cedric B. Baker – Mercer University, Atlanta, USA

Chaerul – Research Center for Biology, National Research and Innovation Agency, Indonesia

Dayar Arbain – Universitas Andalas Padang, Indonesia

Enos Tangke Arung – Universitas Mulawarman, Samarinda, Indonesia

M. Ja'far Luthfi – UIN Sunan Kalijaga, Yogyakarta, Indonesia

Mallappa Kumara Swamy – Universiti Putra Malaysia, Malaysia

Supriyadi – Indonesian Spices and Medicinal Crops Research Institute, Bogor, Indonesia

Suranto – Universitas Sebelas Maret, Surakarta, Indonesia

Syamsul Arifin Achmad – Institut Teknologi Bandung, Indonesia

Taufik Hidayat – Institut Pertanian Bogor, Indonesia

List of reviewers: <https://smujo.id/jnpb/reviewers>



Society for Indonesia
Biodiversity



Universitas Sebelas Maret
Surakarta, Indonesia

GUIDANCE FOR AUTHORS

Aims and Scope *Asian Journal of Natural Product Biochemistry* (**Asian J Nat Prod Biochem**) (formerly *Biofarmasi*) encourages submission of manuscripts dealing with all important contributions to natural product chemistry, include bioactivity compounds of land and sea and of plants, microbes and animals; activities of antimicrobial, antiviral, antiparasite, anti-cancer, etc.; ethnopharmacology, ethnomedicines and medical application; plant-derived drugs and pharmaceutical formulations and evaluation; as well as biochemistry, chemistry, biotechnology, physiology, cell structures, pharmacology, toxicology, and chemical ecology of natural products.

Article types The journal seeks for: (i) **Research papers**, (ii) **Reviews**, and (iii) **Short communications**. Original full-length research manuscripts are limited to 8,000 words (including tables and figures) or proportional to articles in this publication number (beyond that, it should be with notice). Review articles are also limited to 8,000 words, while Short communications should be less than 2,500 words, except for pre-study (can be more).

Submission: The journal only accepts online submissions through the open journal system (<https://smujo.id/jnpb/about/submissions>) or, for login problems, email the editors at unsjournals@gmail.com (or editors@smujo.id). Submitted manuscripts should be the original works of the author(s). Please ensure that the manuscript is submitted using the template, which can be found at (<https://biodiversitas.mipa.uns.ac.id/D/template.doc>). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, and a paragraph describing the claimed novelty of the findings versus current knowledge. Please also provide a list of five potential reviewers in your cover letter. They should come from outside your institution and better from three different countries. Submission of a manuscript implies the submitted work has not been published (except as part of a thesis or report, or abstract) and is not being considered for publication elsewhere. When a group writes a manuscript, all authors should read and approve the final version of the submitted manuscript and its revision; and agree on the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis, drafting the manuscript, and correcting the revision. All authors must be responsible for the work's quality, accuracy, and ethics.

Ethics Author(s) must be obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. For the new invention, authors must manage its patent before publication.

Open Access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance Only articles written in US English are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double-blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. Manuscripts will be rejected if the content does not align with the journal scope, does not meet the standard quality, is in an inappropriate format, or contains complicated grammar, dishonesty (i.e., plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and significance. **Uncorrected proofs** will be sent to the corresponding author by system or email as .doc or .docx files for checking and correcting typographical errors. The corrected proofs should be returned in 7 days to avoid publication delays. The accepted papers will be published online in chronological order at any time but printed at the end of each month.

Free of charge This publication is dedicated entirely to the advancement of science and technology, therefore author(s), or author institution(s) are not subject to publication fees. **Reprints** The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by email and sending back the uncorrected proofs.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering sizes can be applied in presenting tables and figures (9 pt). Word processing program or additional software can be used; however, it must be PC compatible, use the template, and use Microsoft Word based (.doc or .rtf; not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italics, except in italicized sentences. Scientific names (genus, species, author) and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. The genus name can be shortened after the first mention, except in early sentences, or where this may generate confusion; name of the author can be eliminated after the first mention. For example, *Rhizopus oryzae* L. UICC 524 can be written hereinafter as *R. oryzae* UICC 524. Using trivial names should be avoided. **Biochemical and chemical nomenclature** should follow the order of the IUPAC-IUB. For DNA sequences, it is better to use Courier New font. Standard chemical abbreviations can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurements** should use IS denominations, and other systems should use equivalent values with the denomination of IS mentioned first. A dot should not follow abbreviations like g, mg, mL, etc. Minus index (m⁻², L⁻¹, h⁻¹) suggested being used, except in things like "per-plant" or "per-plot." **Mathematical equations** can be written down in one column with text; in that case, they can be written separately. **Numbers** one to ten are written in words, except if it relates to measurement, while values above them are written in number, except in early sentences. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent." Avoid expressing ideas with complicated sentences and verbiage/phrasing, and use efficient and effective sentences.

The title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written, especially for the first and the last name. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. We choose local names in Bahasa Indonesia for universities in Indonesia. The mention of "strata" program, should be avoided. Manuscript written by a group, author for correspondence along with address is required (marked with "*"). **The title page** (first page) should include title of the article, full name(s), institution(s) and address(es) of the author(s); the corresponding authors detailed postage and e-mail addresses (P), and phone (O) and fax numbers (O).

Abstract A concise abstract is required (about 200 words). The abstract should be informative and state briefly the aim of the research, the principal results and major conclusions. An abstract is often presented separately from the article, thus it must be able to stand alone (completely self-explanatory). References should not be cited, but if essential, then cite the author(s) and year(s). Abbreviations should be avoided, but if essential, they must be defined at their first mention. **Keywords** are about five words, covering scientific and local name (if any), research themes, and special methods used; and sorted from A to Z. **Abbreviations** (if any): All important abbreviations must be defined at their first mention there. **Running title** is about five words.

Introduction is about 600 words, covering the aims of the research and provide an adequate background, avoiding a detailed literature survey or a summary of the results. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for a manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgements** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of a maximum of three pages should be clearly presented. The title of a picture is written down below the picture, while the title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; the chart is preferred to use black and white images. The author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned the source. Author is suggested referring to Wikipedia for international boundaries and Google Earth for satellite imagery. If not specifically mentioned, it is assumed to refer to these sources. **There is no appendix**, all data or data analysis is incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Preferably 80% of it comes from scientific journals published in the last 10 years. In the text, give the author names followed by the year of publication and arrange from oldest to newest and from A to Z; in citing an article written by two authors, both of them should be mentioned; however, for three and more authors only the first author is mentioned followed by et al. For example, Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation should be avoided, as shown with the word "cit." Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in alphabetical order. Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). Please include DOI links for journal papers. The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158. DOI: 10.13057/biodiv/d070213.

The usage of "et al." in long author lists will also be accepted:

Smith J, Jones M Jr, Houghton L et al. 1999. Future of health insurance. *N Engl J Med* 325: 325-329. DOI: 10.1007/s002149800025.

Book:

Rai MK, Carpinella C. 2006. *Naturally Occurring Bioactive Compounds*. Elsevier, Amsterdam.

Chapter in the book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds.). *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.). *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. *Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon*. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from the internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. DOI: 10.1038/msb.2008.24. www.molecularsystembiology.com.

THIS PAGE INTENTIONALLY LEFT BLANK

Comparative wound healing potential of *Mitracarpus hirtus* ointment and honey in diabetic albino rats by collagen assessment

BAWA INALEGWU, JACOB AONDONGUSHA JATO*, OUYE AKYENGO, JOHN AKIGHIR

Department of Biochemistry, University of Agriculture Makurdi. P.M.B 2373, 970001 Makurdi, Nigeria.
Tel.: +234-8097037099, *email: jatojack@gmail.com

Manuscript received: 4 March 2021. Revision accepted: 11 June 2021.

Abstract. Inalegwu B, Jato JA, Akyengo O, Akighir J. 2021. Comparative wound healing potential of *Mitracarpus hirtus* ointment and honey in diabetic albino rats by collagen assessment. *Asian J Nat Prod Biochem* 19: 39-44. All humans will experience some type of wound in every lifetime. Most wounds heal quickly with little or no attention, but many people suffer from complex and/or persistent wounds, therefore posing a burden. This study was designed to assess the efficacy of *Mitracarpus hirtus* (L.) DC. ointment against honey in diabetic rats. To achieve this, percentage wound closure and collagen assessments were used to express treatment efficacy. Results show that on day 21, rats treated with *M. hirtus* ointment had the highest percentage closure (94.5%) while honey treated and non-treated recorded 90.0% and 83.3% respectively. Similarly, a significant difference ($p < 0.05$) was observed on day 21 in the total collagen deposited in wounds of diabetic rats (10.57 ± 0.7) and *M. hirtus* ointment treated wounds (11.77 ± 0.4) as compared with the non-treated diabetic rats. *M. hirtus* ointment was efficacious in healing wounds in diabetic rats and heals wound faster than honey and may hold potential for wound healing in diabetes mellitus sufferers. However, the wound healing mechanism of this ointment needs further investigation.

Keyword: Collagen, diabetic rats, honey, *Mitracarpus hirtus*, ointment, wound healing

Abbreviations: DM: Diabetes Mellitus; STZ: Streptozotocin; DMSO: dimethyl sulphoxide; DC: Diabetic Control; DHT: Diabetic honey treatment; DOT: Diabetic ointment treatment; NSSC: Neutral salt-soluble collagen; PSC: Pepsin soluble collagen; ASC: Acid soluble collagen

INTRODUCTION

Diabetes mellitus (DM) is a group of metabolic diseases characterized by hyperglycemia resulting from defects in secretion of insulin, insulin action, or both (ADA 2013). There are three broad categories: type 1, type 2, and gestational diabetes. Although there are other types that are specific to other individual causes (Masharani and German 2011). Symptoms of DM include hyperglycemia, polyuria, polydipsia, and weight loss; sometimes with polyphagia, blurred vision autonomic and peripheral neuropathy which causes poor healing wounds (IDF 2011; Yamany and Sayed 2012).

A wound is defined as a breakdown in the protective function of the skin; the loss of continuity of epithelium, with or without loss of underlying connective tissue following injury to the skin or underlying tissues/organs caused by surgery, a blow, a cut, chemicals, heat/cold, friction shear force, pressure or as a result of disease (Shankar et al. 2014). In the treatment of chronic and other wounds, the primary goals are rapid wound closure and a functional and aesthetically satisfactory scar (Shankar et al. 2014). The wound proceeds through a normal healing cascade of four programmed phases, viz: Homeostasis, inflammation, proliferation and remodelling with specific events to achieve scar formation.

According to Guo and DiPietro (2010), these wound

healing events involve: (i) rapid hemostasis; (ii) appropriate inflammation; (iii) mesenchymal cell differentiation, proliferation, and migration to the wound site; (iv) suitable angiogenesis; (v) prompt re-epithelialization (re-growth of epithelial tissue over the wound surface); and (vi) proper synthesis, cross-linking, and alignment of collagen to provide strength to the healing tissue.

Although the phases and events in wound healing are programmed, healing of wounds may delay. Microbial infections, exercise patterns, stress, nutritional status, and diseases such as diabetes have been identified as causes of delay (Damie 2011). Inflammatory and proliferative phases are the most affected stages during delay (Kurahashi and Fujii 2015). These two phases are characterized by neutrophils, macrophages, cytokines, keratinocytes, and fibroblasts, which ultimately result in collagen deposition and transformation of the extracellular matrix (Wong et al. 2013; Olczyk et al. 2014). Assessing the activities of these parameters is informative on progress of wound healing at a molecular level (Zaja-Milatovic and Richmond 2008; Kim et al. 2011; Upadhyay et al. 2014), but assessment of the end product (collagen) is more valuable, in that it is informative on the progress of healing and confirms the expected end products availability (Olczyk et al. 2014). This study used percentage wound closure, collagen deposition, and solubility in various solvents to compare the efficacy of a new herbal formulation (*M. hirtus* ointment) against honey in diabetic rats.

MATERIALS AND METHODS

Preparation of plant extract and ointment

The plant ointment was prepared according to the method detailed in Jato et al. (2018b). Briefly, harvested and identified *Mitracarpus hirtus* (L.) DC. (syn. *Mitracarpus villosus* (Sw.) Cham. & Schltld. ex DC.) plants were washed and air-dried for 15 days. Dried leaves were ground with a Linsan® blending machine (Lin 319) at 1000 RPM for 10 minutes at 25 °C and methanolic extraction achieved at 250 atm in a rotary evaporator (IKA® EW-28710-22). (50% w/w) *M. hirtus* ointment was prepared by heating 8 g of soft white paraffin in hot water bath for 30 min to melt. Then 8 g of the *M. hirtus* methanolic leaf extract was dissolved in 20 mL dimethyl sulphur oxide (DMSO) from JHD® Ltd and mixed thoroughly with the melted paraffin for 20 min. 500 g of the dried leaf were ground into fine powder with a Linsan® blending machine (Lin 319) at 1000 RPM for 10 minutes at 25°C.

Source and nature of honey used

500 mL of honey was purchased from a local honey farmer from Kwande Local Government area in Benue State and used for the entire research in its raw undiluted state, although no physicochemical analysis was done on the honey, the honey had a dark brown color and was very thick and sticky especially when allowed to make contact with air for a while.

Experimental animals

Care of animals was in line with National Institute of Health, guide for care and use of laboratory animals (Institute for Laboratory Animal Research 1996). Albino

rats purchased from Benue State University Animal House weighing 160-250 g were allowed to acclimatize for 7 days prior to the initiation of the experiment in plastic cages and under laboratory conditions (temperature 22 ± 2 °C and 12 hr light-dark cycle). Animals were fed with balanced diet purchased from UAC foods Nigeria Ltd and water *ad libitum*.

Experimental design

In this Randomized controlled trial (RCT) study design (3³), the rate of wound healing based on reduction in wound size and collagen quantification was studied for 21 days with 27 rats on days 7, 14 and 21. The rats were grouped into 3 (n=3) where group 1= control diabetic rats (non treated), group 2= diabetic rats (Honey treated), group =3 diabetic rats ointment treated.

Induction of diabetes mellitus and wound infliction

In fig 1 the flow chart for diabetes induction and wound healing study is presented. According to the method employed by Jato et al. (2018a), 12-hr fasted rats were administered single 65 mg/kg intraperitoneal injection of Streptozotocin (STZ) freshly prepared in 0.1 M sodium citrate buffer (pH 4.5). On day 8 after STZ injection, blood glucose measurement was performed on tail-vein blood with a glucometer (Accu-Chek® Aviva). Rats whose glucose tolerance test and fasting blood glucose levels exceeded 250 mg/dL (13.9 mmol/dL) were considered diabetic, consistent with the findings of (Mendes et al. 2012). The diabetic rats were then inflicted wounds same day to study wound healing by inflicting single full-thickness wounds (2 cm²) on the dorsum of the rats after disinfection of area with ethanol 90% (Wong et al. 2011).




Figure 1. Flow chart for diabetes induction and wound healing study

Physical examination by measurement of wound size

Wounds were observed postoperatively on days 7, 14 and 21 by measurement of wound size with a rule to determine wound area and then calculating percentage closure as in (eq. 1). Granulated tissues were collected same day for collagen quantification as a measure of wound healing in the diabetic rats as described in Desallais et al. (2014).

$$\% \text{ wound closure} = \frac{IWA - CWA}{IWA} \times 100 \quad \dots\dots\dots (\text{Eq.1})$$

Where: IWA: initial wound area; CWA: current wound area

Collagen assay of granulation tissue

The total collagen content in wounds and solubility of collagen in different solvents obtained from granulation tissues in wounds was evaluated by the hydroxyproline assay method of Bergman and Loxly employed by Desallais et al. (2014).

Estimation of total collagen content in granulation tissue

The collagen content in wounds was evaluated by the hydroxyproline assay method of Bergman and Loxly (Desallais et al. 2014). After digestion of punch biopsy specimens (3 mm diameter) in 6 M HCl for 3 hr at 120°C, evaporated to dryness and then made up with a known volume of water. The pH of the samples was adjusted to 7.0. Afterward, samples were mixed with 0.06 M chloramine T and incubated for 20 min at room temperature. Then, 3.15 M perchloric acid and 20% dimethylaminobenzaldehyde added, and samples were incubated for an additional 20 min at 60°C. The absorbance was determined at 557 nm with a spectrophotometer (UV-vis Jenway® 7305).

Extraction of neutral salt-soluble collagen

Granulation tissue was minced, homogenized in 10 vol of neutral salt solvent (1.0 M NaCl, 0.05 M Tris, pH 7.5) containing 20 mM EDTA and 2.0 mM N-ethyl maleimide and stirred for 24 hr. The suspension was then centrifuged at 35,000 g for 1 hr at 4 °C and the extraction was repeated with the pellet. The supernatants were pooled and an assay of hydroxyproline was done.

Extraction of acid-soluble collagen

The residue obtained from neutral salt soluble collagen was re-suspended in 10 vol of 0.5 M acetic acid and extracted for 24 hrs and centrifuged at 5000 g for 10 min. The pellet was then extracted with acetic acid, supernatants were pooled and an aliquot then used for the determination of hydroxyproline as in neutral salt soluble collagen.

Extraction of pepsin-soluble collagen

The residue obtained after acid extraction was re-suspended in 0.5 M acetic acid containing 100 mg pepsin per g of wet tissue. Digestion was carried out for 24 hr

followed by centrifugation at 5000 g for 10 min and re-extraction. Aliquots of pooled supernatant were then used for hydroxyproline measurement and aldehyde content in the collagen.

Statistical analysis

SPSS software version 21 was used for analysis of results and is expressed as mean ± S.D. Means were compared statistically using ANOVA and Tukey test as post hoc test. A statistically significant *p*-value <0.05 was considered.

RESULTS AND DISCUSSIONS

Wound healing in treated and non-treated STZ-induced diabetic albino rats as indicated by % wound closure

In Figure 2, wound healing results in albino rats treated as measured by percentage wound closure are presented. On day 7, the diabetic rats treated with *M. hirtus* ointment recorded the highest percentage closure of 14.50% while rats treated with honey recorded the least closure (3.3%). On day 14, the non-treated diabetic group (16.3%) whereby they showed lesser percentage of wound closure than the honey-treated rats (32.5%) while the *M. hirtus* ointment treated group maintained the lead with a percentage closure of 45.0%. On day 21, the diabetic rats treated with *M. hirtus* ointment maintained highest percentage closure of 94.5%. The change observed on day 14 was also maintained, an 83.3% and 90.0% wound closure was recorded for non-treated diabetic rats and diabetic rats treated with honey respectively. In all, wound closure and scar formation were recorded on days 16, 17 and 19 for diabetic ointment treatment, diabetic honey treatment, and diabetic control.

Wound healing in treated and non-treated STZ-induced diabetic albino rats as indicated by assessment of collagen formation

Table 1 presents results of total collagen deposited in wounds of treated and non-treated diabetic rats. They show that there is no statistically significant (*p* < 0.05) increase in the total collagen deposited in wounds of diabetic rats treated with *M. hirtus* ointment and with honey at day 7 post wound excision. A statistically significant difference (*p* < 0.05) was observed at day 14 in the deposition of total collagen for diabetic rats treated with honey (5.53 ± 0.2) and diabetic rats treated with *M. hirtus* ointment (5.67 ± 0.3) as compared with the non-treated diabetic rats (5.00 ± 0.1). Similarly, a significant (*p* < 0.05) increase was observed at day 21 in the deposition of total collagen in honey-treated wounds of diabetic rats (10.57 ± 0.7) and *M. hirtus* ointment treated wounds (11.77 ± 0.4) as compared with the non-treated diabetic rats. Thus all diabetic rats had increased total collagen formation with increased days of healing.




Figure 2. Wound healing in albino rats treated as measured by percentage wound closure

Table 1. Total collagen deposited at 7, 14, and 21 days post wound excision in STZ-induced diabetic rats in various treatment groups

Group	Duration of wounds		
	Day 7	Day 14	Day 21
DC	1.70 ± 0.4 ^a	5.00 ± 0.1 ^{a*}	9.83 ± 0.4 ^{a*}
DHT	1.93 ± 0.9 ^a	5.53 ± 0.2 ^b	10.57 ± 0.7 ^{ab}
DOT	2.40 ± 0.4 ^a	5.67 ± 0.3 ^b	11.77 ± 0.4 ^b

DC-Diabetic Control, DHT-Diabetic honey treatment, and DOT-Diabetic ointment treatment. Data are expressed as the mean ± SD from three animals in each group. Significant differences between the diabetic control and treated groups were assessed using Turkey test. *: significant difference ($p < 0.05$), ^a, ^b: values with same alphabets belongs to the same Homogeneous subset

Table 2. Wound healing in treated and non-treated STZ-induced diabetic albino rats as indicated by solubility of collagen in acid

Days	7	14	21
Group 1: DC (Mean ± SD)	524.00 ± 11.10 ^b	506.00 ± 9.00 ^b	492.67 ± 11.40 ^b
Group 2: DHT (Mean ± SD)	513.67 ± 5.70 ^{b*}	499.00 ± 7.00 ^b	473.67 ± 13.10 ^{b*}
Group 3: DOT (Mean ± SD)	393.67 ± 11.00 ^{a*}	356.67 ± 49.00 ^{a*}	342.00 ± 44.00 ^{a*}

Note: DC-Diabetic Control, DHT-Diabetic honey treatment, DOT-Diabetic ointment treatment. *: significantly different down the column at $p < 0.05$ when compared to the control, Values with the same alphabetical superscripts (^a, ^b) belong to the same Homogeneous subset as compared by Turkey post Hoc test

Table 3. Wound healing in treated and non-treated STZ-induced diabetic albino rats as indicated by solubility of collagen in neutral salt

Days	7	14	21
Group 1: DC (Mean ± SD)	116.67 ± 16.80 ^a	92.33 ± 10.80 ^a	78.67 ± 8.10 ^a
Group 2: DHT (Mean ± SD)	106.0 ± 4.00 ^a	85.33 ± 13.60 ^a	73.00 ± 9.60 ^a
Group 3: DOT (Mean ± SD)	101.33 ± 6.80 ^a	80.67 ± 13.60 ^a	68.33 ± 9.00 ^a

Note: DC-Diabetic Control, DHT-Diabetic honey treatment, DOT-Diabetic ointment treatment. *: significantly different down the column at $p < 0.05$ when compared to the control, Values with the same alphabetical superscripts (^a, ^b) belong to the same Homogeneous subset as compared by Turkey post Hoc test

Tables 2, 3 and 4 show solubility of collagen in various solvents namely: acid, neutral salt and pepsin respectively for the treated and non-treated groups. Results show that, no significant ($p < 0.05$) decrease was recorded in the pepsin soluble collagen (PSC) and neutral salt soluble collagen (NSSC) at day 7, 14 and 21 in the diabetic rats treated with honey, and diabetic rats treated with *M. hirtus* ointment when compared to the non-treated diabetic rats. Conversely, a significant ($p < 0.05$) decrease was observed in the solubility of acid-soluble collagen (ASC) on days 7, 14 and 21 in the diabetic rats treated with honey, and diabetic rats treated with *M. hirtus* ointment when compared to the non-treated diabetic rats. There was a decreasing trend in the solubility of ASC, NSSC and PSC in all the groups, indicating maturation and toughness of collagen deposits through the healing period. Therefore, in all the groups of diabetic rats there was a decrease in solubility with increase in days of experiment.

The *M. hirtus* ointment-treated diabetic rat wounds showed higher percentage closure than the honey-treated diabetic wounds in the albino rats while the non-treated diabetic rat wounds recorded the least percentage closure. The findings agree with that of Dwivedi and Chaudhary (2012) on amputures in diabetic wounds in rats while the findings of Barua et al. (2013) (89.76%) agree with findings using honey on day 21. Also, Sazegar et al. (2011) and Nisbet et al. (2010) reported a significant percentage closure in honey-treated wounds than the non-treated wounds.

Table 4. Wound healing in treated and non-treated STZ-induced diabetic albino rats as indicated by solubility of collagen in pepsin

Days	7	14	21
Group 1: DC (Mean ± SD)	2730.33 ± 333.00 ^a	2480.00 ± 385.10 ^a	2349.00 ± 201.80 ^a
Group 2: DHT (Mean ± SD)	2483.00 ± 404.50 ^a	2266.33 ± 185.00 ^a	2188.33 ± 424.90 ^a
Group 3: DOT (Mean ± SD)	2278.67 ± 294.40 ^a	2075.33 ± 168.20 ^a	1996.00 ± 314.00 ^a

Note: DC-Diabetic Control, DHT-Diabetic honey treatment, DOT-Diabetic ointment treatment. *: significantly different down the column at $p < 0.05$ when compared to the control, Values with the same alphabetical superscripts (^{a, b}) belong to the same Homogeneous subset as compared by Turkey post Hoc test

The higher percentage closure observed in the treated wounds of the diabetic rats can be attributed to increased deposition of total collagen and reduction in oxidative stress arising from the infliction of wounds to the rats. Close interaction of cells is necessary for synthesis and deposition of collagen facilitated by contracture of the skin tissue (Lembong et al. 2017). Cardiac glycosides have been implicated for contraction in tissues by inhibiting the Na^+ - K^+ ATPase pump in the sarcolemmal membrane of the myocyte and other cells (Cunningham et al. 2018). This inhibition causes intracellular accumulation of Na^+ , which makes the Na^+ - Ca^{2+} pump extrude less Ca^{2+} , causing Ca^{2+} to accumulate inside the cell resulting in increased force of contraction necessary for wound closure (Lembong et al. 2017). In addition to excellent water-absorbing quality, biodegradability, biocompatibility and outstanding film-forming properties, interactions of Ca^{2+} ions with the cells to facilitate contractions are the basis for using calcium alginate in wound dressing (Wang et al. 2015). Therefore, we submit that increased closures associated with the plant ointment may be the effect of cardiac glycosides in the plant on the wounds.

Collagen obtained from the treated and non-treated diabetic rats showed decreased solubility in the various solvents through the days of treatment. Also, the collagen of *M. hirtus* and honey exhibited lesser solubility in the acid, neutral salt buffer and pepsin solvents. This decrease in solubility was a result of collagen maturation (Weston and Althen 2002). The mature collagens have lesser crosslinks because the initial collagen produced is reducible and over time are replaced by mature, thermally stable and less soluble crosslinks. These mature crosslinks rather than total collagen are the key factors in collagen-related toughness (Weston and Althen 2002; Ponrasu and Suguna 2014). In an earlier submission, we reported that there was decreased antioxidant enzyme activity associated with increased wound healing. The development of mature tough collagen and the epidermal layer stratum corneum have been implicated for this negative correlation and the resultant challenges in transdermal drug delivery (Yang et al. 2018; Jato et al. 2021). Results obtained for the acid-soluble collagen were in agreement with the findings of Ponrasu and Suguna (2014) and Ponrasu et al. (2018) while that of the NSSC was in agreement with reports by Kirubanandan et al. (2013) and Ponrasu and Suguna (2014). PSC also recorded very high solubility more than the ASC and NSSC as observed by other studies (Ponrasu and Suguna 2014; Ponrasu et al. 2018; Kirubanandan et al. 2013). This is because the acid and the neutral salt only

carry out partial solubility, while the insoluble collagen is soluble in pepsin (Kirubanandan et al. 2013).

In conclusion, from the findings in this study, it was concluded that wounds of diabetic rats treated with *M. hirtus* ointment healed faster than honey. Thus diabetic rats treated with the ointment recorded the highest percentage of wound closure. Also, total collagen deposited in diabetic rats was highest in the ointment treated rats followed by honey treated rats while the non-treated was least and the solubility of collagen in the various solvents was decreased concomitantly with increase in healing and was lesser in the ointment treated diabetic rats, suggesting faster maturation of the deposited collagen. It is our view however that, the mechanism of action of the ointment be studied in addition to the identification and isolation of the bioactive compounds responsible for the efficacy of the ointment.

ACKNOWLEDGEMENTS

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors. However, the authors are thankful to Vincent Upev of Biochemistry Department, College of Veterinary Medicine, University Agriculture Makurdi for assistance with Streptozotocin. The authors declare that they have no conflict of interest. The authors also declare that all institutional and national guidelines for the care and use of laboratory animals were followed. This research was approved by the University's research and ethics committee with the approval letter ref No: FUAM/REC/19/1253612.

REFERENCES

- American Diabetes Association (ADA). 2013. Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care Suppl 1*: 36. DOI: 10.2337/dc13-S067
- Barua CC, Begum SS, Pathak DC, Bora RS. 2013 Wound healing activity of *Alternanthera brasiliana* Kuntze and its antioxidant profiles in experimentally induced diabetic rats. *J App Pharm Sci* 3 (10): 161-166. DOI: 10.7324/JAPS.2013.31029.
- Cunningham L, Levine G, Bozkurt B. 2018. Heart Failure: Long-term Management. In: *Cardiol Sec 5th ed.* 241-252. Elsevier, New York. DOI: 10.1016/B978-0-323-47870-0.00027-1.
- Damir A. 2011. Why Diabetic Foot Ulcers do not heal? *JIMSA* 24 (4): 205-206.
- Desallais L, Avouac J, Fréchet M, Elhai, M, Ratsimandresy R, Montes M, Mouhsine H, Do H, Zagury J, Allanore Y. 2014. Targeting IL-6 by

- both passive or active immunization strategies prevents bleomycin-induced skin fibrosis. *Arthritis Res Ther* 16 (4): 1-12. DOI: 10.1186/ar4672.
- Dwivedi VK, Chaudhary M. 2012. Comparative wound healing efficacy of ampicillin and becaplermin in diabetic rat. *Afr J Pharm Pharmacol* 6 (1): 883-892. DOI: 10.5897/AJPP11.748.
- Guo S, DiPietro LA. 2010. Factors affecting wound healing. *J Dent Res* 89 (3): 219-229. DOI: 10.1177/0022034509359125.
- Institute for Laboratory Animal Research. 1996. Guide for care and use of Laboratory Animal. National Academies Press [PubMed], Washington DC.
- International Diabetes Federation. 2011. The IDF Diabetes Atlas. 5th ed. Brussels: International Diabetes Federation. www.idf.org/idf-diabetes-atlas-fifth-edition
- Jato JA, Bawa I, Onyezili FN. 2018a. Diabetes induction with streptozotocin and insulin action on blood glucose levels in albino rats. *Intl J Mod Sci Technol* 3 (10): 208-212.
- Jato JA, Bawa I, Onyezili FN. 2018b. Phytochemical analysis of *Mitracarpus villosus*, and comparative toxicity of *Mitracarpus villosus* ointment and honey. *Intl J Mod Sci Technol* 3 (11): 220-227.
- Jato JA, Inalegwu B, Onyezili FN. 2021. Antioxidant enzymes in assessment of wound healing efficacy of *Mitracarpus villosus* ointment in diabetic state. *Chronic Dis J* 9 (1): 7-13. DOI: 10.22122/cdj.v9i1.581.
- Kim YS, Cho IH, Jeong MJ, Jeong SJ, Nah SY, Cho YS, Kim SH, Go A, Kim SE, Kang SS, Moon CJ. 2011. Therapeutic effect of total ginseng saponin on skin wound healing. *J Ginseng Res* 35 (3): 360-367. DOI: 10.5142/jgr.2011.35.3.360.
- Kirubanandan S, Swethkamal K, Renganathan S. 2013. Activities of triphala towards promoting the synthesis at wound site and inhibiting methicillin-resistant *Staphylococcus aureus* and its enzymes. *Intl J Pharm Pharmaceut Sci* 5 (2): 54-62.
- Kurahashi T, Fujii J. 2015. Roles of antioxidative enzymes in wound healing. *J Dev Biol* 3: 57-70. DOI: 10.3390/jdb3020057.
- Lembong, J, Sabass B, Stone HA. 2017. Calcium oscillations in wounded fibroblast monolayers are spatially regulated through substrate mechanics. *Phys Biol* 14 (4): 045006. DOI: 10.1101/116426.
- Masharani U, German MS. 2011. Pancreatic Hormones and Diabetes mellitus. In: Gardner DG, Shoback D (eds) Greenspan's basic & clinical endocrinology. 9th ed. McGraw-Hill Medical, New York.
- Mendes JJ, Leandro CI, Bonaparte DP, Pinto AL. 2012. A rat model of diabetic wound infection for the evaluation of topical antimicrobial therapies. *Comp Med* 62 (1): 37-48.
- Nisbet HO, Nisbet C, Yarim M, Guler, A, Ahmet O. 2010. Effects of three types of honey on cutaneous wound healing. *Wounds: Compendium Clin Res Pract* 22 (11): 275-283.
- Olczyk P, Mencner A, Komosinska-Vassev K. 2014. The role of the extracellular matrix components in cutaneous wound healing. *BioMed Res Int* 1-8. DOI: 10.1155/2014/747584.
- Ponrasu T, Madhukumar KN, Ganeshkumar M, Iyappan K, Sangeethapriya V, Gayathri VS, Suguna L. 2014. Efficacy of *Acorus calamus* on collagen maturation on full-thickness cutaneous wounds in rats. *Pharm Mag* 10 (2): S299-S305. DOI: 10.4103/0973-1296.133283.
- Ponrasu T, Suguna L. 2014. Efficacy of *Annona squamosa* L in the synthesis of glycosaminoglycans and collagen during wound repair in streptozotocin-induced diabetic rats. *BioMed Res Intl* (9): 1-10. DOI: 10.1155/2014/124352.
- Sazegar A, Reza A, Hosseini A, Behravan E. 2011. The effects of supplemental zinc and honey on wound healing in rats. *Iran J Bas Med Sci* 14 (4): 391-398.
- Shankar M, Ramesh B, Roopa KD, Niranjan BM. 2014. Wound healing and it's importance- a review. *Der Pharmacol Sin* 1 (1): 24-30.
- Upadhyay A, Chattopadhyay P, Goyary D, Mazumder PM, Veer V. 2014. *Ixora coccinea* enhances cutaneous wound healing by upregulating the expression of collagen and basic fibroblast growth factor. *Intl Scholarly Res Notices*. DOI: 10.1155/2014/751824.
- Wang T Gu Q, Zhao J, Mei J, Shao M, Pan Y Zhang J, Wu H, Zhang Z, Liu F. 2015. Calcium alginate enhances wound healing by up-regulating the ratio of collagen types I/III in diabetic rats. *Intl J Clin Exp Pathol* 8 (6): 6636-6645.
- Weston AR, Althen TG. 2002. Review: The role of collagen in meat tenderness. *Professional Anim Sci* 18: 107-111. DOI: 10.15232/S1080-7446(15)31497-2.
- Wong VW, Geoffrey C, Gurtner GC, Longaker MT. 2013. Wound healing: a paradigm for regeneration. *Mayo Clin Proc* 88 (9): 1022-1031. DOI: 10.1016/j.mayocp.2013.04.012.
- Wong VW, Sorkin M, Glotzbach JP, Longaker MT, Gurtner GC. 2011. Surgical approaches to create murine models of human wound healing. *J Biomed Biotechnol* 2011: 969618. DOI: 10.1155/2011/969618
- Yamany AA, Sayed HM. 2012. Effect of low-level laser therapy on neurovascular function of diabetic peripheral neuropathy. *J Adv Res* 3: 21-28. DOI: 10.1016/j.jare.2011.02.009.
- Yang J, Chen Z, Ye R, Li J, Lin Y, Gao J, Ren L, Liu B, Jiang L. 2018. Touch-actuated microneedle array patch for closed loop transdermal drug delivery. *Drug Delivery* 25 (1): 1728-39. DOI: 10.1080/10717544.2018.1507060.
- Zaja-Milatovic S, Richmond A. 2008. CXC chemokines and their receptors: A case for a significant biological role in cutaneous wound healing. *Histol Histopathol* 23 (11): 1399-1407. DOI: 10.14670/HH-23.1399.

Anticancer mechanism of artonin E and related prenylated flavonoids from the medicinal plant *Artocarpus elasticus*

CHRISTIAN BAILLY

OncoWitan, Scientific Consulting Office. Lille (Wasquehal), 59290, France. *email: christian.bailly@oncowitan.com

Manuscript received: 23 June 2021. Revision accepted: 25 July 2021.

Abstract. Bailly C. 2021. Anticancer mechanism of artonin E and related prenylated flavonoids from the medicinal plant *Artocarpus elasticus*. *Asian J Nat Prod Biochem* 19: 44-56. Plants of the *Artocarpus* genus are largely distributed throughout tropical Asia and Oceania. Species such as *A. altilis* (Parkinson) Fosberg and *A. heterophyllus* Lam. are popular trees known as breadfruit and jackfruit, respectively. They contain a large structural diversity of bioactive prenylated flavonoids. Here we have focused on the less known species *Artocarpus elasticus* Reinw. ex Blume which is well distributed in southeast Asia, and used in traditional medicine to treat dysentery, tuberculosis, and other diseases. Numerous prenylated flavonoids have been isolated from the leaves and bark of *A. elasticus*, such as artocarpesin, artocarpin, artelastin, and many others. They are endowed with antioxidant, anti-inflammatory, and anticancer properties. A focus is made of the subgroup of compounds designated artonins, with the derivative artonin E as a lead anticancer agent. Art-E has revealed marked anticancer effects *in vitro* and *in vivo*, after oral administration. The mechanism of action of Art-E is discussed, to highlight the structural and functional analogy between Art-E and the antitumor natural product morusin. Both compounds trigger TRAIL-mediated apoptosis of cancer cells. They can be considered further for the development of novel anticancer agents.

Keywords: *Artocarpus*, cancer therapeutics, natural products, phytotherapy, prenylated flavonoids

Abbreviations: iNOS, inducible nitric oxide synthase; ROS, reactive oxygen species

INTRODUCTION

The genus *Artocarpus* refers to a large group of plants belonging to the Moraceae family (Zerega et al. 2010). The internet database www.theplantlist.org has recorded 68 taxonomically accepted species, while about 90 additional plant names are now in synonymy or have even been moved as synonyms to other genera. The name *Artocarpus* derives from the Greek *artos* for “bread” and *karpos* for “fruit”. Some of the *Artocarpus* species are relatively well-known, such as *A. altilis* (Parkinson) Fosberg known as the breadfruit tree, and *A. heterophyllus* Lam. known as the jackfruit tree, which are two economically important *Artocarpus* species (Sikarwar et al. 2014; Mohammed Haleel et al. 2018; Sahu et al. 2019; Buddhisuharto et al. 2021). Their fruits are used for food production and several parts of these two trees including fruits, leaves, and barks have been used in traditional medicine in India and surrounding countries (Ranasinghe et al. 2019). Other *Artocarpus* species are used in traditional medicine, such as (i) decoctions prepared from the leaves of the tree *A. tonkinensis*, used in northern Vietnam to treat arthritis and backache (Adorisio et al. 2016), and (ii) extracts of *Artocarpus lacucha* (also known as Monkey jack) used in traditional Thai medicine as an anthelmintic agent (Aneklaphakij et al. 2020; Gupta et al. 2020), for examples. The ethnopharmacological use of *Artocarpus* species is widespread in subtropical and tropical regions of Asia where the trees are largely distributed. *Artocarpus* are believed to originate from the island of Borneo, from which species dispersed and diversified in several directions

(Williams et al. 2017). *Artocarpus*-based remedies are used to treat multiple diseases and conditions, including malarial fever, diarrhea, diabetes, parasitic infections, and different inflammatory diseases (Jagtap and Bapat 2010).

These plants contain a large diversity of bioactive compounds, including many prenylated flavonoids and cyclized derivatives which have revealed interesting antimicrobial, anti-inflammatory and anticancer properties (Hakim et al. 2006; Hari et al. 2014). One particular species, *Artocarpus elasticus* Reinw. ex Blume (Figure 1) has shown marked anticancer effects against different cancer cell lines and tumor models, attributed to the presence of prenylated flavonoids isolated from the root bark or other parts of the tree. The lead compound is artonin E (Art-E), thoroughly investigated as an antitumor agent. Over the past ten years, the mechanism of action of Art-E has been delineated and the activated signaling pathways at the origin of antitumor activity have been defined. These studies gave us the impetus to analyze in deep the mode of action of Art-E and structurally related compounds. The present review will address successively the plant *A. elasticus* and plant extracts, the phytochemical compounds isolated from the plant, the specific flavonoid group known as artonins, and the molecular targets and pathways activated by Art-E, responsible for its pharmacological properties. The role of TRAIL in the mode of action of Art-E and related compounds is discussed, opening the door to the identification of novel anticancer molecules. The purpose of the study was to promote the knowledge of artonins and the use of artonin E as a lead anti-cancer compound.




Figure 1. *Artocarpus elasticus* Reinw. ex Blume as a source of anticancer prenylated flavonoids. Different *Artocarpus* species are used in traditional medicine and as food products, such as *A. altilis* (Parkinson ex F.A.Zorn) Fosberg, *A. heterophyllus* Lam., *A. tonkinensis* A.Chev. ex Gagnep. and *A. lacucha* Buch.-Ham. Similarly, *A. elasticus* is also a source of bioactive mono-, bis- and tris-prenylated flavonoids, endowed with anticancer properties. *A. blumei* Trécul and *A. kunstleri* King are synonyms for *A. elasticus* Reinw. ex Blume (<http://www.theplantlist.org/>)

ARTOCARPUS ELASTICUS AND ITS MEDICINAL USES

The plant *Artocarpus elasticus* Reinw. ex Blume is native to Southeast Asia and can easily be found in countries like Myanmar, Thailand, Malaysia, Indonesia and the Philippines. In some countries like Malaysia and Indonesia, *A. elasticus* is cultivated to maintain its sustainability (Susiarti et al. 2020). It is an evergreen, robust and tall tree (up to 45-65 m). Various vernacular names are used locally, such as Benda (Javanese Indonesia), Terap nasi (Peninsular Malaysia), and others (Teo and Nasution 2003)

The wood of *A. elasticus* is sought as a building material. Upon wounding, it exudes a white and thick latex which is elastic or flexible, hence the name *elasticus*. The sticky latex is used to catch birds but also to treat dysentery. Different parts of the tree are used in traditional medicines, the latex for dysentery, the bark for female contraception, and the young leaves mixed with rice to treat tuberculosis. The flesh of the fruits is eaten raw or cooked. The seeds are consumed fried or roasted (Susiarti et al. 2020).

The bark and the leaves of the plant are often used to identify new phytochemicals. In addition, an extract of the stem bark of *A. elasticus* has been used as a reducing and stabilizing agent for the synthesis of silver nanoparticles (Abdullah et al. 2015). The same type of antibacterial silver nanoparticles have been obtained with seed extract of *A. hirsutus* (Shobana et al. 2020).

PRENYL FLAVONOIDS FROM ARTOCARPUS ELASTICUS

Artocarpus elasticus is a rich reservoir of bioactive natural products, notably prenyl flavonoids particularly abundant in the bark of the tree and the roots. There is a large structural diversity of prenylated flavonoids isolated from *A. elasticus*, with compounds bearing one, two or three prenyl groups, with a classical flavonoid skeleton or a bulkier pentacyclic core (Nomura et al. 1998; Kijjoa et al. 1998; Cidade et al. 2001). The main isolated compounds are shown in Figure 2, with the exception of the artonins discussed below. Most of these compounds can be found in other *Artocarpus* species. Here, we will discuss only those found in *A. elasticus*.

The tetracyclic compound artelastin, with three prenyl side chains, has been isolated in 1996 from the wood of *A. elasticus* together with artelastochromene, artelasticin, and artocarpesin (Kijjoa et al. 1996). Artelastin has revealed marked cytotoxic properties against different cancer cell lines *in vitro*. It was found to disturb the microtubule network of MCF7 breast cancer cells and to interfere with the cell cycle progression. Artelastin caused an accumulation in S phase due to drug-induced delay in DNA replication (Pedro et al. 2005). It is a potent inhibitor of both T- and B lymphocyte mitogen-induced proliferation and an inhibitor of cytokines production, such as interferon- γ , interleukins IL-2, -4 and -10, in stimulated splenocytes (Cerqueira et al. 2003). Artelastin displays marked antioxidant properties, inhibiting the production of reactive oxygen species (ROS) and the expression of the inducible nitric oxide synthase (iNOS) in lipopolysaccharide-stimulated macrophages (Cerqueira et al. 2008).




Figure 2. Chemical structures of prenylated flavonoids which have been isolated from *A. elasticus* Reinw. ex Blume

Artelasticin can be found in *A. elasticus* (Kijjoa et al. 1996), in *A. lanceifolius* (Syah et al. 2001) and other plants, such as the Cameroon medicinal plant *Dorstenia psilurus*. It was found to activate AMPK (AMP-activated protein kinase) and stimulate glucose uptake in rat skeletal muscles. Its marked blood glucose-lowering effect can be useful for the treatment of type-2 diabetes (Choumessi et al. 2019). Artocarpesin was found to be moderately cytotoxic toward a panel of cancer cell lines (IC_{50} values in the 60-100 μ M range) but its analog cycloartocarpesin is more potent (IC_{50} values in the 15-50 μ M range) (Kuetee et al. 2015). This compound can be found in many other plants. Like artelasticin, it presents marked anti-inflammatory and antioxidant properties (Fang et al. 2008). Its antioxidant activity contributes to the anti-browning effect observed with extracts of *A. heterophyllum* on fresh-cut apple slices. Remarkably, artocarpesin was found to exert a powerful inhibitory activity against mushroom tyrosinase ($IC_{50} = 0.52 \mu$ M) whereas the related product artocarpin was totally inactive against the enzyme (Zheng et al. 2008, 2009). Artocarpesin, found in the twigs and woods of *A. heterophyllum*, *A. incisus*, and *A. elasticus*, has been considered for the design of skin whitening agents (Arung et al. 2008, 2011). Artocarpin has been reported in a dozen of *Artocarpus* species and displays antioxidant, anti-inflammatory and anti-parasitic properties, as well as skin-whitening activities (Chan et al. 2018a; Morrison et al. 2021).

The bis-prenyl flavonoid cyclocommunin is known for a long time for its potent capacity to inhibit platelet aggregation induced by collagen or arachidonic acid (Lin et al. 1993). It was first isolated from *A. communis*, hence its

name cyclocommunin (Lin and Shieh, 1992), but it is also called isocyclomulberrin (Chen et al. 1993; Ma et al. 2010). Cyclocommunin is also an activator of protein kinase C (PKC) and a compound capable to elevate the level of intracellular calcium in rat neutrophils. As a result, cyclocommunin was found to stimulate respiratory burst in neutrophils (Wang et al. 1999), enhancing the production of superoxide anion, unlike artonin B which inhibited superoxide anion formation in these cells (Wei et al. 2005). Cyclocommunin displays modest anti-proliferative effects against hepatocellular carcinoma cells ($IC_{50} = 14-33 \mu$ M depending on cell line) whereas Art-A and Art-B were inactive against these HCC cell lines (Ma et al. 2010). Recently, it was reported that cyclocommunin displays a significant antimycobacterial activity against *Mycobacterium tuberculosis* (laboratory strain H37Ra) with a minimum inhibitory concentration (MIC) of 12.3 μ M (Boonyaketgson et al. 2020).

Artelastocarpin and carpelastofuran were isolated from the wood of *A. elasticus*. They both revealed cytotoxic properties toward cancer cells, with an efficacy like that of artelastin, artelasticin, and artelastochromene ($IC_{50} = 7-12 \mu$ M depending on cell lines) (Cidade et al. 2001). A series of compounds named artoindonesianins has been obtained from different *Artocarpus* species (Musthapa et al. 2010). Most of them have been isolated from species other than *A. elasticus*, such as the cytotoxic compounds artoindonesianins Z-4 and Z-5 found in the bark of *A. lanceifolius* (Musthapa et al. 2009a) and artoindonesianins A and B found in the roots of *A. champeden* (Hakim et al. 1999) and many other artoindonesianins, as indicated in Table 1.

Table 1. Artoindonesianin compounds isolated from *Artocarpus* species

Compound	<i>Artocarpus</i> species	References
Artoindonesianin A	<i>A. champeden</i>	Hakim et al. 1999
Artoindonesianin B	<i>A. champeden</i>	Hakim et al. 1999
Artoindonesianin C	<i>A. teysmanii</i> <i>A. rigidus</i> <i>A. kemando</i>	Makmur et al. 2000 Namdaung et al. 2006 ; Ee et al. 2011
Artoindonesianin D	<i>A. maingayi</i>	Hakim et al. 2000
Artoindonesianin E	<i>A. champeden</i>	Hakim et al. 2001
Artoindonesianin F	<i>A. heterophyllum</i>	Rao et al. 2010
Artoindonesianin G	<i>A. lanceifolius</i>	Syah et al. 2001
Artoindonesianin H	<i>A. lanceifolius</i>	Syah et al. 2001
Artoindonesianin I	<i>A. lanceifolius</i>	Syah et al. 2001
Artoindonesianin J	<i>A. bracteata</i>	Ersam et al. 2002
Artoindonesianin K	<i>A. champeden</i>	Achmad et al. 2005
Artoindonesianin L	<i>A. rotunda</i>	Suhartati et al. 2001
Artoindonesianin M	<i>A. champeden</i>	Syah et al. 2002b
Artoindonesianin N	<i>A. gomezianus</i>	Hakim et al. 2002a
Artoindonesianin O	<i>A. gomezianus</i>	Hakim et al. 2002a
Artoindonesianin P	<i>A. lanceifolius</i> <i>A. elasticus</i>	Hakim et al. 2002b Jenis et al. 2019
Artoindonesianin Q	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin R	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin S	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin T	<i>A. champeden</i>	Syah et al. 2002a
Artoindonesianin U	<i>A. champeden</i>	Syah et al. 2004
Artoindonesianin V	<i>A. champeden</i> <i>A. altilis</i>	Syah et al. 2004 Shamaun et al. 2010
Artoindonesianin W	<i>A. elasticus</i>	Jenis et al. 2019
Artoindonesianin X	<i>A. fretessi</i>	Soekamto et al. 2003
Artoindonesianin Y	<i>A. fretessi</i>	Soekamto et al. 2003
Artoindonesianin Z1	<i>A. lanceifolius</i> <i>A. anisophyllum</i>	Hakim et al. 2006 Noraini et al. 2013
Artoindonesianin Z2	<i>A. lanceifolius</i>	Hakim et al. 2006
Artoindonesianin Z3	<i>A. lanceifolius</i>	Hakim et al. 2006
Artoindonesianin Z4	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin Z5	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin Z4	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin Z5	<i>A. lanceifolius</i>	Musthapa et al. 2009b
Artoindonesianin A1	<i>A. champeden</i>	Syah et al. 2006b
Artoindonesianin A2	<i>A. champeden</i>	Syah et al. 2006a
Artoindonesianin A3	<i>A. champeden</i>	Syah et al. 2006a
Artoindonesianin B1	<i>A. altilis</i> <i>A. heterophyllum</i>	Syah et al. 2006b Lang et al. 2016
Artoindonesianin E1	<i>A. elasticus</i>	Musthapa et al. 2009a

For the record, the structure of artoindonesianin C, first isolated from *A. teysmanii* (Makmur et al. 2000), has been printed on a post stamp on the occasion of the international year of chemistry 2011 (Figure 3). The oxepinoflavone derivative artoindonesianin E1 was isolated from the bark of *A. elasticus*, together with the related compounds artocarpin, cycloartocarpin, and cudraflavones A and C (Figure 2). The derivative artoindonesianin A-3 was initially isolated from *A. champeden* (Syah et al. 2006a) and latter re-discovered from the bark of *A. elasticus* tree collected from Alor Island in Indonesia (Kuran and Ersam 2017). Other artoindonesianins have been found in *A. elasticus*, such as artoindonesianin P which was found to inhibit the enzyme α -glucosidase (IC_{50} = 25.4 μ M) but

with a reduced efficacy compared to Art-E and artobiloxanthone (IC_{50} = 16.2 and 8.6 μ M, respectively) (Jenis et al. 2019). Recently, artoindonesianins P and W were isolated from *A. elasticus* and found to inhibit human neutrophil elastase (IC_{50} = 28.7 and 11.2 μ M, respectively) whereas artobiloxanthone was slightly more active (IC_{50} = 9.8 μ M) (Ban et al. 2020).

Other prenylated flavonoids have been identified, such as artelastoheterol, artelasticinol, artelastoxanthone, artobiloxanthone, cycloartelastoxanthone, cycloartobiloxanthone, cycloartelastoxanthendiol, elastixanthone, and artonol A, all isolated from the root bark of *A. elasticus* (Figure 2). Artelastoxanthone has revealed modest antiproliferative activities against different cancer cell lines *in vitro* (Ko et al. 2005). Cycloartelastoxanthone and cycloartobiloxanthone and artobiloxanthone have shown protective effects on DNA damage caused by superoxide anion radicals $O_2^{\bullet-}$ (Lin et al. 2009). Elastixanthone and cycloartobiloxanthone have shown antimicrobial activities (Ramli et al. 2016). Artobiloxanthone and cycloartobiloxanthone have been found in many *Artocarpus* species, not limited to *A. elasticus*. The latter compound is interesting because it was found to potentially inhibit migration and invasion of H460 lung cancer cells, via inhibition of the phosphorylation of focal adhesion kinase (FAK) and the expression of cell division cycle 42 (CDC42) (Tungsukruthai et al. 2017). Cycloartobiloxanthone dose-dependently reduces proliferation of other lung cancer cell lines (H23, H292 and A549) and triggers caspase-dependent apoptosis with an efficacy comparable to that of the standard DNA-damaging anticancer drugs cisplatin and etoposide, at least *in vitro* (Losuwannarak et al. 2018). But the ROS scavenging activity of cycloartobiloxanthone is much weaker than that of artobiloxanthone and Art-E (Sritularak et al. 2010). A recent molecular docking analysis has predicted that artobiloxanthone can bind to the active site of the enzyme transglutaminase 2 (TG2), possibly inhibiting this enzyme considered as an anticancer target (Parvatikar and Madagi 2021).




Figure 3. The structure of artoindonesianin C, isolated from *A. teysmanii*. A post stamp with the structure of the compound was emitted to celebrate the international year of chemistry 2011 in Indonesia

Here we focused mostly on prenylated flavonoids isolated from *A. elasticus* but the whole *Artocarpus* genus the structural diversity is huge, as previously reported in specific review articles on *Artocarpus* species (Achmad et al. 2005; Hakim et al. 2006; Hakim 2010; Veitch and Grayer 2008). A recent analysis of a root extract of *A. heterophyllus* led to the identification of 47 prenylated flavonoids (Ye et al. 2019). There must be that many in the related species *A. elasticus*. An important group of prenylated flavonoids commonly found in *Artocarpus* species is called artonin, with the lead compound Art-E. Given the importance of the lead, this family is presented hereafter.

THE ARTONIN GROUP OF PRENYL FLAVONOIDS

There are 24 compounds named artonins A to Y (Table 2). The first two compounds, Art-A and -B, were isolated

in 1989 for the root bark of *A. heterophyllus* (Hano et al. 1989) followed by Art-C and -D one year later from the same plant (Hano et al. 1990a). They can be found in a few other *Artocarpus* species and display antioxidant properties (Ko et al. 2005). Art-B is an interesting compound, much more potent than Art-A at inhibiting the proliferation of nasopharyngeal cancer KB cells and the growth of the parasite *Trypanosoma brucei brucei*, *in vitro* (Bourjot et al. 2010). Art-B can reduce proliferation of different types of cancer cells, such as human CCRF-CEM leukemia cells, via the induction of mitochondria-dependent apoptosis (Lee et al. 2006). Art-A is less cytotoxic than Art-B, nevertheless, it is a more potent inhibitor of the papain-like cysteine protease cathepsin K (IC₅₀ = 1.9 and 9.0 μM, for Art-A and Art-B respectively) largely implicated in bone resorption (Zhai et al. 2017). Art-A also displays antimalarial activity, at least *in vitro* (Widyawaruyanti et al. 2007).

Table 2. Artonin compounds isolated from *Artocarpus* species

Compound	CID ^a	Formula (g/mol) ^b	<i>Artocarpus</i> species	References
Artonin A	14557102	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. heterophyllus</i> <i>A. styracifolius</i> <i>A. hypargyrea</i> <i>A. xanthocarpus</i> <i>A. champeden</i>	Hano et al. 1989 Bourjot et al. 2010 Qiao et al. 2011 Jin et al. 2015 Widyawaruyanti et al. 2007
Artonin B	11964501	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. heterophyllus</i> <i>A. champeden</i> <i>A. styracifolius</i>	Hano et al. 1989 Hakim et al. 1999s Bourjot et al. 2010
Artonin C	14681571	C ₄₀ H ₃₈ O ₁₀ (678.7)	<i>A. heterophyllus</i>	Hano et al. 1990a
Artonin D	14681573	C ₄₀ H ₃₆ O ₁₀ (676.7)	<i>A. heterophyllus</i>	Hano et al. 1990a
Artonin E	5481962	C ₂₅ H ₂₄ O ₇ (436.5)	<i>A. communis</i> <i>A. kemando</i> <i>A. lanceifolius</i> <i>A. chama</i> <i>A. nobilis</i> <i>A. lanceifolius</i> <i>A. gomezianus</i> <i>A. rotunda</i> <i>A. rigida</i> <i>A. rigidus</i>	Hano et al. 1990b Seo et al. 2003 Cao et al. 2003 Wang et al. 2004 Jayasinghe et al. 2008 Musthapa et al. 2009 Plaibua et al. 2013 Suhartati et al. 2001 Suhartati et al. 2018 Namdaung et al. 2006 Bourjot et al. 2010
Artonin F	14680593	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. integer</i> var. <i>silvestris</i> Corner <i>A. rigida</i>	Shaha et al. 2016 Hano et al. 1990
Artonin G	46887714	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigida</i>	Hano et al. 1990
Artonin H	21595104	C ₃₀ H ₃₄ O ₇ (506.6)	<i>A. rigida</i>	Hano et al. 1990
Artonin I	57335177	C ₄₀ H ₃₆ O ₁₁ (692.7)	<i>A. heterophyllus</i>	Hano et al. 1992
Artonin J	44258663	C ₂₅ H ₂₄ O ₇ (436.5)	<i>A. heterophyllus</i>	Aida et al. 1993
Artonin K	15340661	C ₂₁ H ₁₈ O ₇ (382.4)	<i>A. heterophyllus</i>	Aida et al. 1993
Artonin L	44258662	C ₂₂ H ₂₀ O ₇ (396.4)	<i>A. heterophyllus</i>	Aida et al. 1993
Artonin M	44258661	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. altilis</i> <i>A. rotunda</i>	Hano et al. 1993 Suhartati et al. 2001
Artonin N	44258669	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigida</i>	Hano et al. 1993
Artonin O	46887814	C ₃₀ H ₃₀ O ₇ (502.6)	<i>A. rigida</i> <i>A. rotunda</i>	Hano et al. 1993 Suhartati et al. 2001
Artonin P	44258658	C ₂₅ H ₂₀ O ₈ (448.4)	<i>A. rigida</i> <i>A. communis</i>	Hano et al. 1993 Chan et al. 2018b
Artonin Q	131753034	C ₃₁ H ₃₀ O ₈ (530.6)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin R	131753035	C ₃₁ H ₃₀ O ₁₀ (562.6)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin S	44258666	C ₂₆ H ₂₈ O ₇ (452.5)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin T	44258664	C ₂₆ H ₂₆ O ₇ (450.5)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin U	44258358	C ₂₁ H ₂₀ O ₅ (352.4)	<i>A. heterophyllus</i>	Aida et al. 1994
Artonin V	129687399	C ₂₅ H ₂₆ O ₇ (438.5)	<i>A. altilis</i>	Hano et al. 1994
Artonin X	(FDB021143)	C ₄₀ H ₃₈ O ₉ (662.7)	<i>A. heterophyllus</i>	Shinomiya et al. 1995
Artonin Y	15541482	C ₂₀ H ₁₈ O ₆ (354.4)	<i>A. heterophyllus</i>	Shinomiya et al. 2000

Note: ^aCompound Identity number (PubChem CID). ^bFormula and molecular weight.

The most important compound in the series is arguably Art-E (also known as 5'-hydroxymorusin), initially isolated from the stem bark of *A. communis* Forst., together with Art-F (Hano et al. 1990b). It was later found in several other *Artocarpus* species, including *A. elasticus* (Ramli et al. 2016). The compound, bearing an isoprenoid group at position C-3, was initially characterized as a potent and selective inhibitor of arachidonate 5-lipoxygenase ($IC_{50} = 0.36 \mu\text{M}$) expressed on cells involved in regulation of immune responses (Reddy et al. 1991). Art-E has revealed a very modest capacity to inhibit ADP-induced platelet aggregation *in vitro*, with an IC_{50} of $192 \mu\text{M}$ (Jantan et al. 2010). Later, Art-E was found to exert potent antiproliferative activities against different cancer cell lines, notably against breast adenocarcinoma MCF-7 and MDA-MB-231 cells ($ED_{50} = 2.2$ and $3.0 \mu\text{g/ml}$, respectively) (Wang et al. 2004). It is slightly less active against another type of cancer cells derived from solid tumors, but also exerts a potent cytotoxic action against P-388 leukemia cells (Musthapa et al. 2009b). Art-E is the lead product in the series, with interesting anticancer and antimicrobial properties (see below). The anticancer effects have been characterized using different experimental models *in vitro* and *in vivo*. In particular, the compound has revealed a robust dose-dependent effect in a 4T1 breast carcinoma xenograft model. The oral administration of Art-E at 25, 50 and 100 mg/kg reduced drastically the growth of tumor in mice and reduced the appearance of metastasis, without any apparent toxicity (Etti et al. 2017a). The compound is well tolerated; it can be administered orally for up to consecutive 10 days at 30 mg/kg, without any significant effect on cholesterol, creatinine and blood urea nitrogen levels (Fukai et al. 2003).

Modest antibacterial effects have been reported with Art-E against *Escherichia coli* and *Bacillus subtilis* (Suhartati et al. 2008), and with Art-O against *B. subtilis* (Suhartati et al. 2016). Mild antibacterial effects have been noted with Art-E against *Pseudomonas aeruginosa* strain PA01 (MIC = of $32 \mu\text{g/ml}$) whereas the compound was essentially inactive against the microorganisms *Providencia stuartii*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Salmonella typhi* and *Escherichia coli* (Kuete et al. 2011). Art-E showed little activity against *S. aureus* strain ATCC25922 (MIC = $256 \mu\text{g/ml}$) (Kuete et al. 2011) whereas a significant antibacterial effect has been reported with the strain ATCC 25923 (MIC = $3.9 \mu\text{g/ml}$) and the methicillin-resistant strain ATCC BAA-1720 (MIC = $13.3 \mu\text{g/ml}$) (Zajmi et al. 2015). A structural analysis using transmission electron microscopy revealed that Art-E disrupted the architecture of the bacterial cell, so as to facilitate the penetration of antibiotics (Zajmi et al. 2015). A similar type of effect on the bacterial cell wall has been reported with Art-I, despite it is structurally distinct from Art-E. Importantly, the membrane destabilizing effects of Art-E are reminiscent of those reported with morusin,

which is able to disrupt the cell membrane architecture and inhibiting the phosphatidic acid biosynthesis pathway of *S. aureus* (Pang et al. 2019). In many ways, morusin and Art-E behave similarly.

The flavonoid Art-F displays little cytotoxic effects against cancer cells but has shown antimycobacterial activity against *Mycobacterium tuberculosis* (MIC = $6.25 \mu\text{g/ml}$) (Namdaung et al. 2006). It has also revealed significant anti-plasmodial activity against the chloroquine-resistant strain FcB1, being only slightly less active than Art-B and more active than Art-A ($IC_{50} = 4.9, 1.56, 2.20 \mu\text{M}$ for Art-A, B and F, respectively) (Bourjot et al. 2010). Art-F, isolated from *A. integer* var. *silvestris* Corner, has revealed a weak capacity to inhibit 15-lipoxygenase (Shaha et al. 2016). Art-F can be found in different *Artocarpus* species, including a hybrid between *A. heterophyllus* (jackfruit) and *A. integer* (champedak) which has been specifically developed in Thailand to produce tastier and larger fruits, and for disease resistance (Panthong et al. 2013).

The isolation and structural characterization of several artonins have been reported but no specific effect has been pointed out, such as with Art-G and -H (Hano et al. 1990b), Art-J, -K and -L (Aida et al. 1993), and Art-M, -N, O, -P (Hano et al. 1993). Art-O, found in *A. rigida*, has shown a modest anti-proliferative action against HT-29 human colon cancer cell ($ED_{50} = 3.2 \mu\text{M}$) (Ren et al. 2010).

Art-I is much more interesting because the compound has revealed activities against multidrug-resistant (MDR) *Staphylococcus aureus* strains. It is a complex molecule (Figure 4) initially isolated from *A. heterophyllus* (Hano et al. 1992) but also from the leaves of *Morus mesozygia* (Fozing et al. 2012). Art-I has been found to inhibit the enzyme phosphodiesterase I *in vitro* ($IC_{50} = 15.4 \mu\text{M}$) (Fozing et al. 2012) and to function as an efflux pump inhibitor and a generator of ROS in *S. aureus*. Interestingly, the compound revealed an unanticipated capacity to promote considerably the activity of antibiotics (reducing the MIC 500- to 1000 fold), via Art-I-induced cell membrane damages (Farooq et al. 2014). It could be a useful natural product to treat multi-drug resistant *Staphylococcus* infections. The total synthesis of these complex molecules has been reported recently, opening the door to the design of simpler, more active analogs (Liu et al. 2020).

Art-V has been first isolated from *A. altalis* (Hano et al. 1994). This plant, known as paparagua tree, is largely used as a traditional medicine in Amazonian Ecuador and other countries in Oceania and South America. Many flavones and terpenoids have been isolated from this plant (Luzuriaga-Quichimbo et al. 2019). Art-X is a rare compound, isolated from *A. heterophyllus*, analogous to the diarylheptanoid kuwanon R which has been characterized as an inhibitor of protein tyrosine phosphatase 1B (PTP1B) (Shinomiya et al. 1995; Hoang et al. 2009).




Figure 4. Chemical structures of artonins

ANTICANCER MECHANISM AND MOLECULAR TARGETS OF ARTONIN-E

The mechanism of action of Art-E has been thoroughly investigated, at least at the cellular level. The compound triggers a down-regulation of the anti-apoptotic protein MCL1 (myeloid leukemia cell sequence-1) in human A549 and H292 non-small cell lung cancer cells and sensitizes them to anoikis (detachment-induced apoptosis) (Wongpankam et al. 2012). In these NSCLC cells, Art-E down-regulates several proteins involved in cell migration and invasion, such as FAK (focal adhesion kinase) and CDC42 (Cell division cycle-42). At non-cytotoxic low concentrations (0.05-05 $\mu\text{g/ml}$), the compound markedly reduced the migration of H460 NSCLC cells *in vitro* (Plaibua et al. 2013). But the main characteristic of Art-E is certainly its capacity to overcome resistance of cancer cells induced by the ligand TRAIL (tumor necrosis factor-related apoptosis-inducing ligand). In gastric cancer cells, the effect is coupled with an Art-E-induced up-regulation of the death receptor 5 (DR5) and tumor suppressor p53 (Toume et al. 2015). A similar enhancement of TRAIL-induced apoptosis by Art-E has been reported recently with LoVo colorectal cancer cells. The compound down-regulated DR5 while it up-regulated the other major regulator of TRAIL-induced apoptosis called cFLIP

(cellular FADD-like-IL-1beta-converting enzyme-inhibitory protein) on LoVo cells (Sophonnithprasert et al. 2019). Remarkably, Art-E can trigger cell cycle arrest (S phase) and a massive mitochondria-dependent apoptosis in SKOV-3 ovarian cancer cells (Rahman et al. 2016) and in HCT116 colon cancer cells, Art-E induced apoptosis through upregulation of p-ERK1/2 (Nimmuan-ngam et al. 2020).

The capacity of Art-E to overcome TRAIL-resistance in cancer cells echoes the effects reported many years ago with Art-B and three other prenylated flavonoids isolated from *A. champedan*. Art-B and the structural analog heterophyllin were found to potently overcome TRAIL resistance in human gastric adenocarcinoma cells and to enhance the expression of DR5 (Minakawa et al. 2014). Many other bioactive natural products have been found to suppress TRAIL resistance, or to enhance TRAIL-induced apoptosis via the death receptor pathway (Dai et al 2015; Ahmed and Ishibashi 2016; Shahwar et al. 2019) but the potency of Art-E and Art-B is noticeably high. A similar capacity to induce TRAIL sensitization by regulating DR5 has been observed with morusin in glioblastoma cells (Park et al. 2016), suggesting thus a class effect. TRAIL emerges as a master element of the antitumor action of Art-E and its analogs (Figure 5).




Figure 5. A schematic illustration of the anticancer mechanism of action common to morusine and artonin E (5-hydroxy-morusin). Both compounds trigger an up-regulation of the tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) and a down-regulation of the death receptor 5 (DR5). Upon trimerization of TRAIL and binding to DR5, the complex promotes the sequential recruitment of Fadd (Fas-associated protein with death domain) and caspase-8 (Casp-8) to form the death-inducing signaling complex (DISC). Processing of Bid (a BH3 domain-containing proapoptotic Bcl2 protein family member) by Casp-8 mediates mitochondrial damages. Activation of Casp-8 also directly induced Casp-3 activation, which concurs to the amplification of apoptosis.

Art-E has revealed marked activities against breast cancer cell lines. The compound inhibited the growth of estrogen receptor (ER) positive breast cancer cells MCF7 with an efficacy superior to that of the reference estrogen receptor modulator tamoxifen, at least *in vitro* ($IC_{50} = 3.8$ and $18.9 \mu\text{M}$ for Art-E and tamoxifen, at 72 hours, respectively). Molecular modeling predicted that Art-E can interact directly with the ligand-binding domain of hER α . The docking analysis indicated that Art-E can form more stable complexes with hER α compared to its analogs Art-U, -L, -S, and -T, and artoelastin (Etti et al. 2016). The analog isocycloporosin is predicted to exhibit an even better binding capacity to hER α than Art-E (Fitriah et al. 2018). Art-E induces caspase-dependent apoptosis of MCF7 cells, associated with elevated production of ROS and an up-regulation of the cyclin-dependent protein kinase inhibitor p21 (Etti et al. 2017b). Similar results have been reported using MDA-MB 231 triple-negative breast cancer cells: up-regulation of p21, G2/M cell cycle arrest, and caspase-dependent apoptosis (Etti et al. 2017c). Binding of Art-E to hER α is entirely plausible but this computational hypothesis has not been validated experimentally. Other targets have been proposed for Art-E, based on *in silico* modeling, such as the inflammatory cytokine-induced ubiquitin-like modifier FAT10 which is known to target hundreds of proteins for degradation by the 26S proteasome. A molecular modeling analysis has predicted that Art-E could bind and inhibit FAT10 implicated in hepatic carcinoma, but here again, there is no experimental validation of this hypothesis (Chaturvedi 2015). In brief, from a mechanistic point of view, Art-E can be considered as a TRAIL regulator in cancer cells, but more work is needed to better evidence the upstream molecular targets. Proteins such as hER α and FAT10 may provide a receptor for the compound but more works are needed to validate these molecular modeling proposals.

DISCUSSION

Artocarpus species have been largely investigated to characterize their pharmacological properties, in line with the traditional medicinal uses of these plants. Thus far, most phytochemical studies have been centered around *A. altilis* and *A. heterophyllus* (Nomura et al. 1998; Sikarwar et al. 2014; Mohammed Haleel et al. 2018; Mainasara and Abu Bakar 2019; Buddhisuharto et al. 2021). The species *A. elasticus* has been less investigated, although this bark-fiber producing plant is used traditionally to treat inflammatory conditions, but also to design crafts, clothes, ropes, and building materials (Veriyan et al. 2019). The phytochemical analysis indicates that like its congeners, *A. elasticus* is rich in prenylated flavones, in particular compounds with a C-3 prenyl side chain, sometimes cyclized into a 5 or 6 membered ring, and occasionally with an additional prenyl chain on ring A or B. These types of C-prenylated flavonoids are frequent in *Artocarpus* species (Šmejkal et al. 2014; Molčanová et al. 2019).

The present analysis sheds light on the artonin group of prenylated compounds and points out the prenylflavone Art-E as an interesting anticancer natural product. Art-E displays a range of pharmacological activities, chiefly antioxidant and anti-inflammatory effects, similar to those reported with the analogous isoprene flavonoid morusin (Choi et al. 2020). It is a potent anticancer agent, inspiring for the design and development of novel anticancer agents. The structural analogy between morusin and Art-E deserves further attention. Morusin has been much more studied than Art-E as an anticancer agent but also for its antiviral and antidiabetic effects (Choi et al. 2020; Kim et al. 2021). The *in vitro* and *in vivo* anticancer effects of morusin have been extensively characterized and the compound appeared promising against cancer stem cells (Zoofishan et al. 2018). We have a lot to learn from morusin to better understand the mode of action of Art-E.

Art-E and related prenylated flavonoids represent interesting anticancer agents, potentially useful to design novel bioactive compounds. Promising data have been reported with morusin-loaded nanoparticles for the treatment of glioblastoma (Agarwal et al. 2019; Zheng et al. 2021) and with artocarpin (Tzeng et al. 2016). A similar potency could be anticipated with Art-E using nanoparticle delivery systems. This compound deserves further attention.

CONCLUDING REMARK

In conclusion, this review exposes the potential health benefit of *Artocarpus elasticus* Reinw. ex Blume used as a traditional medicine in Asia and presents for the first time the complete series of artonin bioactive natural products. An extensive analysis of the scientific information led us to underline the value of the lead compound artonin E for the design of novel anticancer agents. The anticancer properties of Art-E have been well-established, demonstrating its capacity to trigger apoptosis of different types of cancer cells, but additional research efforts are required to better characterize its molecular targets. Another takeaway message is that Art-E can be considered as a morusin derivative, endowed with similar, if not superior, anticancer properties. This isoprene flavonoid warrants further studies.

FUNDING AND DECLARATION OF COMPETING INTEREST

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The author declares no conflict of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

REFERENCES

- Abdullah NI, Ahmad MB, Shameli K. 2015. Biosynthesis of silver nanoparticles using *Artocarpus elasticus* stem bark extract. *Chem Cent J* 9: 61.
- Achmad SA, Hakim EH, Juliawaty LD, Makmur L, Syah YM. 2005. Indonesian rainforest plants – Chemodiversity and bioactivity. *Malaysian J Sci* 24: 7-16.
- Adorisio S, Fierabracci A, Rossetto A, Muscari I, Nardicchi V, Liberati AM, Riccardi C, Van Sung T, Thuy TT, Delfino DV. 2016. Integration of Traditional and Western Medicine in Vietnamese Populations: A Review of Health Perceptions and Therapies. *Nat Prod Commun* 11: 1409-1416.
- Agarwal S, Mohamed MS, Mizuki T, Maekawa T, Sakthi Kumar D. 2019. Chlorotoxin modified morusin-PLGA nanoparticles for targeted glioblastoma therapy. *J Mater Chem B* 7: 5896-5919.
- Ahmed F, Ishibashi M. 2016. Bio-active Natural Products with TRAIL-Resistance Overcoming Activity. *Chem Pharm Bull (Tokyo)* 64: 119-127.
- Aida M, Shinomiya K, Hano Y, Nomura T. 1993. Artonins J, K, and L, three new isoprenylated flavones from the root bark of *Artocarpus heterophyllus* Lamk. *Heterocycles* 36: 575-583.
- Aida M, Shinomiya K, Matsuzawa K, Hano Y, Nomura T. 1994. Artonins Q, R, S, T, and U, Five New Isoprenylated Phenoles from the Bark of *Artocarpus heterophyllus* Lamk. *Heterocycles* 37: 847-858.
- Aneklaphakij C, Bunsupa S, Sirichamorn Y, Bongcheewin B, Satitpatipan V. 2020. Taxonomic Notes on the 'Mahat' (*Artocarpus lacucha* and *A. thailandicus*, Moraceae) Species Complex in Thailand. *Plants (Basel)* 9: 391.
- Arung ET, Muladi S, Sukaton E, Shimizu K, Kondo R. 2008. Artocarpin, A Promising Compound as Whitening Agent and Anti-skin Cancer. *J Tropic Wood Sci Technol* 6: 33-36
- Arung ET, Shimizu K, Kondo R. 2011. Artocarpus plants as a potential source of skin whitening agents. *Nat Prod Commun* 6: 1397-1402.
- Ban YJ, Baiseitova A, Nafiah MA, Kim JY, Park KH. 2020. Human neutrophil elastase inhibitory dihydrobenzoxanthones and alkylated flavones from the *Artocarpus elasticus* root barks. *Appl Biol Chem* 63: 63.
- Boonyaketgason S, Du Y, Valenciano Murillo AL, Cassera MB, Kingston DGI, Trisuwan K. 2020. Flavanones from the Twigs and Barks of *Artocarpus lakoocha* Having Antiplasmodial and Anti-TB Activities. *Chem Pharm Bull (Tokyo)* 68: 671-674.
- Bourjot M, Apel C, Martin MT, Grellier P, Nguyen VH, Guéritte F, Litaudon M. 2010. Antiplasmodial, antitrypanosomal, and cytotoxic activities of prenylated flavonoids isolated from the stem bark of *Artocarpus styracifolius*. *Planta Med* 76: 1600-1604.
- Buddhisuharto AK, Pramastya H, Insanu M, Fidriani I. 2021. An Updated Review of Phytochemical Compounds and Pharmacology Activities of *Artocarpus* genus. *Biointerface Res Applied Chem* 11: 14898-14905.
- Cao S, Butler MS, Buss AD. 2003. Flavonoids from *Artocarpus lanceifolius*. *Nat Prod Res* 17: 79-81.
- Cerqueira F, Cidade H, van Ufford L, Beukelman C, Kijjoa A, Nascimento MS. 2008. The natural prenylated flavone artelastin is an inhibitor of ROS and NO production. *Int Immunopharmacol* 8: 597-602.
- Cerqueira F, Cordeiro-da-Silva A, Araújo N, Cidade H, Kijjoa A, Nascimento MS. 2003. Inhibition of lymphocyte proliferation by prenylated flavones: artelastin as a potent inhibitor. *Life Sci* 73: 2321-2334.
- Chan EWC, Wong SK, Tangah J, Chan HT. 2018a. Chemistry and Pharmacology of Artocarpin: An Isoprenyl Flavone from *Artocarpus* Species. *Sys Rev Pharm* 9: 58-63
- Chan STS, Popplewell WL, Bokesch HR, McKee TC, Gustafson KR. 2018b. Five New Stilbenes from the Stem Bark of *Artocarpus communis*. *Nat Prod Sci* 24: 266-271.
- Chaturvedi I. 2015. A Molecular Docking study to find Natural Inhibitor Against FAT10 Protein for curing Hepatic Carcinoma. *J Sci* 1: 1-9.
- Chen CC, Huang YL, Ou JC, Lin CF, Pan TM. 1993. Three new prenylflavones from *Artocarpus altilis*. *J Nat Prod* 56: 1594-1597.
- Choi DW, Cho SW, Lee SG, Choi CY. 2020. The Beneficial Effects of Morusin, an Isoprene Flavonoid Isolated from the Root Bark of *Morus*. *Int J Mol Sci* 21: 6541.
- Choumessi AT, Johanns M, Beaufay C, Herent MF, Stroobant V, Vertommen D, Corbet C, Jacobs R, Herinckx G, Steinberg GR, Feron O, Quetin-Leclercq J, Rider MH. 2019. Two isoprenylated flavonoids from *Dorstenia psilurus* activate AMPK, stimulate glucose uptake, inhibit glucose production and lower glycemia. *Biochem J* 476: 3687-3704.
- Cidade HM, Nascimento MS, Pinto MM, Kijjoa A, Silva AM, Herz W. 2001. Artelastocarpin and carpelastofuran, two new flavones, and cytotoxicities of prenyl flavonoids from *Artocarpus elasticus* against three cancer cell lines. *Planta Med* 67: 867-870.
- Dai X, Zhang J, Arfuso F, Chinnathambi A, Zayed ME, Alharbi SA, Kumar AP, Ahn KS, Sethi G. 2015. Targeting TNF-related apoptosis-inducing ligand (TRAIL) receptor by natural products as a potential therapeutic approach for cancer therapy. *Exp Biol Med (Maywood)* 240: 760-773.
- Ee GC, Teo SH, Rahmani M, Lim CK, Lim YM, Go R. 2011. Artomandin, a new xanthone from *Artocarpus kemando* (Moraceae). *Nat Prod Res* 25: 995-1003.
- Ersam T, Achmad SA, Ghisalberti E, Hakim EH, Makmur L, Syah YM. 2002. A New Isoprenylated Chalcone, Artoindonesianin J, from the Root and Tree Bark of *Artocarpus bracteata*. *J Chem Res* 2002: 186-187.
- Etti I, Abdullah R, Hashim NM, Kadir A, Abdul AB, Etti C, Malami I, Waziri P, How CW. 2016. Artonin E and Structural Analogs from *Artocarpus* Species Abrogates Estrogen Receptor Signaling in Breast Cancer. *Molecules* 21: 839.
- Etti IC, Abdullah R, Hashim NM, Yeap SK, Imam MU, Ramli F, Malami I, Lam KL, Etti U, Waziri P, Rahman M. 2017c. The molecular mechanism of the anticancer effect of Artonin E in MDA-MB 231 triple-negative breast cancer cells. *PLoS One* 12: e0182357.
- Etti IC, Abdullah RB, Hashim NM, Arifah K, Yeap SK, Sani D, Ramli F, Malami I, Waziri P. 2017a. Reduction of breast tumor burden in mice by a prenylated flavonoid, Artonin E. *Australian Med J* 10: 681-693.
- Etti IC, Rasheed A, Hashim NM, Abdul AB, Kadir A, Yeap SK, Waziri P, Malami I, Lim KL, Etti CJ. 2017b. Artonin E induces p53-independent G1 cell cycle arrest and apoptosis through ROS-mediated mitochondrial pathway and living suppression in MCF-7 cells. *Drug Des Devel Ther* 11: 865-879.
- Fang SC, Hsu CL, Yen GC. 2008. Anti-inflammatory effects of phenolic compounds isolated from the fruits of *Artocarpus heterophyllus*. *J Agric Food Chem* 56: 4463-4468.
- Farooq S, Wahab AT, Fozing CD, Rahman AU, Choudhary MI. 2014. Artonin I inhibits multidrug resistance in *Staphylococcus aureus* and potentiates the action of inactive antibiotics in vitro. *J Appl Microbiol* 117: 996-1011.
- Fitriah A, Holil K, Syarifah U, Fitriyah F, Utomo DH. 2018. In silico Approach for Revealing the Anti-Breast Cancer and Estrogen Receptor Alpha Inhibitory Activity of *Artocarpus altilis*. *AIP Conference Proceedings* 2021: 070003.
- Fozing CD, Ali Z, Ngadjui BT, Choudhary MI, Kapche GD, Abegaz BM, Khan IA. 2012. Phosphodiesterase I-inhibiting Diels-Alder adducts from the leaves of *Morus mesozygia*. *Planta Med* 78: 154-159.
- Fukai T, Satoh K, Nomura T, Sakagami H. 2003. Antinephritis and radical scavenging activity of prenylflavonoids. *Fitoterapia* 74: 720-724.
- Gupta AK, Rather MA, Kumar Jha A, Shashank A, Singhal S, Sharma M, Pathak U, Sharma D, Mastinu A. 2020. *Artocarpus lakoocha* Roxb. and *Artocarpus heterophyllus* Lam. Flowers: New Sources of Bioactive Compounds. *Plants (Basel)* 9: 1329.
- Hakim A. Diversity of secondary metabolites from Genus *Artocarpus* (Moraceae). 2010. *Nusantara Biosci* 2: 146-156.
- Hakim EH, Achmad SA, Juliawaty LD, Makmur L, Syah YM, Aimi N, Kitajima M, Takayama H, Ghisalberti EL. 2006. Prenylated flavonoids and related compounds of the Indonesian *Artocarpus* (Moraceae). *J Nat Med* 60: 161-184.
- Hakim EH, Afrida A, Eliza E, Achmad SA, Aimi N, Kitajima N, Makmur L, Mujahidin D, Syah YM, Takayama H. 2000. Artoindonesianin-D: senyawa baru bioaktif turunan piranoflavon dan chaplasi tumbuhan *Artocarpus maingayi*. *Proc ITB* 32, 13-19.
- Hakim EH, Aripin A, Achmad SA, Aimi N, Kitajima N, Makmur L, Mujahidin D, Syah YM, Takayama H. 2001. Artoindonesianin-E suatu senyawa baru turunan flavanon dari tumbuhan *Artocarpus champeden*. *Proc ITB* 33: 69-73.
- Hakim EH, Asnizar, Yurnawilis, Aimi N, Kitajima M, Takayama H. 2002b. Artoindonesianin P, a new prenylated flavone with cytotoxic activity from *Artocarpus lanceifolius*. *Fitoterapia* 73: 668-673.
- Hakim EH, Fahriyati A, Kau MS, Achmad SA, Makmur L, Ghisalberti EL, Nomura T. 1999. Artoindonesianins A and B, two new prenylated

- flavones from the root of *Artocarpus champeden*. J Nat Prod 62: 613-615.
- Hakim EH, Ulinnuha UZ, Syah YM, Ghisalberti EL. 2002a. Artoindonesianins N and O, new prenylated stilbene and prenylated arylbenzofuran derivatives from *Artocarpus gomezianus*. Fitoterapia 73: 597-603.
- Hano Y, Aida M, Nomura T, Ueda S. 1992. A novel way of determining the structure of artonin I, an optically active Diels-Alder type adduct, with the aid of an enzyme system of *Morus alba* cell cultures. J Chem Soc. Chem Commun 17: 1177-1178.
- Hano Y, Aida M, Nomura T. 1990a. Two new natural Diels-Alder-type adducts from the root bark of *Artocarpus heterophyllum*. Heterocycles 53: 391-395.
- Hano Y, Aida M, Shiina M, Nomura T, Kawai T, Ohe H, Kagei K. 1989. Artonins A and B, Two New Prenylflavones from the Root Bark of *Artocarpus heterophyllum* Lamk. Heterocycles 29: 1447-1453.
- Hano Y, Inami R, Nomura T. 1994. Constituents of Moraceae plants. 20. a novel flavone, artonin V, from the root bark of *Artocarpus altilis*. J Chem Res Syn 9: 348-349.
- Hano Y, Inami R, Nomura T. 1990b. Components of the Bark of *Artocarpus rigida* Bl. 1. Structures of Two New Isoprenylated Flavones, Artonins G and H. Heterocycles 31: 2173-2179.
- Hano Y, Inami R, Nomura T. 1993. Components of the Bark of *Artocarpus rigida* Bl. 2. Structures of Four New Isoprenylated Flavone Derivatives Artonins M, N, O and P. Heterocycles 35: 1341-1350.
- Hano Y, Yamagami Y, Kobayashi M, Isohata R, Nomura T. 1990b. Artonins E and F, two new prenylflavones from the bark of *Artocarpus communis* Forst. Heterocycles 31: 877-882.
- Hari A, Revikumar KG, Divya D. 2014. *Artocarpus*: A review of its phytochemistry and pharmacology. J Pharma Search 9: 7-12.
- Hoang DM, Ngoc TM, Dat NT, Ha do T, Kim YH, Luong HV, Ahn JS, Bae K. 2009. Protein tyrosine phosphatase 1B inhibitors isolated from *Morus bombycis*. Bioorg Med Chem Lett 19: 6759-6761.
- Jagtap UB, Bapat VA. 2010. *Artocarpus*: a review of its traditional uses, phytochemistry and pharmacology. J Ethnopharmacol 129: 142-166.
- Jantan I, Mohd Yasin YH, Jamil S, Sirat H, Basar N. 2010. Effect of prenylated flavonoids and chalcones isolated from *Artocarpus* species on platelet aggregation in human whole blood. J Nat Med 64: 365-369.
- Jayasinghe UL, Samarakoon TB, Kumarihamy BM, Hara N, Fujimoto Y. 2008. Four new prenylated flavonoids and xanthenes from the root bark of *Artocarpus nobilis*. Fitoterapia 79: 37-41.
- Jenis J, Baiseitova A, Yoon SH, Park C, Kim JY, Li ZP, Lee KW, Park KH. 2019. Competitive α -glucosidase inhibitors, dihydrobenzoxanthenes, from the barks of *Artocarpus elasticus*. J Enzyme Inhib Med Chem 34: 1623-1632.
- Jin YJ, Lin CC, Lu TM, Li JH, Chen IS, Kuo YH, Ko HH. 2015. Chemical constituents derived from *Artocarpus xanthocarpus* as inhibitors of melanin biosynthesis. Phytochemistry 117: 424-435.
- Kijjoa A, Cidade H, Pinto M, Gonzalez MJTG, Anantchoke C, Gedris TE, Herz W. 1996. Prenylflavonoids from *Artocarpus elasticus*. Phytochemistry 43: 691-694.
- Kijjoa A, Cidade HM, Gonzalez MJTG. 1998. Further prenylflavonoids from *Artocarpus elasticus*. Phytochemistry 47: 875-878.
- Kim TI, Kwon EB, Oh YC, Go Y, Choi JG. 2021. Mori ramulus and its Major Component Morusin Inhibit Herpes Simplex Virus Type 1 Replication and the Virus-Induced Reactive Oxygen Species. Am J Chin Med 49: 163-179.
- Ko HH, Lu YH, Yang SZ, Won SJ, Lin CN. 2005. Cytotoxic prenylflavonoids from *Artocarpus elasticus*. J Nat Prod 68: 1692-1695.
- Kuete V, Ango PY, Fotso GW, Kapche GD, Dzoyem JP, Wouking AG, Ngadju BT, Abegaz BM. 2011. Antimicrobial activities of the methanol extract and compounds from *Artocarpus communis* (Moraceae). BMC Complement Altern Med 11: 42.
- Kuete V, Mbaveng AT, Zeino M, Fozing CD, Ngameni B, Kapche GD, Ngadju BT, Effert T. 2015. Cytotoxicity of three naturally occurring flavonoid-derived compounds (artocarpesin, cycloartocarpesin and isobavachalcone) towards multi-factorial drug-resistant cancer cells. Phytomedicine 22: 1096-102.
- Kuran RY, Ersam T. 2017. Prenylated Flavone of the Bark of *Artocarpus elasticus* from Alor Island-NTT Indonesia. J Applied Chem Sci 4: 322-324.
- Lang L, Dong N, Wu D, Yao X, Lu W, Zhang C, Ouyang P, Zhu J, Tang Y, Wang W, Li J, Huang J. 2016. 2-Arylbenzo[b]furan derivatives as potent human lipoxygenase inhibitors. J Enzyme Inhib Med Chem 31: 98-105.
- Lee CC, Lin CN, Jow GM. 2006. Cytotoxic and apoptotic effects of prenylflavonoid artonin B in human acute lymphoblastic leukemia cells. Acta Pharmacol Sin 27: 1165-1174.
- Lin CN, Shieh WL, Ko FN, Teng CM. 1993. Antiplatelet activity of some prenylflavonoids. Biochem Pharmacol 45: 509-5012.
- Lin CN, Shieh WL. 1992. Pyranoflavonoids from *Artocarpus communis*. Phytochemistry 31: 2922-2924.
- Lin KW, Liu CH, Tu HY, Ko HH, Wei BL. 2009. Antioxidant prenylflavonoids from *Artocarpus communis* and *Artocarpus elasticus*. Food Chem 115: 558-562.
- Liu X, Yang J, Gao L, Zhang L, Lei X. 2020. Chemoenzymatic Total Syntheses of Artonin I with an Intermolecular Diels-Alderase. Biotechnol J 15: e2000119.
- Losuwannarak N, Sritularak B, Chanvorachote P. 2018. Cycloartobiloxanthone Induces Human Lung Cancer Cell Apoptosis via Mitochondria-dependent Apoptotic Pathway. In Vivo 32: 71-78.
- Luzuriaga-Quichimbo CX, Blanco-Salas J, Cerón-Martínez CE, Ruiz-Téllez T. 2019. Providing added value to local uses of paparagua (*Artocarpus altilis*) in Amazonian Ecuador by phytochemical data review. Revista Brasileira de Farmacognosia 29: 62-68.
- Ma JP, Qiao X, Pan S, Shen H, Zhu GF, Hou AJ. 2010. New isoprenylated flavonoids and cytotoxic constituents from *Artocarpus tonkinensis*. J Asian Nat Prod Res 12: 586-592.
- Mainasara MM, Abu Bakar MF. 2019. Phytochemical constituents, antioxidant and antiproliferative properties of *Artocarpus altilis* (sukun) from Endau Rompin, Johor, Malaysia. Sci World J 14: 124-135.
- Makmur L, Syamsurizal S, Tukiran T, Achmad SA, Aimi N, Hakim EH, Kitajima M, Takayama H. 2000. Artoindonesianin C, a new xanthone derivative from *Artocarpus teysmanii*. J Nat Prod 63: 243-244.
- Minakawa T, Toume K, Arai MA, Koyano T, Kowithayakorn T, Ishibashi M. 2013. Prenylflavonoids isolated from *Artocarpus champeden* with TRAIL-resistance overcoming activity. Phytochemistry 96: 299-304. Corrigendum in Phytochemistry 2014, 106: 207.
- Mohammed Haleel PM, Rashid K, Senthil Kumar S. 2018. *Artocarpus Heterophyllum*: Review Study on Potential Activities. Res J Pharmacol Pharmacodyn 10: 24-28.
- Molčanová L, Janošiková D, Dall'Acqua S, Smejkal KS. 2019. C-prenylated flavonoids with potential cytotoxic activity against solid tumor cell lines. Phytochem Rev 18: 1051-1100.
- Morrison JJ, Zhang J, Lin J, Murray JE, Porter R, Langat MK, Sadgrove NJ, Barker J, Zhang G, Delgoda R. 2021. Potential chemopreventive, anticancer and anti-inflammatory properties of a refined artocarpin-rich wood extract of *Artocarpus heterophyllum* Lam. Sci Rep 11: 6854.
- Musthapa I, Hakim EH, Juliawaty LD, Syah YM, Achmad SA. 2010. Prenylated flavones from some Indonesian *Artocarpus* and their antimalarial properties. Med Plant 2: 157-160.
- Musthapa I, Juliawaty LD, Syah YM, Hakim EH, Latip J, Ghisalberti EL. 2009a. An oxepinoflavone from *Artocarpus elasticus* with cytotoxic activity against P-388 cells. Arch Pharm Res 32: 191-194.
- Musthapa I, Latip J, Takayama H, Juliawaty LD, Hakim EH, Syah YM. 2009b. Prenylated flavones from *Artocarpus lanceifolius* and their cytotoxic properties against P-388 cells. Nat Prod Commun 4: 927-930.
- Namdaung U, Aroonrerk N, Suksamram S, Danwisetkanjana K, Saenboonrueng J, Arjchomphu W, Suksamram A. 2006. Bioactive constituents of the root bark of *Artocarpus rigidus* subsp. *rigidus*. Chem Pharm Bull (Tokyo) 54: 1433-1436.
- Nimmuan-ngam S, Yangnok K, Innajak S, Watanapokasin R. 2020. Effect of artonin E on apoptosis cell death induction in colon cancer cells. Cancer Epidemiol Biomarker Prev 29, B061.
- Nomura T, Hano Y, Aida M. 1998. Isoprenoid-substituted flavonoids from *Artocarpus* plants (Moraceae). Heterocycles 47: 1179-1205.
- Noraini J. Phytochemicals and bioactivities of *Artocarpus anisophyllum* MIQ. 2013. Masters thesis, Universiti Teknologi Malaysia, Faculty of Science (<http://eprints.utm.my/id/eprint/33249/>).
- Pang D, Liao S, Wang W, Mu L, Li E, Shen W, Liu F, Zou Y. 2019. Destruction of the cell membrane and inhibition of cell phosphatidic acid biosynthesis in *Staphylococcus aureus*: an explanation for the antibacterial mechanism of morusin. Food Funct 10: 6438-6446.
- Panthong K, Tohdee K, Towatana NH, Voravuthikunchai SP, Chusrib S. 2013. Prenylated Flavone from Roots of a Hybrid between

- Artocarpus heterophyllus* and *Artocarpus integer* and its Biological Activities. *J Braz Chem Soc* 24: 1656-1661.
- Park D, Ha JJ, Park SY, Choi M, Lim SL, Kim SH, Lee JH, Ahn KS, Yun M, Lee SG. 2016. Morusin Induces TRAIL Sensitization by regulating EGFR and DR5 in Human Glioblastoma Cells. *J Nat Prod* 79: 317-323.
- Parvatikar PP, Madagi SB. 2021. Molecular Docking Analysis: Interaction Studies of Natural Compounds with Human TG2 Protein. In S.-I. Ao et al. (Eds.): WCECS 2018, *Transactions on Engineering Technologies*. Springer Nature Singapore Pte Ltd., Singapore
- Pedro M, Ferreira MM, Cidade H, Kijjoa A, Bronze-da-Rocha E, Nascimento MS. 2005. Artelastin is a cytotoxic prenylated flavone that disturbs microtubules and interferes with DNA replication in MCF-7 human breast cancer cells. *Life Sci* 77: 293-311.
- Plaibua K, Pongrakhananon V, Chunhacha P, Sritularak B, Chanvorachote P. 2013. Effects of artonin E on migration and invasion capabilities of human lung cancer cells. *Anticancer Res* 33: 3079-3088.
- Qiao X, Zhao T, Wang M, Lei C, Hou A. 2011. [Flavonoids from *Artocarpus hypargyreus*]. *Zhongguo Zhong Yao Za Zhi* 36: 2975-2979.
- Rahman MA, Ramli F, Karimian H, Dehghan F, Nordin N, Ali HM, Mohan S, Hashim NM. 2016. Artonin E Induces Apoptosis via Mitochondrial Dysregulation in SKOV-3 Ovarian Cancer Cells. *PLoS One* 11: e0151466.
- Ramli F, Rahmani M, Ismail IS, Sukari MA, Abd Rahman M, Zajmi A, Akim AM, Hashim NM, Go R. 2016. A New Bioactive Secondary Metabolite from *Artocarpus elasticus*. *Nat Prod Commun* 11: 1103-1106.
- Ranasinghe RASN, Maduwanthi SDT, Marapana RAUJ. 2019. Nutritional and Health Benefits of Jackfruit (*Artocarpus heterophyllus* Lam.): A Review. *Int J Food Sci* 2019: 4327183.
- Rao GV, Mukhopadhyay T, Radhakrishnan N. 2010. Artoindonesianin F, a potent tyrosinase inhibitor from the roots of *Artocarpus heterophyllus* Lam. *Indian J Chem* 49B: 1264-1266.
- Reddy GR, Ueda N, Hada T, Sackeyfio AC, Yamamoto S, Hano Y, Aida M, Nomura T. 1991. A prenylflavone, artonin E, as arachidonate 5-lipoxygenase inhibitor. *Biochem Pharmacol* 41: 115-118.
- Ren Y, Kardono LB, Riswan S, Chai H, Farnsworth NR, Soejarto DD, Carcache de Blanco EJ, Kinghorn AD. 2010. Cytotoxic and NF-kappaB inhibitory constituents of *Artocarpus rigida*. *J Nat Prod* 73: 949-955.
- Sahu SK, Liu M, Yssel A, Kariba R, Muthemba S, Jiang S, Song B, Hendre PS, Muchugi A, Jamnadass R, Kao SM, Featherston J, Zerega NJC, Xu X, Yang H, Deynze AV, Peer YV, Liu X, Liu H. 2019. Draft Genomes of Two *Artocarpus* Plants, Jackfruit (*A. heterophyllus*) and Breadfruit (*A. altilis*). *Genes (Basel)* 11: 27.
- Seo EK, Lee D, Shin YG, Chai HB, Navarro HA, Kardono LB, Rahman I, Cordell GA, Farnsworth NR, Pezzuto JM, Kinghorn AD, Wani MC, Wall ME. 2003. Bioactive prenylated flavonoids from the stem bark of *Artocarpus kemando*. *Arch Pharm Res* 26: 124-127.
- Shaha MKK, Sirata HM, Jamil S, Jalil J. 2016. Flavonoids from the Bark of *Artocarpus integer* var. *silvestris* and their Anti-inflammatory Properties. *Nat Prod Commun* 11: 1275-1278.
- Shahwar D, Iqbal MJ, Nisa MU, Todorovska M, Attar R, Sabitaliyevich UY, Farooqi AA, Ahmad A, Xu B. 2019. Natural Product Mediated Regulation of Death Receptors and Intracellular Machinery: Fresh from the Pipeline about TRAIL-Mediated Signaling and Natural TRAIL Sensitizers. *Int J Mol Sci* 20: 2010.
- Shamaun SS, Rahmani M, Hashim NM, Ismail HB, Sukari MA, Lian GE, Go R. 2010. Prenylated flavones from *Artocarpus altilis*. *J Nat Med* 64: 478-81.
- Shinomiya K, Aida M, Hano Y, Nomura T. 1995. A diels-alder-type adduct from *Artocarpus heterophyllus*. *Phytochemistry* 40: 1317-1319.
- Shinomiya K, Hano Y, Nomura T. 2000. Mechanism on one-sided Wessely-Moser rearrangement reaction. *Heterocycles* 53: 877-886.
- Shobana S, Veena S, Sameer SSM, Swarnalakshmi K, Vishal LA. 2020. Green Synthesis of Silver Nanoparticles Using *Artocarpus hirsutus* Seed Extract and its Antibacterial Activity. *Curr Pharm Biotechnol* 21: 980-989.
- Sikarwar MS, Hui BJ, Subramaniam K, Valeisamy BD, Yean LK, Balaji K. 2014. A Review on *Artocarpus altilis* (Parkinson) Fosberg (breadfruit). *J Applied Pharm Sci* 4: 91-97.
- Šmejkal K. 2014. Cytotoxic potential of C-prenylated flavonoids. *Phytochem Rev* 13: 245-275.
- Soekamto NH, Achmad SA, Ghisalberti EL, Hakim EH, Syah YM. 2003. Artoindonesianins X and Y, two isoprenylated 2-arylbenzofurans, from *Artocarpus fretessi* (Moraceae). *Phytochemistry* 64: 831-834.
- Sophonmithprasert T, Mahabusarakam W, Watanapokasin R. 2019. Artonin E sensitizes TRAIL-induced apoptosis by DR5 upregulation and cFLIP downregulation in TRAIL-refractory colorectal cancer LoVo cells. *J Gastrointest Oncol* 10: 209-217.
- Sritularak B, Tantivanont A, Chanvorachote P, Meksawan K, Miyamoto T, Kohno Y, Likhitwitayawuid K. 2010. Flavonoids with free radical scavenging activity and nitric oxide inhibitory effect from the stem bark of *Artocarpus gomezianus*. *J Med Plants Res* 4: 387-392.
- Suhartati T, Achmad SA, Aimi N, Hakim EH, Kitajima M, Takayama H, Takeya K. 2001. Artoindonesianin L, a new prenylated flavone with cytotoxic activity from *Artocarpus rotunda*. *Fitoterapia* 72: 912-918.
- Suhartati T, Hernawan H, Suwandi JF, Yandri Y, Hadi S. 2018. Isolation of Artonin E from the root bark of *Artocarpus rigida*, synthesis of Artonin E acetate and evaluation of anticancer activity. *Macedonian J Chem & Chem Engineer* 37: 35-42.
- Suhartati T, Wulandari A, Suwandi JF, Yandri Y, Hadi S. 2016. Artonin O, a Xanthone Compound from Root Wood of *Artocarpus rigida*. *Orient J Chem* 32: 2777-2784.
- Suhartati T, Yandri Y, Hadi S. 2008. The Bioactivity Test of Artonin E from the Bark of *Artocarpus Rigida* Blume. *Eur J Sci Res* 23: 330-337.
- Susiarti S, Rahayu M, Kuncari ES, Astuti IP. 2020. Utilization of "Benda" (*Artocarpus elasticus* Reinw. ex Blume) in Bogor, West Java, Indonesia: An Ethnobotanical Case Study. *J Trop Biol Conserv* 17: 297-307.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Makmur L, Mujahidin D. 2001. Artoindonesianins G-I, three new isoprenylated flavones from *Artocarpus lanceifolius*. *Fitoterapia* 72: 765-773.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Makmur L, Mujahidin D. 2002b. Artoindonesianin M, a new prenylated flavone from *Artocarpus champeden*. *Bull Soc Nat Prod Chem (Indonesia)* 2: 31-36.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Makmur L, Mujahidin D. 2002a. Artoindonesianins Q-T, four isoprenylated flavones from *Artocarpus champeden* Spreng. (Moraceae). *Phytochemistry* 61: 949-953.
- Syah YM, Achmad SA, Ghisalberti EL, Hakim EH, Mujahidin D. 2004. Two new cytotoxic isoprenylated flavones, artoindonesianins U and V, from the heartwood of *Artocarpus champeden*. *Fitoterapia* 75: 134-140.
- Syah YM, Juliawaty LD, Achmad SA, Hakim EH, Ghisalberti E. 2006a. Cytotoxic prenylated flavones from *Artocarpus champeden*. *J Nat Med* 60: 308-312.
- Syah YM, Hakim EH, Makmur L, Kurdi VA, Ghisalberti E, Aimi N, Achmad SA. 2006b. Prenylated 2-arylbenzofurans from two Species of *Artocarpus*. *Nat Product Commun* 1: 549-552.
- Teo SP, Nasution RE. 2003. *Artocarpus elasticus* Reinw. ex Blume. In: Brink M, Escobin RP. (eds.). *Plant Resources of South-East Asia No 17: Fibre plants*. PROSEA Foundation, Bogor, Indonesia. prota4u.org/prosea
- Toume K, Habu T, Arai MA, Koyano T, Kowithayakorn T, Ishibashi M. 2015. Prenylated flavonoids and resveratrol derivatives isolated from *Artocarpus communis* with the ability to overcome TRAIL resistance. *J Nat Prod* 78: 103-110. Erratum in *J Nat Prod* 78: 968.
- Tungsukruthai S, Sritularak B, Chanvorachote P. 2017. Cycloartobioxanthone Inhibits Migration and Invasion of Lung Cancer Cells. *Anticancer Res* 37: 6311-6319.
- Tzeng CW, Tzeng WS, Lin LT, Lee CW, Yen FL, Lin CC. 2016. Enhanced autophagic activity of artocarpin in human hepatocellular carcinoma cells through improving its solubility by a nanoparticle system. *Phytomedicine* 23: 528-540.
- Veitch NC, Grayer RJ. 2008. Flavonoids and their glycosides, including anthocyanins. *Nat Prod Rep* 25: 555-611.
- Veriyan A, Raffinal R, Linda R. 2019. Kajian Etnobotani Serat Kulit Kayu Kepuak (*Artocarpus elasticus* Reinw. ex Blume) Pada Suku Dayak Desa'di Desa Kunyai Kecamatan Sungai Tebelian Kabupaten Sintang. *Protobiont* 8: 41-45.
- Wang JP, Tsao LT, Raung SL, Lin PL, Lin CN. 1999. Stimulation of respiratory burst by cyclocommunin in rat neutrophils is associated with the increase in cellular Ca²⁺ and protein kinase C activity. *Free Radic Biol Med* 26: 580-588.
- Wang YH, Hou AJ, Chen L, Chen DF, Sun HD, Zhao QS, Bastow KF, Nakanish Y, Wang XH, Lee KH. 2004. New isoprenylated flavones,

- artochamins A--E, and cytotoxic principles from *Artocarpus chama*. *J Nat Prod* 67: 757-761.
- Wei BL, Weng JR, Chiu PH, Hung CF, Wang JP, Lin CN. 2005. Anti-inflammatory flavonoids from *Artocarpus heterophyllus* and *Artocarpus communis*. *J Agric Food Chem* 53: 3867-3871.
- Widyawaruyanti A, Subehan S, Kalauni SK, Awale S, Nindatu M, Zaini NC, Syafruddin D, Asih PBS, Tezuka Y, Kadota S. 2007. New prenylated flavones from *Artocarpus champeden*, and their antimalarial activity in vitro. *J Nat Med* 61: 410-413.
- Williams EW, Gardner EM, Harris R 3rd, Chaveerach A, Pereira JT, Zerega NJ. 2017. Out of Borneo: biogeography, phylogeny and divergence date estimates of *Artocarpus* (Moraceae). *Ann Bot* 119: 611-627.
- Wongpankam E, Chunhacha P, Pongrakhananon V, Sritularak B, Chanvorachote P. 2012. Artonin E mediates MCL1 down-regulation and sensitizes lung cancer cells to anoikis. *Anticancer Res* 32: 5343-5351.
- Ye JB, Ren G, Li WY, Zhong GY, Zhang M, Yuan JB, Lu T. 2019. Characterization and Identification of Prenylated Flavonoids from *Artocarpus heterophyllus* Lam. Roots by Quadrupole Time-Of-Flight and Linear Trap Quadrupole Orbitrap Mass Spectrometry. *Molecules* 24: 4591.
- Zajmi A, Mohd Hashim N, Noordin MI, Khalifa SA, Ramli F, Mohd Ali H, El-Seedi HR. 2015. Ultrastructural Study on the Antibacterial Activity of Artonin E versus Streptomycin against *Staphylococcus aureus* Strains. *PLoS One* 10: e0128157.
- Zerega NJC, Nur Supardi MN, Motley TJ. 2010. Phylogeny and Recircumscription of Artocarpeae (Moraceae) with a Focus on *Artocarpus*. *Systematic Bot* 35: 766-782.
- Zhai XX, Xiao CY, Jiang W, Chen YL, Ding YQ, Ren G. 2017. Isoprenylated Phenolics from Roots of *Artocarpus heterophyllus*. *Nat Prod Commun* 12: 921-924.
- Zheng R, Zhao L, Chen X, Liu L, Liu Y, Chen X, Wang C, Yu X, Cheng H, Li S. 2021. Metal-coordinated nanomedicine for combined tumor therapy by inducing paraptosis and apoptosis. *J Control Release* 336: 159-168.
- Zheng ZP, Chen S, Wang S, Wang XC, Cheng KW, Wu JJ, Yang D, Wang M. 2009. Chemical components and tyrosinase inhibitors from the twigs of *Artocarpus heterophyllus*. *J Agric Food Chem* 57: 6649-6655.
- Zheng ZP, Cheng KW, To JT, Li H, Wang M. 2008. Isolation of tyrosinase inhibitors from *Artocarpus heterophyllus* and use of its extract as anti-browning agent. *Mol Nutr Food Res* 52: 1530-1538.
- Zoofishan Z, Hohmann J, Hunyadi A. 2018. Phenolic antioxidants of *Morus nigra* roots, and antitumor potential of morusin. *Phytochemistry Rev* 17: 1031-1045.

Phytochemical analysis and in-vitro antimicrobial screening of the leaf extract of *Senna occidentalis* (Fabaceae)

A.A. TAMASI¹, M.O. SHOGE^{1,✉}, T.T. ADEGBOYEGA², E.C. CHUKWUMA³

¹Department of Chemistry, Faculty of Science, Air Force Institute of Technology, PMB 2014, Nigerian Air Force Base, Kaduna, Kaduna State, Nigeria.

✉email: moshachemist@yahoo.com, m.shoge@afit.edu.ng

²Biology Unit, Faculty of Science, Air Force Institute of Technology, PMB 2014, Nigerian Air Force Base, Kaduna, Kaduna State, Nigeria

³Forest Herbarium Ibadan, Forestry Research Institute of Nigeria, Jericho Hills, Ibadan, Oyo State, Nigeria

Manuscript received: 25 February 2021. Revision accepted: 31 August 2021.

Abstract. Tamasi AA, Shoge MO, Adegboyega TT, Chukwuma EC. 2021. Phytochemical analysis and in-vitro antimicrobial screening of the leaf extract of *Senna occidentalis* (Fabaceae). *Asian J Nat Prod Biochem* 19: 57-64. Due to the use of *Senna occidentalis* as an antimalarial, anti-inflammatory, antioxidant, hepatoprotective, and antibacterial agent traditionally, this study examined the phytochemical composition and antimicrobial activity of the isolated methanol, ethyl acetate, and hexane fractions of the leaves of *Senna occidentalis*. Qualitative and quantitative phytochemical analyses were carried out using standard procedures. Antimicrobial activity was performed by using standard procedures against known common clinical isolates. Qualitative phytochemical analysis showed the abundant presence of saponins, tannins, flavonoids, and terpenoids in the ethyl acetate fraction. Cardiac glycosides were not detected in all the tested isolated fractions while steroids were found to be present and abundant in ethyl acetate and hexane fractions respectively. Quantitative phytochemical analysis showed that the methanol fraction contained 4%w/w alkaloids, 0.51%w/w flavonoids, 9.5%w/w saponins, 1.3%w/w tannins 5.5%w/w terpenoids, and 0.6%w/w total phenol content. The antimicrobial susceptibility test shows a range of inhibitory zone of 10-16 mm. Hexane fraction has the highest zone of inhibition against *Candida albicans*. The lowest MIC values of 0.6mg/mL and 0.2mg/mL were observed in ethyl acetate fraction against *Candida albicans* and *Escherichia coli* respectively. The highest MIC was 20mg/mL was observed in the ethyl acetate fraction against *Staphylococcus aureus*. The MMC values were varied widely. The MMC value against *Bacillus subtilis*, *Candida albicans*, and *Trichophyton rubrum* was 5mg/mL while the MMC value against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Escherichia coli* was 20mg/mL. The results from this study show that the leaves of *Senna occidentalis* can serve as a potential source of some phytochemicals and also have the potential to be developed as a source of antibiotics.

Keywords: Alternate medicine, antimicrobial, phytochemicals, *Senna occidentalis*

INTRODUCTION

Medicinal plants are traditionally used globally for the treatment of many diseases, including but not limited to; asthma, gastrointestinal problems, skin disorders, respiratory and urinary problems, hepatic and cardiovascular disease (Tian et al. 2014). Plants synthesize a wide range of biologically active compounds (Cushnie et al. 2014) that are important for them to survive and flourish in the natural environment, protecting them from abiotic stress, insect pests as well as disease infections (Vardhini and Anjum 2015). Since medicinal plants possess an effective source of antimicrobial natural products with proven potential of treating infectious diseases compared to the synthetic drug, there has been a gradual revival of interest in the use of medicinal plants in developed and developing countries (Akinyemi et al. 2006; Ajayi and Akintola 2010; Valarmathy et al. 2010). The natural products from medicinal plants are used in pharmaceutical preparations either as pure or as extracts (Gogtay et al. 2002).

Senna is a large genus of flowering plants in the legume family; Fabaceae (Leguminosae). This genus is distributed throughout the tropics comprising more than

300 species (Gebrelibanos et al. 2014). *Senna* species are commonly used as ornamentals, famine foods, and many species are used in both traditional and modern medicines (Haraguchi et al. 1998). *Senna occidentalis* (L.) Link. (Syn. *Cassia occidentalis* L.) is commonly known as coffee *Senna* (Odeja et al. 2014). It is found in open pastures and fields, cultivated with cereals such as soybean, corn, sorghum, and others (Barbosa-Ferreira et al. 2005). Various parts of *S. occidentalis* (seeds, roots, leaves, and stems) are traditionally claimed to be useful in treating different medical conditions and infections (Bekele and Reddy 2015). It has been used as a folklore medicine for hepatotoxicity treatment. The aqueous-ethanolic leaves extract was tested for hepatoprotective activity on liver damage in paracetamol-induced hepatotoxicity in rats shows that the paracetamol-induced hepatotoxic indications were to some degree reduced after 21 days of treatment with *Senna occidentalis* leaves extract in a dose-dependent manner (Uzzi and Grillo 2013). Aqueous extract of *Senna occidentalis* tested for antidiabetic activity showed a significant reduction in fasting blood glucose levels in the normal and alloxan-induced diabetic rats (Laxmi 2010).

Previous studies on *S. occidentalis* extract in four different solvents (methanol, aqueous, benzene, petroleum

ether, and chloroform) showed that methanol extract had antimicrobial activity against *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Escherichia coli*, *Staphylococcus aureus*, and *Staphylococcus epidermidis*. The aqueous extract was effective against *Proteus vulgaris*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa*; while benzene and petroleum ether extracts were active against *Proteus mirabilis* and *Escherichia coli*; and chloroform extract was inactive against all tested strains (Arya et al. 2010). Further, a study by Daniyan et al. (2011) showed that flower extract of *S. occidentalis* had maximum inhibition against *Klebsiella pneumoniae* and no activity against *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Pseudomonas aeruginosa*. Thus, the flower extract of *S. occidentalis* can be used to treat *Klebsiella*-associated ailments such as pneumonia, bronchitis, and other diseases caused by *Klebsiella pneumoniae*.

The antibacterial and antimalarial activities of leaves and root bark extracts of this plant have also been reported (Tona et al. 2001; Chukwujekwu et al. 2006). Studies on the ethanolic, dichloromethane and lyophilized aqueous extracts of *S. occidentalis* root bark were tested for antimalarial activity against *Plasmodium berghei*. The extracts produced significant chemo suppressions of parasitemia with 200 mg/kg dose when administered orally in experimental animals (mice). The species was found to be potential with 60% chemo suppression of the parasites.

Ibrahim et al. (2010) attempt to demonstrate the *in vitro* and *in vivo* activities of *S. occidentalis* against *Trypanosoma brucei*. Although the exact mechanism for the observed *in vivo* trypano-suppressive effect is not known, it is obvious that the extract contains some phytochemicals that could interfere with the survival of the parasites *in vivo*. The presence of alkaloids, flavonoids, tannins, and anthraquinones have been reported in this extract (Ogunkunle and Ladejobi 2006) and previous reports attributed the antitrypanosomal activity of several tropical plants to the flavonoids (azaanthraquinone), highly aromatic planar quaternary alkaloids, barbarine and harmaine (Nok 2001). The trypanosuppressive action of this extract could be due to the presence of one or more of these bioactive compounds. The present study examined the phytochemical compounds of *S. occidentalis* leaves which could be useful for the synthesis of orthodox drugs, and also its antimicrobial activity.

MATERIALS AND METHODS

Sample collection and preparation

The plant was collected at Unguwar Dosa, Kaduna, Nigeria and identified by Prof Gabriel Ajibade of Biological Sciences Department, Nigerian Defence Academy, Kaduna. It was authenticated at the Forest Herbarium Ibadan (FHI) (Holmgren et al. 1990). Upon collection, the leaves of the plant were rinsed with distilled water, air-dried at room temperature for two weeks, and then blended into fine powder. Active compounds from the

plant were extracted in a crude form through cold maceration extraction method using methanol for 72 hours, concentrated at 40°C using rotary evaporator and the crude extract fractionated with hexane, ethyl acetate, and methanol successively in a 500ml separating funnel.

Figure 1.0 illustrates the morphology of *S. occidentalis*, while Figure 2 shows the current distribution of the species in Nigeria.

Qualitative phytochemical screening

The hexane, ethyl acetate, and methanol fractions of *S. occidentalis* were assessed for the existence of some phytochemicals using the following standard methods:

Test for Alkaloids

(i) Dragendroff's reagent: 8 g of Bi (NO₃.5H₂O) was dissolved in 20 ml of nitric acid and 2.72g of potassium iodide in 50 ml of water, mixed and makeup to 100ml with distilled water.

Test: To 0.1g of extract, added 2.0 ml of HCl. To this acidic medium, 1 ml of reagent was added. An orange-red precipitate produced immediately indicates the presence of alkaloids.

(ii) Wagner's reagent: 1.0 g of iodine and 2.0 g of potassium iodide were dissolved in 5 ml sulphuric acid and the solution was diluted to 100 ml.

Test: 0.1g extract was acidified by adding 1.5% v/v HCl and a few drops of Wagner's reagent. The formation of yellow or brown precipitate confirmed the presence of alkaloids.

(iii) Meyer's reagent: 1.36 g Mercuric chloride was dissolved in 60 ml of distilled water and 5g of potassium iodide in 10 ml of water. The two solutions were mixed and diluted to 100 ml with distilled water.

Test: To 0.1g ml of extract, few drops of reagent were added. The formation of white or pale precipitate showed the presence of alkaloids.



Figure 1. Morphology of *Senna occidentalis* (L.) Link.




Figure 2. Distribution of *S. occidentalis* in Nigeria

Test for flavonoids: ammonia/H₂SO₄

The sample was added to 5 ml of diluted ammonia solution and concentrated H₂SO₄. The disappearance of yellow color on standing indicates the presence of flavonoids.

Test for saponins: Frothing Test

3 ml of the aqueous solution of the extract was mixed was stoppered in a test tube and shaken vigorously for about 30sec, then allowed to stand for 3min and observed for a persist honeycomb froth, which indicates the presence of saponins.

Test for tannins: Ferric chloride method

2 ml of the aqueous extract was added to 3 drops of 10% Ferric chloride solution. The occurrence of blue-black color showed the presence of tannins

Anthraquinone test: Borntrager's test

0.1g of the dried extract was placed in a dry test tube and 5 ml of chloroform was added and was heated in a steam bath for 1 min. The extract was filtered while hot and allowed to cool. A 10% ammonia solution was added to the filtrate and then shaken. The upper aqueous layer was observed for bright pink indicates the presence of anthraquinones.

Terpenoids test: Salkowski test

0.1g of the sample was mixed with 2ml of CHCl₃ in a test tube 3ml of concentrated H₂SO₄ was carefully added to the mixture to form a layer. An interface with a reddish-brown coloration was formed if terpenoids constituent is present.

Cardiac glycosides Test: Keller-Killiani test

A solution of glacial acetic acid (4.0 ml) with 1 drop of 2.0% FeCl₃ was mixed with 10 ml aqueous plant extract and 1 ml concentrated H₂SO₄. A brown ring formed between the layers showed the presence of cardiac glycosides.

Phenol test: Ferric chloride test

To 1 ml aliquot of each of the fractions, was added with 3-4 drops of neutral 5% ferric chloride solution. The formation of dark green color indicated the presence of phenols.

Steroids test: Liebermann-Burchard test

The extracts were added 5ml of chloroform, shake and filtered. and few drops of acetic anhydride along with a few drops of concentrated sulphuric acid were added through the side of the tube. The appearance of green color indicates the presence of sterols in the extracts.

Quantitative phytochemical screening

Quantitative determination of alkaloids

Quantitative determination of alkaloids was performed according to the phytochemical methods described by Harborne (1973). Exactly 200 cm³ of 10% acetic acid in ethanol was added to each fraction (1g) in a 250 cm³ beaker and allowed to stand for 4 hours. The fractions were concentrated on a water bath to ¼ of its original volume and 10ml of concentrated ammonium hydroxide was added until the precipitation was complete and filtered after 3 hours with a pre-weighed filter paper. The residues were dried in an oven and the percentage of alkaloid for each fraction is calculated mathematically as follows:

$$\% \text{ Alkaloids} = \frac{\text{Weight of alkaloid} \times 100}{\text{Weight of samples}}$$

Quantitative determination of terpenoids

Dried plant extracts (1g) of each fraction (Wi) were soaked in 20 ml of ethanol for 24 hours (Indumathi et al. 2014). After filtration, the extracts were extracted with 10 ml of petroleum ether using a separating funnel. The ether extract was separated into pre-weighed glass vials and kept completely dry (Wf). The yield (%) of total terpenoids contents was measured by the formula:

$$\% \text{ Terpenoids} = \frac{W_i - W_f \times 100}{W_i}$$

Quantitative determination of saponins

Quantitative determination of saponins was carried out using the method by Ejikeme et al. (2014). 100 cm³ of 20% aqueous ethanol was added to 1g of each fraction in 250 cm³ conical flasks. The mixtures were heated over a hot water bath for 4 hours with continuous stirring at a temperature of 55°C. The residue of the mixtures was re-extracted with another 100 cm³ of 20% aqueous ethanol after filtration and heated for 4 hours at a constant temperature of 55°C with constant stirring. The extracts were evaporated to 40 cm³ over a water bath at 90°C. 20 cm³ of diethyl ether was added to the concentrate in a 250 cm³ separating funnel and vigorously agitated from which the aqueous layer was recovered while the ether layer was discarded. This purification process was repeated twice. 40 cm³ of n-butanol was added to the aqueous portion, shook, allowed to partition and the butanol layer collected after 10min. this process was also repeated twice and the combined butanol layer was washed with 20 cm³ of 5% sodium chloride and heated in a pre-weighed crucible. The saponin content was calculated as follows:

$$\% \text{ Saponins} = \frac{\text{Weight of saponins} \times 100}{\text{Weight of samples}}$$

Quantitative determination of tannin (using the spectrophotometric method)

Tannins contents were quantitatively determined by the Folin-Ciocalteu method for ethyl acetate and methanol fractions. 0.1ml of each fraction were added to volumetric flasks (10 ml) both containing 7.5 ml of distilled water and

0.5 ml of Folin-Ciocalteu phenol reagent, 1 ml of 35 % Na₂CO₃ solution was diluted to 10 ml with distilled water. The mixtures were shaken well and kept at room temperature for 30 minutes. Reference standard solutions of gallic acid (32, 63, 125, 250, 500, and 1000 µg/mL) were prepared. Absorbance for test and standard solutions were measured against the blank at 760 nm with a UV/Visible spectrophotometer (Miean and Mohamed 2001; Ribarova and Atanassova 2005).

Determination of total flavonoid content

Total flavonoids content was determined from the calibration curve of quercetin and expressed as a microgram of quercetin equivalent per milligram of extract. Total flavonoid content was determined according to the procedure of (Chang et al. 2002), validated by (Nugroho et al. 2013) with some modifications. 1mg/1 ml standard solution was prepared, and series concentrations of quercetin were then obtained. The sample solution (0.5ml) was added with 1.5 ml methanol, 0.1 ml of 10% AlCl₃, 0.1 ml Potassium acetate 1M, and 2.8 ml of distilled water, and then incubated for 30 minutes. Absorbance was measured at 415 nm. Distilled water was used as blank. Total flavonoid content was expressed in mg quercetin per 1g of plant extract.

Total phenolic content

The total phenolic content of ethyl acetate and methanol extracts (200-1000 µg/mL) were quantified using the Folin-Ciocalteu's phenol reagent (Singleton and Rossi 1965) with gallic acid (50-200µg/mL) as standard. Each extract (1 ml) was added to 1 ml Folin-Ciocalteu 's reagent (diluted tenfold in distilled water) in separate test tubes. The content of each test tube was mixed and allowed to stand for five minutes at 25°C in an incubator. 1 ml of 2 % sodium carbonate solution (Na₂CO₃) was added to the mixture. This was allowed to stand for 2 hours at 25°C in an incubator and centrifuged at 1000 g for 10 minutes to get a clear solution. The absorbance of the supernatant was then determined at 760 nm using a UV spectrophotometer. Distilled water (1 ml) was added to 1 ml Folin-Ciocalteu's reagent (diluted ten-fold in distilled water) processed in the same way as done for the test samples and used as blank. All measurements were done in triplicates. The total phenolics were expressed as micrograms per milliliter of gallic acid equivalents through the calibration curve with gallic acid.

Antimicrobial susceptibility test of *Senna occidentalis*

Determination of antimicrobial activity by standard procedures

Isolates of micro-organisms were obtained from the Department of Pharmaceutical Microbiology, University of Ibadan, Nigeria. The antimicrobial activity of methanol, ethyl acetate, and hexane fractions of *S. occidentalis* was determined against *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Candida albicans*, and *Trichophyton rubrum*. A suspension of micro-organism pure culture by disk diffusion was spread evenly over the face of a sterile agar plate using a sterile swab. For all the fractions, 100 mg/mL and 50

mg/mL each were applied to the center of the various agar plates (in a fashion such that the antimicrobial doesn't spread out from the center). Gentamicin, 10µg/mL and 1% Ketoconazole were inoculated as controls. The agar plate was incubated for 24 hours at a temperature of 37°C. The zone of inhibition appears when the plant extract fractions exert a growth-inhibiting effect. The diameter of the inhibitory zone was measured in millimeters and it is related to the level of antimicrobial activity present in the fraction; the larger the inhibitory zone usually means the better the antimicrobial potency.

Determination of Minimum Inhibitory Concentration (MIC) & Minimum Microbicidal Concentration (MMC)
(Pramila et al. 2012)

The minimal inhibitory concentration (MIC) is defined as the lowest concentration of the compounds which inhibit the growth of the microorganisms. Minimal bactericidal concentration (MBC) is defined as the lowest concentration of the compounds that kill and show no growth of the microorganisms on an agar plate. The test microbes were grown in nutrient broth and potato dextrose broth for bacteria and fungus respectively as previously described and the cultures were adjusted to 0.5 McFarland standard turbidity. MIC and MBC values of the plant extract against *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Candida albicans*, and *Trichophyton rubrum* were determined based on a micro-well dilution method. The 24-well microtiter plates were prepared by dispensing 2 ml of MH broth into each well and 3 µl of the bacterial and fungal inoculum. The plant extract was dissolved in the fractions to obtain the concentration of 1 mg/mL and was serially diluted and mixed thoroughly. Then the plates were incubated at 37°C for 24h

RESULTS AND DISCUSSION

The screening of plant parts for medicinal properties starts with crude extract preparation using a range of solvents with different polarities to dissolve different phytochemicals compounds in the plant parts. This is a vital step in drug development from the plant. Phytochemical constituents vary quantitatively and qualitatively in different plant species and even in plant parts of the same species depending upon various atmospheric conditions and storage conditions (Jahan et al. 2008). Three solvents (hexane, ethyl acetate, and methanol) were used for the extract preparation. The qualitative phytochemical screening of the three fractions is shown in Table 1.

Saponins and terpenoids were found to be present in all three extracts of *S. occidentalis*. Tannins, flavonoids, alkaloids, and phenols were found to be present in ethyl acetate and methanol extracts only. Steroids and anthraquinones were positive in hexane and ethyl acetate extracts, while cardiac glycoside was completely negative in all three extracts.

The occurrence of this wide range of phytochemicals shows the medicinal potentials of *S. occidentalis*. Alkaloids, saponins, tannins, flavonoids, and steroids have been known to be biologically active thus partially responsible for the antimicrobial activities of plants (Nethathe and Ndip 2011). Flavonoids are known to be synthesized by plants in response to microbial infection (Cowan 1999), antimicrobial properties of saponins are due to the leakage of proteins and certain enzymes from the cell (Zablotowicz et al. 1996) while tannins bind to proline-rich proteins and interfere with the protein synthesis (Shimada 2006).

As revealed in Table 2, the three fractions had a range of terpenoids from 5.5% to 7.5%w/w. Malik et al. (2017) also reported a wide range of terpenoid content, i.e. *Cassia absus* contains 40%, and *Carissa carandas* 80%, in which these two plants were used as medicinal plants in Punjab. Terpenoids are chemically interesting groups of secondary metabolites (Ajaib et al. 2016) of important therapeutic use as antimicrobial, antifungal, antiviral, antihyperglycemic, anti-inflammatory, antioxidants, and antiparasitic (Brahmkshatriya and Brahmkshatriya 2013). Jaeger and Cuny (2016) reported that terpenoids are naturally formed in a stereospecific manner and can be obtained in enantiopure form by extraction from natural sources. Thus, the pure enantiomeric form is needed for the production of highly active medical compounds of high pharmaceutical value.

Saponins have a wider range of 1.5% to 9.5% w/w across the tested fractions. A previous study by Ezeonu and Ejikeme (2016) also reported a wide range of saponin content between *Protea elliottii* (1.6%) to *Anogeissus leiocarpus* (12.5%). The extraction of saponins from these plants serves as a good source of saponins since saponins protect plants against microbial attacks such as yeast and fungal infections (Sheikh et al. 2013). Oyekunle et al. (2006) reported that saponins show antimicrobial activity by inhibiting the growth of Gram-positive and Gram-negative microorganisms. However, some saponins are not effective against Gram-negative microorganisms because saponins are not able to penetrate the cell membranes of Gram-negative bacteria.


Table 1. Phytochemicals compounds of three fractions of *S. occidentalis* leaves

Test	Hexane Fraction	Ethyl acetate Fraction	Methanol Fraction
Saponins	+ve	++ve	++ve
Tannins	-ve	++ve	+ve
Flavonoids	-ve	++ve	+ve
Cardiac glycosides	-ve	-ve	-ve
Anthraquinones	+ve	+ve	-ve
Terpenoids	+ve	++ve	+ve
Steroids	++ve	+ve	-ve
Alkaloids	-ve	+ve	+ve
Phenol	-ve	+ve	+ve

Note: ++ve: Abundant, +ve: Present, -ve: Absent

Table 2. Percentage chemical composition of *S. occidentalis* leaf fractions

Sample	% Alkaloids	% Flavonoid	% Saponin	% Tannin	% Terpenoid	% Total Phenol
Hexane fraction	0	0	1.5%w/w	0	6%w/w	0
Ethylacetate fraction	8%w/w	0.2%w/w	2.5%w/w	1%w/w	7.5%w/w	0.73%w/w
Methanol fraction	4%w/w	0.51%w/w	9.5%w/w	1.3%w/w	5.5%w/w	0.6%w/w

**Figure 3.** Diameter of inhibitory zone of *S. occidentalis* leaf fractions against several microbes. A. Fraction concentrations of 50 mg/mL; B. Fraction concentrations of 100 mg/mL

Flavonoids and phenols content in ethyl acetate and methanol were low, i.e. 0.2%w/w, 0.7%w/w, and 0.51%w/w, 0.6%w/w respectively. These results are also supported by previous studies on the flavonoid content of *Erythrina velutina* (0.2%) and *Mimosa tenuiflora* (0.21%) both of the plants that belong to the Fabaceae family with reported antimicrobial activity (Siqueira et al. 2012). Also, the low phenol content is similar to the report on stem bark (0.75%) and leaves (0.09%) of *Mangifera indica* (Okwu and Ezenagu 2008). Phenols protect plants from oxidative damage and do the same for humans (Okwu 2005).

Tannins content in ethyl acetate and methanol fractions were 1%w/w and 1.3%w/w respectively. Alkaloids were present in ethyl acetate fraction at 8%w/w and methanol fraction at 4%w/w. These values are much higher than reported values of alkaloids in *Cucumis sativus* (cucumber) and *Momordica charantia* (bitter melon) leaves with 1.23%w/w where bitter melon ethanolic leaf extract showed 70% percent inhibition against *E. coli*, *S. dysenteriae* 1, and *S. pneumoniae* while *Cucumis sativus* leaf extract was

found to have less antibacterial activity (Debnath et al. 2015). The differences in methanol and ethyl acetate differences could be due to their inherent polarity and hence influences their performance.

The antimicrobial activities of *S. occidentalis* against *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Candida albicans*, and *Trichophyton rubrum* were examined in this study and their potency was qualitatively and quantitatively assessed by the presence or absence of inhibitory zones, minimum inhibitory concentration (MIC), and minimum microbicidal concentrations (MMC).

The zones of inhibition of the tested fractions against the selected bacterial and fungal strains are shown in Figure 3. in a concentration-dependent manner. The result showed that methanol and ethyl acetate fractions at 100 mg/mL and 50 mg/mL did not inhibit the growth of *Pseudomonas aeruginosa*. However, an inhibitory zone of 12 mm was observed for both hexane fraction and the tested control (Gentamicin) at 100 mg/mL and 10 µg/mL respectively. The zone produced by the standard antibiotic was similar to that produced by the hexane fraction of *S. occidentalis*. Similar to the present work, studies on the antibacterial activity of methanolic leaf extract of *Plukenetia conophora* against *Pseudomonas aeruginosa* isolated from urinary tract infection showed a concentration-dependent activity with *Pseudomonas aeruginosa* having zones of inhibition of 10, 12, 14, and 18 mm at 50, 100, 150 and 200 mg/mL respectively (Enitan et al. 2014).

The methanol fraction did not show any inhibition zone against *Staphylococcus aureus* and *Escherichia coli* at the concentrations of 100mg/mL and 50mg/mL. The diameter of inhibition of the ethyl acetate and hexane fractions at the concentration of 100mg/mL and 50mg/mL against *Staphylococcus aureus* was the same, i.e., 12mm and 10mm, respectively. These values were lower than positive control Gentamicin (16 mm). A study by Kumar et al. (2010) showed that the inhibitory zone of cardamom extract against *S. aureus* was 6 mm to 9 mm. The inhibitory zone of hexane, ethyl acetate, and dichloromethane extract against *Escherichia coli* was 12 mm.

The result showed a similar antibacterial activity of methanol and ethyl acetate fractions at the concentration of 100 mg/mL against *Bacillus subtilis* was 14mm, while hexane fraction has the inhibitory zone of 12mm. These values were lower compared to that of Gentamicin (18mm at 10ug/mL). The inhibitory zone of hexane fraction against *C. albicans* at the concentrations of 100 mg/mL and 50 mg/mL were 16mm and 14mm.




Figure 4. The Minimum Inhibitory Concentration (MIC) and Minimum Microbicidal Concentration (MMC) of *S. occidentalis* leaf fractions

This fraction has a similar zone of inhibition for all tested bacteria at the concentrations of 100 mg/mL and 50 mg/mL, i.e. 12 mm and 10mm respectively. However, these values are much lower than the positive control Ket 1% (18 mm). Methanol extract of *S. occidentalis* showed a zone of inhibition of 14mm at 100mg/mL against *Trichophyton rubrum* and 12 mm for ethyl acetate and hexane fractions respectively, which are lower than that of positive control Ket 1% (18mm).

The MIC and MMC results of the tested fractions are depicted in Figure 4. Ethyl acetate fraction shows the lowest MIC value of 0.2mg/mL and 10 mg/mL MMC against *Escherichia coli*. This same fraction also has a MIC value of 0.6mg/mL and 10 mg/mL MMC against *Candida albicans*. The MIC and MMC values of hexane fractions were 2.5mg/mL and 10mg/mL against *Pseudomonas aeruginosa*. The MIC value (5 mg/mL) of methanol fraction was similar against all tested microbial strains except for *Staphylococcus aureus* with the MIC and MMC value of 10 mg/mL. The highest MIC value (20mg/mL) was recorded for ethyl acetate fraction against *Staphylococcus aureus*. Ethyl acetate fraction could be described as having a better antimicrobial activity than other fractions. The results of this study have added to the existing information on *S. occidentalis*, it is also important to consider its sustainability. On the distribution of *S. occidentalis*, the species seem to be more abundant around the transitional equatorial states in Nigeria (Oyo, Ogun, Lagos, Kwara, Osun, Edo, Ekiti, Ondo, Rivers, etc) and less in the Northern regions. Hence, its natural habitats need some attention, to ensure its sustainable collection and use.

In conclusion, the phytochemical analysis of the leaves of *Senna occidentalis* shows the presence of flavonoids, tannins, and phenols and the abundance of alkaloids, saponins, and terpenoids. Antimicrobial activities of the *S. occidentalis* fractions were more potent against the tested fungi; *Candida albicans* and *Trichophyton rubrum* than Gram-positive and Gram-negative bacteria. These results

could be used as scientific evidence of the use of the plant as a source of remedy for various ailments traditionally. Further pharmacognostic studies are suggested to ascertain its potency in the treatment of diseases.

REFERENCES

- Ajaib M, Almas M, Khan KM, Perveen S, Shah S. 2016. Phytochemical screening, antimicrobial and antioxidant activities of *Ficus natalensis*. *J Chem Soc Pak* 38 (2): 345-351.
- Ajayi AO, Akintola TA. 2010. Evaluation of antibacterial activity of some medicinal plants on common enteric food-borne pathogens. *Afr J Microbiol Res* 4 (4): 314-316.
- Akinoyemi KO, Oluwa OK, Omomigbehin EO. 2006. Antimicrobial activity of crude extracts of three medicinal plants used in southwest Nigerian folk medicine on some food-borne bacterial pathogens. *Afr J Tradit Complement Altern Med* 3 (4): 13-22. DOI: 10.4314/ajtcam.v3i4.31173
- Arya V, Yadav S, Kumar S, Yadav JP. 2011. Antioxidant activity of organic and aqueous leaf extracts of *Cassia occidentalis* L. in relation to their phenolic content. *Nat Prod Res* 25 (15): 1473-1479. DOI: 10.1080/14786419.2010.545351
- Barbosa-Ferreira M, Dagli ML, Maiorka PC, Górnica SL. 2005. Sub-acute intoxication by *Senna occidentalis* seeds in rats. *Food Chem Toxicol* 43 (4): 497-503. DOI: 10.1016/j.fct.2004.11.017
- Bekele G, Reddy PR. 2015. Ethnobotanical study of medicinal plants used to treat human ailments by Guji Oromo tribes in Abaya District, Borana, Oromia, Ethiopia. *Universal J Plant Sci* 3 (1): 1-8. DOI: 10.13189/ujps.2015.030101.
- Brahmkshatriya PP, Brahmkshatriya PS. 2013. Terpenes: Chemistry, biological role, and therapeutic applications. In: Ramawat K, Mérillon JM. (eds) *Natural Products*. Springer, Berlin, Heidelberg. DOI: 10.1007/978-3-642-22144-6_120.
- Chang CC, Yang MH, Wen HM, Chern JC. 2002. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal* 5 (2): 46-52. DOI: 10.38212/2224-6614.2748.
- Cowan MM. 1999. Plant products as antimicrobial agents. *Clin Microbiol Rev* 12: 564-582. DOI: 10.1128/CMR.12.4.564
- Cushnie TPT, Cushnie B, Lamb AJ. 2014. Alkaloids: an overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *Intl J Antimicrob Agents* 44: 377-386. DOI: 10.1016/j.ijantimicag.2014.06.001
- Daniyan SY, Oloruntimelehin JB, Ifeadi O. 2011. Antibacterial activity of *Cassia occidentalis* flower vegetable extract on selected bacteria. *Asian J Biomed Pharm Sci* 1 (1). DOI:10.15272/AJBPS.V1I1.5.

- Debnath B, Uddin MJ, Patari P, Das M, Maiti D, Manna K. 2015. Estimation of alkaloids and phenolics of five edible cucurbitaceous plants and their antibacterial activity. *Intl J Pharm Pharm Sci* 7 (12): 223-227.
- Enitan SS, Olley M, Uhumwango SE, Akele YR, Ehiaghe AF, Enitan CB. 2014. Antibacterial activity of methanolic leaf extract of *Plukenetia conophora* Mull. arg. against selected bacteria isolated from urinary tract infection. *Intl J Microbiol Appl* 1 (1): 1-10.
- Ejikeme C, Ezeonu CS, Eboatu AN. 2014. Determination of physical and phytochemical constituents of some tropical timbers indigenous to Niger delta area of Nigeria. *Eur Sci J* 10 (18): 247-270.
- Ezeonu CS, Ejikeme CM. 2016. Qualitative and quantitative determination of phytochemical contents of indigenous Nigerian softwoods. *New J Sci*. DOI: 10.1155/2016/5601327
- Gebrelibanos M, Periyasamy G, Sintayehu B. 2014. *Senna occidentalis* seed: is it health risk or potential medicine? *Intl J Pharm* 1 (3): 161-167.
- Gogtay NJ, Bhatt HA, Dalvi SS, Kshirsagar NA. 2002. The use and safety of non-allopathic Indian medicines. *Drug Saf* 25: 1005-1019. DOI: 10.2165/00002018-200225140-00003.
- Haraguchi M, Gómiak SL, Calore EE, Cavaliere MJ, Raspantini PC, Calore NM, Dagli ML. 1998. Muscle degeneration in chicks caused by *Senna occidentalis* seeds. *Avian Pathol* 27 (4): 346-351. DOI: 10.1080/03079459808419350.
- Harborne JB. 1973. *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. Chapman and Hall, London, UK. DOI: 10.1007/978-94-009-5921-7_1.
- Holmgren PK, Keuken W, Schofield EK. 1990. *Index Herbariorum Part I. The Habaria of the World*. 8th ed. Reg. Veg., New York.
- Ibrahim MA, Aliyu AB, Sallau AB, Bashir M, Yunusa I, Umar TS. 2010. *Senna occidentalis* leaf extract possesses antitrypanosomal activity and ameliorates trypanosome-induced anemia and organ damage. *Pharm Res* 2 (3): 175. DOI: 10.4103/0974-8490.65513.
- Indumathi C, Durgadevi G, Nithyavani S, Gayathri PK. 2014. Estimation of terpenoid content and its antimicrobial property in *Enicostemma littorale*. *Intl J Chem Tech Res* 6 (9): 4264-4267.
- Jaeger R, Cuny E. 2016. Terpenoids with special pharmacological significance: A review. *Nat Prod Commun* 11 (9): 1934578X1601100946. DOI: 10.1177/1934578X1601100946.
- Jahan N, Afaque SH, Khan NA, Ahmad G, Ansari AA. 2008. Physico-chemical studies of the gum acacia.
- Kumar R, Yadav BR, Singh RS. 2010. Genetic determinants of antibiotic resistance in *Staphylococcus aureus* isolates from milk of mastitic crossbred cattle. *Curr Microbiol* 60 (5): 379-386. DOI: 10.1007/s00284-009-9553-1.
- Laxmi V, Singour PK, Chaurasiya PK, Rajak H, Pawar RS, Patil UK. 2010. Effect of ethanolic extract of *Cassia occidentalis* Linn. for the management of alloxan-induced diabetic rats. *Pharm Res* 2 (3): 132-137. DOI: 10.4103/0974-8490.65506.
- Malik SK, Ahmad M, Khan F. 2017. Qualitative and quantitative estimation of terpenoid contents in some important plants of Punjab, Pakistan. *Pak J Sci* 69 (2): 150-154.
- Miean KH, Mohamed S. 2001. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem* 49 (6): 3106-3112. DOI: 10.1021/jf000892m
- Nethathe BB, Ndip RN. 2011. Bioactivity of *Hydnora africana* on selected bacterial pathogens: Preliminary phytochemical screening. *Afr J Microbiol Res* 5 (18): 2820-2826. DOI: 10.5897/AJMR11.566
- Nok AJ. 2002. Azaanthraquinone inhibits respiration and in vitro growth of long slender bloodstream forms of *Trypanosoma congolense*. *Cell Biochem Funct: Cell Biochem Its Modulation Active Agents Dis* 20 (3): 205-212. DOI: 10.1002/cbf.948.
- Nugroho AE, Malik A, Pramono S. 2013. Total phenolic and flavonoid contents, and in vitro antihypertension activity of purified extract of Indonesian cashew leaves (*Anacardium occidentale* L.). *Intl Food Res J* 20 (1): 299-305.
- Odeja OO, Obi G, Ogwuche CE, Elemike EE, Oderinlo OO. 2014. Phytochemical screening, antioxidant and antimicrobial activities of *Senna occidentalis* (L.) leaves. *Intl J Herbal Med* 2 (4): 26-30. DOI: 10.1186/s40816-016-0028-1.
- Ogunkunle AT, Ladejobi TA. 2006. Ethnobotanical and phytochemical studies on some species of *Senna* in Nigeria. *Afr J Biotechnol* 5 (21): 2020-2023.
- Okwu DE, Ezenagu VI. 2008. Evaluation of the phytochemical composition of mango (*Mangifera indica* Linn) stem bark and leaves. *Intl J Chem Sci* 6 (2): 705-716.
- Okwu DE. 2005. Phytochemicals, vitamins and mineral contents of two Nigerian medicinal plants. *Intl J Mol Med Adv Sci* 1 (4): 375-381.
- Oyekunle MA, Aiyelaagbe OO, Fafunso MA. 2006. Evaluation of the antimicrobial activity of saponins extracts of *Sorghum bicolor* L. Moench. *Afr J Biotechnol* 5 (23): 2405-2407.
- Ribarova F, Atanassova M. 2005. Total phenolics and flavonoids in Bulgarian fruits and vegetables. *J Univ Chem Technol Metallurgy* 40 (3): 255-260.
- Sheikh N, Kumar Y, Misra AK, Pfoze L. 2013. Phytochemical screening to validate the ethnobotanical importance of root tubers of *Dioscorea* species of Meghalaya, North East India. *J Med Plants* 1 (6): 62-9.
- Shimada T. 2006. Salivary proteins as a defense against dietary tannins. *J Chem Ecol* 32 (6): 1149-1163. DOI: 10.1007/s10886-006-9077-0.
- Singleton VL, Rossi JA. 1965. Colorimetry of total phenolics with phosphomolybdic phosphotungstic acid reagents. *Am J Enol Vitic* 16 (3): 144-158.
- Siqueira CF, Cabral DL, Peixoto Sobrinho TJ, de Amorim EL, de Melo JG, Araújo TA, de Albuquerque UP. 2012. Levels of tannins and flavonoids in medicinal plants: evaluating bioprospecting strategies. *Evid-Based Compl Altern Med* 2012. DOI: 10.1155/2012/434782.
- Tian XR, Feng JT, Ma ZQ, Xie N, Zhang J, Zhang X, Tang HF. 2014. Three new glycosides from the whole plant of *Clematis lasiantha* Maxim and their cytotoxicity. *Phytochem Lett* 10: 168-172. DOI: 10.1016/j.phytol.2014.09.004.
- Tona L, Mesia K, Ngimbi NP, Chrimwami B, Okond'Ahoka, Cimanga K, Bruyne TD, Apers S, Hermans N, Totte J, Pieters L. 2001. In-vivo antimalarial activity of *Cassia occidentalis*, *Morinda morindoides* and *Phyllanthus niruri*. *Ann Tropic Med Parasitol* 95 (1): 47-57. DOI: 10.1080/00034983.2001.11813614.
- Uzzi HO, Grillo DB. 2013. The hepatoprotective potentials of aqueous leaf extract of *Cassia occidentalis* against paracetamol induced hepatotoxicity in adult Wistar rats. *Intl J Herbs and Pharmacol Res* 2 (2): 6-13.
- Valarmathy K, Gokulakrishnan M, Kausar MS, Paul K. 2010. A study of antimicrobial activity of ethanolic extracts of various plant leaves against selected microbial species. *Int J Pharma Sci Res* 1 (8): 293-295.
- Vardhini BV, Anjum NA. 2015. Brassinosteroids make plant life easier under abiotic stresses mainly by modulating major components of antioxidant defense system. *Front Environ Sci* 2: 67. DOI: 10.3389/fenvs.2014.00067.
- Zablutowicz RM, Hoagland RE, Wagner SC. 1996. Effect of saponins on the growth and activity of rhizosphere bacteria. In: *Saponins Used in Food and Agriculture*. Springer, Boston, MA. DOI: 10.1007/978-1-4613-0413-5_8.

Detection of terpenoids and steroids in *Lindsaea obtusa* with thin layer chromatography

YUSFIA URWATUL WUTSQA¹, SURATMAN^{2,*}, SITI LUSI ARUM SARI²

¹Department of Pharmacy, STIKes KHAS Kempek. Jl. Tunggul Pegagan, Kempek, Gempol, Cirebon 45161, West Java, Indonesia

²Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sebelas Maret. Jl. Ir. Sutami 36A Surakarta 57126, Central Java, Indonesia. Tel./fax. +62-271-663375, *email: suratman@staff.uns.ac.id

Manuscript received: 4 August 2021. Revision accepted: 21 September 2021.

Abstract. Wutsqa YU, Suratman, Sari SA. 2021. Detection of terpenoids and steroids in *Lindsaea obtusa* with thin layer chromatography. *Asian J Nat Prod Biochem* 19: 65-69. The diversity of ferns on this earth is very high, as well as the potential and benefits for human life, namely as ornamental plants, food material and medicines. The benefits of not all ferns are known because the lack of information on these ferns, one of ferns is *Lindsaea obtusa* J. Sm. Ex Hook. The phytochemical test is the first step in research on the search for new active compounds derived from natural ingredients. This study was conducted to determine the differences of terpenoid and steroid compounds profile in *L. obtusa* extract using Thin Layer Chromatography (TLC). Extraction of *L. obtusa* using maceration techniques with methanol and n-hexane solvents. The extraction results will be used for phytochemical screening using TLC technique with GF254 silica gel plate as a stationary phase and a mobile phase are chloroform: n-hexane (9: 1 v/v), n-hexane: ethyl acetate (4: 1 v/v), and methanol: ethyl acetate (5: 1 v/v). Detection of terpenoid and steroids using Lieberman Burchard reagent. The result shows that *L. obtusa* extract contains terpenoid and steroid compounds. Quantitatively, the secondary metabolites of *L. obtusa* extract were soluble in methanol solvent more than the compounds that were soluble in n-hexane solvent. However, qualitatively, steroid compounds that dissolve in n-hexane solvents have a more varied R_f value than terpenoids, especially in the mobile phase of chloroform: n-hexane (9: 1 v/v) and n-hexane: ethyl acetate (4: 1 v/v). *L. obtusa* contain more varied steroid compounds than terpenoid compounds.

Keywords: *Lindsaea obtusa*, phytochemistry screening, thin layer chromatography, terpenoids, steroids

INTRODUCTION

The diversity of ferns on this earth is very high. There are around 13,000 species of ferns on this earth, while the ferns in Indonesia, including the Malesiana area, are estimated to be 1250-1500 species (Machfira et al. 2016). The distribution of ferns is very large, as well as the potential and benefits that are high enough for human life, namely as ornamental plants, foodstuffs and medicines. The benefits of not all ferns are known because of the lack of information about the potentials of these ferns, one of which is the ferns *Lindsaea obtusa* J. Sm. Ex Hook.

Lindsaea obtusa is a terrestrial fern with 20-30 cm tall. These ferns have root fibers with a diameter of 1-2.5 mm, brown in color. *L. obtusa* stem is hard, stiff, and black with a length of 4-25 cm. The leaves of *L. obtusa* are folium compositum pinnatus leaves, and have black round petiolus. Leaflets are round or oval in shape, asymmetrical at the tip and base, smooth and green in color. *L. obtusa* has a round, brownish-green sorus located on the underside of the leaf. On one of leaf there are approximately 10-13 sorus. *L. obtusa* grows in forest shrubs especially in humid areas. This fern lives at an altitude of approximately 1000 meters above sea level (Dong et al. 2016). The true ferns *L. obtusa* live in temperate Asian regions such as Taiwan, and tropical climates such as Indonesia, parts of Java and Bali.

The phytochemical test is an important first step in research on the search for new active compounds derived from natural ingredients. This information can be used as

initial information for the synthesis of new drugs or to become prototypes of certain active compounds (Kruk et al. 2021). Terpenoids are also called isoprenoids, this is because the carbon skeleton is the same as the isoprene compound. Synthesis and accumulation of terpenoid compounds contribute to medicinal properties for the treatment respiratory inflammation, atopic dermatitis, arthritis, and neuroinflammation among various inflammatory diseases (Kim et al. 2020).

Steroids are lipids that do not have fatty acid groups and are not ester derivatives. Steroids are complex, fat-soluble organic molecules. This compound acts as a major component of cell tissue (Rashidinejad et al. 2021). Synthetic steroids that are often used are glucocorticosteroids, estrogen, methylprednisolone, corticosteroids, androgens, squalamine and hydrocortisone. These steroid compounds are commonly used for the treatment of diseases due to neurodegenerative disorder (Bansal and Singh, 2018). Ferns from the Equisetaceae, Osmundaceae, Lygodiaceae, Dennstaedtiaceae, Woodsidiaceae, Thelypteridaceae and Dryopteridaceae families also contain steroid compounds (Yokota et al. 2017).

The method that can be used for phytochemical screening is chromatography. Chromatography is the separation of a mixture of compounds in a sample based on differences in the interaction of the sample with the stationary and mobile phases. One of the chromatography methods that can be used for phytochemical screening is Thin Layer Chromatography (TLC). Separation in Thin

Layer Chromatography is based on differences in polarity between the sample and the solvent. The stationary phase is a solid that is applied flat in the form of glass or aluminum as a support. The mobile phase is a mixture of several liquids of different polarity. Research on the content of terpenoids and steroid in *L. obtusa* was very limited. Therefore this study was conducted to determine the difference of terpenoids and steroid profile in *L. obtusa* J. Sm. Ex Hook.

MATERIALS AND METHODS

Sample preparation

Sampling of *L. obtusa* was carried out using the cruise method, namely exploring the area in the KGPAA Mangkunagoro I Botanical Forest Park, Ngargoyoso, Karanganyar, Central Java, Indonesia which generally contained these ferns. Extraction process and phytochemistry screening was conducted in the Biology Laboratory, Faculty of Mathematics and Natural Sciences, Universitas Sebelas Maret, Surakarta, Indonesia.

The samples of *L. obtusa* ferns were separated from dirt or foreign materials and then cleaned with running water until clean. These sample dried until the fern has reduced its moisture content. After drying, the sample is mashed using blender until it becomes a powder and then weighed which is then used for the extraction process (Seremet et al. 2020).

Secondary metabolites extraction

The extraction of secondary metabolites was carried out by maceration method using methanol and n-hexane as solvents. 100 g of *L. obtusa* powder was macerated using methanol as a solvent with a ratio of 1 : 5 which was carried out for overnight. The same is done with the n-hexane solvent. The extraction process is carried out by soaking in a solvent and then filtered using filter paper. The solvent is evaporated with a rotary evaporator at 400C, a speed of 40 rpm and a pressure of 0.06-0.08 MPa until a thick extract is obtained for approximately 1 hour (Sati et al. 2019). The viscous extract is weighed and the yield is calculated against the weight of the initial simplicia. The formula for the percentage of extract yield according to Wahyuni and Widjanarko (2015) is :

$$\text{Yield Extract} = \left| \frac{\text{extract weight obtained (g)}}{\text{extracted weight (g)}} \right| \times 100\%$$

Thin layer chromatography (TLC) profile analysis

Chemical compound profile analysis was performed on each extract using the Thin Layer Chromatography (TLC) method. The n-hexane and methanol extracts were eluted using the same stationary and mobile phases. This was done to determine the results indicated by the presence of different color spots on TLC for each extract (Rubiyanto, 2013). The n-hexane and methanol extracts were spotted on the TLC plate with the silica gel GF254 as stationary phase and eluted using the mobile phase in the form of

chloroform: n-hexane (9:1 v/v), n-hexane: ethyl acetate (4:1 v/v) and methanol: ethyl acetate (5:1 v/v) in the developer vessel. GF254 silica gel plate was made with a width of 2 cm and a length of 9 cm and given an initial limit of 1 cm and an end limit of 0.5 cm. The limit of the solvent is below the line where the spots are. After the eluent reaches the finish line of elution, the plates are removed and dried.

Analysis of secondary metabolite compounds

The detection of terpenoid and steroid content from n-hexane extract and methanol extract was used by Lieberman Burchard spray reagent. The parameters were observed by the terpenoid and steroid compounds from *L. obtusa* and the Rf (Retardation Factor) value (Yin et al. 2017). In thin layer chromatography, the degree of retention is expressed as Rf which can be formulated:

$$\text{Retardation Factor (Rf)} = \left| \frac{\text{Distance of movement of the solute}}{\text{Distance of movement of the solvent}} \right|$$

Lieberman Burchard is a spotting reagent for detecting steroids and terpenoids. The Lieberman burchard spray reagent was prepared by mixing 5 ml of acetic acid anhydride with 5 ml of concentrated sulfuric acid, then adding this mixture to 50 ml of absolute ethanol. Each substance mixing was carried out by cooling. The application of this method to the TLC plate was sprayed with Lieberman Burchard reagent and then heated for 10 minutes at a temperature of 100°C. The presence of terpenoids is indicated by the appearance of a red violet color, while the presence of steroids is indicated by the appearance of a reddish brown color (Gummadi et al. 2021).

RESULTS AND DISCUSSION

Secondary metabolite extraction

Extraction of secondary metabolites from *L. obtusa* was carried out by maceration method using methanol and n-hexane as solvents. The results showed that the extraction with methanol as a solvent produced a higher yield than n-hexane (Table 1).

Extraction with methanol solvent resulted of 5.8 g so that the yield percentage was 5.8%. The extraction result with n-hexane solvent resulted of 0.7 g so that the yield percentage was 0.7%. The difference in yield percentage is due to differences in the solubility of the extracted compound in each solvent used, so that it can affect the yield and characteristics of the extracted chemical compound. The n-hexane solvent is intended to dissolve compounds that are semi-polar to nonpolar, while methanol is used to dissolve compounds that are more polar. Compounds can be said to be nonpolar compounds if they have bonds between atoms that have attractive ability to gain electrons together, resulting in a distant state where the electrons are shared equally. Meanwhile, if a compound has one atom capable of attracting electrons stronger than other atoms and the electrons from the bond will not be

used together equally, then the compound is called a polar compound (Tunega et al. 2020). These results are similar to those of (Azka and Abdullah, 2012) who showed that the secondary metabolites of water clover ferns *Marsilea crenata* most of them also dissolved in polar solvents with the yield of methanol extract of 11.98%, ethyl acetate extract of 1.37%, and chloroform extract of 0.31%. Methanol is a polar solvent while ethyl acetate and chloroform are semi polar and nonpolar solvents.

Thin layer chromatography profile analysis

Research shows that *L. obtusa* extract contains terpenoid and steroid compounds. This is indicated by a positive reaction to Liebermann-Burchard reagent (Table 2). The Liebermann-Burchard reaction was first described by Liebermann and then extensively developed by Burchard (Xiong et al. 2007). The Liebermann-Burchard reaction has been studied to determine the sterol content. Liebermann Burchard reagent produces a varies color response depending on the double bond system, other functional groups, and the presence of non-polar bonds (Xiong et al. 2002). Liebermann-Burchard is a spotting reagent for detecting steroids and terpenoids. This analysis was based on the ability of terpenoid and steroid compounds forming colour with concentrated H_2SO_4 in anhydride acetic acid (Parbuntari et al. 2018). Liebermann-Burchard reagent will show a red-violet stain which indicates the presence of terpenoid compounds, while the presence of steroids is indicated by the appearance of a reddish brown stain.

There are two fractions that give negative results on the detection of terpenoids with no visible red violet spots, namely in the mobile phase methanol extract of


chloroform: n-hexane (9: 1 v/v) and n-hexane extract in the mobile phase of methanol: ethyl acetate (5 : 1 v/v) (Table 2). In the detection of steroids, there were also two fractions that gave negative results with no visible reddish brown spots, namely the mobile phase methanol extract n-hexane: ethyl acetate (4: 1 v/v) and n-hexane extract in the mobile phase of methanol: ethyl. acetate (5: 1 v/v).

Table 3 shows that the Rf values for detection of terpenoid compounds in the mobile phase of chloroform: n-hexane (9: 1 v/v) n-hexane extract are found at Rf 0.82 and 0.93. The methanol extract in the mobile phase of n-hexane: ethyl acetate (4: 1 v/v) was found at Rf 0.90 and for n-hexane extract 0.42; 0.76; and 0.85. Methanol extract in the mobile phase of methanol: ethyl acetate (5: 1 v/v) was found at Rf 0.73 and 0.88. Meanwhile, the Rf value for the detection of steroid compounds in the mobile phase of chloroform: n-hexane (9: 1 v/v) methanol extract was found at Rf 0.93 and for n-hexane extract 0.49; 0.64; 0.73; 0.86; and 0.90. The n-hexane extract in the mobile phase of n-hexane: ethyl acetate (4: 1 v/v) was found at Rf 0.34; 0.48; 0.58; and 0.96. Methanol extract in the mobile phase of methanol: ethyl acetate (5: 1 v/v) was found at Rf 0.78. These results indicate that the n-hexane extract gave clearer results than the methanol extract, especially in the mobile phase of n-hexane: ethyl acetate (4: 1 v/v).

Table 1. Maceration result of *Lindsaea obtusa* extract

Solvent	Powder mass (g)	Extract mass (g)	Yield (%)
Methanol	100	5.8	5.8
N-hexana	100	0.7	0.7

Table 2. Thin Layer Chromatography Profile (TLC) of *Lindsaea obtusa* extract with Liebermann-Burchard reagent



Note: *KH = chloroform: n-hexane (9: 1 v/v); HE = n-hexane: ethyl acetate (4: 1 v/v); ME = methanol: ethyl acetate (5: 1 v/v). The yellow sign indicates the presence of a terpenoid compound and the orange sign indicates the presence of a steroid compound

Table 3. Retardation Factor (Rf) value of *Lindsaea obtusa* extract with Liebermann-Burchard reagent

Senyawa	KH		HE		ME	
	Methanol	N-Hexane	Methanol	N-Hexane	Methanol	N-Hexane
Terpenoid	-	0.82 0.92	0.9	0.42 0.76 0.85	0.73 0.88	-
Steroid	0.93	0.49 0.64 0.73 0.86 0.9	-	0.34 0.48 0.58 0.96	0.78	-

Note: *KH = chloroform: n-hexane (9: 1 v/v); HE = n-hexane: ethyl acetate (4: 1 v/v); ME = methanol: ethyl acetate (5: 1 v/v).

The Rf value of 0.90 indicates the existence of a terpenoid compound. This is also found in medicinal Herb *Hypochaeris radiata* using High Performance Thin Layer Chromatography (HPTLC) technique (Senguttuvan and Subramaniam, 2016). The results of the thin layer chromatography profile showed that red-violet spots were more visible in the n-hexane extract than in the methanol extract. This is consistent with the nonpolar secondary metabolites of terpenoids and steroids. Most of the terpenoid compounds also contain an -OH group so that the presence of a hydroxyl group substituent attached to the hydrocarbon chain can be attracted by semi-polar and even polar solvents (Stachowiak et al. 2020). So it can be concluded that *L. obtusa* positively contains secondary metabolites in the form of terpenoids and steroids. Other ferns such as *Azolla microphylla* also contains terpenoid and steroid compounds (Rashad, 2021).

In conclusion, *Lindsaea obtusa* J. Sm. ex Hook extract contains terpenoid and steroid compounds. Quantitatively, the secondary metabolites of *L. obtusa* extract that were soluble in methanol solvent were more than the compounds that were soluble in n-hexane solvent. However, qualitatively, steroid compounds that dissolve in n-hexane solvents have a more varied Rf value than terpenoids, especially in the mobile phase of chloroform: n-hexane (9: 1 v/v) and n-hexane: ethyl acetate (4: 1 v/v).

REFERENCES

- Azka A, Abdullah A. 2012. Aktivitas antioksidan dan komponen bioaktif semanggi air (*Marsilea crenata*). Jurnal Inovasi dan Kewirausahaan 1 (3): 152-158. [Indonesian]
- Bansal R, Singh R. 2018. Exploring the potential of natural and synthetic neuroprotective steroids against neurodegenerative disorders: A literature review. Med Res Rev 38 (4) : 1126-1158. DOI: 10.1002/med.21458
- Dong SY, Zuo ZY, Chao YS, Damas K, Sule B. 2016. New species of the fern genus *Lindsaea* (Lindsaeaceae) from New Guinea with notes on the phylogeny of *L. Sect. Synaphlebium*. PLoS One 11 (10): e0163686. DOI:10.1371/journal.pone.0163686
- Gummadi SB, Belvotagi VAR, Kommidi DR, Bobbala RK, Achanta VNAR, Sterner, O. 2021. Steroidal glycosides from *Caralluma pauciflora*. Phytochem Lett 41: 43-48. DOI: 10.1016/j.phytol.2020.10.016
- Kim T, Song B, Cho KS, Lee IS. 2020. Therapeutic potential of volatile terpenes and terpenoids from forests for inflammatory diseases. Int J Mol Sci 21 (6): 2187. DOI: 10.3390/ijms21062187
- Kruk A, Piwowarski JP, Pawłowska KA, Popowski D, Granica S. 2021. High molecular pyrogens present in plant extracts interfere with examinations of their immunomodulatory properties in vitro. Sci Rep 11 (1): 799. DOI: 10.1038/s41598-020-79579-2
- Machfira Y, Nery S, Fitmawati. 2016. Keanekaragaman Pteridaceae berdasarkan karakter morfologi dan fitokimia di Hutan PT. Chevron Pacific Indonesia (PT. CPI) Rumbai. Jurnal Riau Biologi 1 (2): 165-172. [Indonesian]
- Parbuntari H, Prestica Y, Gunawan R, Nurman MN, Adella F. 2018. Preliminary phytochemical screening (qualitative analysis) of cacao leaves (*Theobroma Cacao* L.). Eksakta 19 (2): 1-8. DOI: 10.24036/eksakta/vol19-iss2/142
- Rashad S. 2021. An overview on the aquatic fern *Azolla* spp. as a sustainable source of nutrients and bioactive compounds with resourceful applications. Egyptian J Aquat Biol Fish 25 (1): 775-782. DOI: 10.21608/ejabf.2021.150205
- Rashidinejad A, Marze S, Singh H. 2021. Lipid digestion and bioaccessibility of lipid-soluble compounds. In: Grundy MML, Wilde PJ (eds) Bioaccessibility and Digestibility of Lipids from Food. Springer, Cham. DOI: 10.1007/978-3-030-56909-9_10
- Sati P, Dhyani P, Bhatt ID, Pandey A. 2019. *Ginkgo biloba* flavonoid glycosides in antimicrobial perspective with reference to extraction method. J Trad Complm Med 9 (1): 15-23. DOI: 10.1016/j.jtcm.2017.10.003
- Senguttuvan J, Subramaniam P. 2016. HPTLC fingerprints of various secondary metabolites in the traditional medicinal herb *Hypochaeris radiata* L. J Bot 2016: 5429625. DOI: 10.1155/2016/5429625
- Seremet L, Nistor OV, Andronoiu DG, Mocanu GD, Barbu VV, Măidan A, Rudi L, Botez E. 2020. Development of several hybrid drying methods used to obtain red beetroot powder. Food Chem 310: 125637. DOI: 10.1016/j.foodchem.2019.125637
- Stachowiak W, Rzemieniecki T, Klejdysz T, Pernak J, Niemczak M. 2020. "Sweet" ionic liquids comprising the acesulfame anion - synthesis, physicochemical properties and antifeedant activity towards stored product insects. New J Chem 44 (17): 7017-7028.
- Tunega D, Gerzabek MH, Haberhauer G, Lischka H, Solc R, Aquino AJA. 2020. Adsorption process of polar and nonpolar compounds in a nanopore model of humic substances. Europe J Soil Sci 71 (5): 845-855. DOI: 10.1039/C9NJ06005G
- Wahyuni DT, Widjanarko SB. 2015. Pengaruh jenis pelarut dan lama ekstraksi terhadap ekstrak karotenoid labu kuning dengan metode gelombang ultrasonik. Jurnal Pangan dan Agroindustri 3 (2): 390-401. [Indonesia]
- Xiong Q, Ruan B, Whitby FG, Tuohy RP, Belanger TL, Kelley RL, Wilson WK, Schroepfer GJ Jr. 2002. A colorimetric assay for 7-dehydrocholesterol with potential application to screening for Smith-Lemli-Opitz syndrome. Chem Phys Lipids 115 (1-2):1 -15. DOI: 10.1016/s0009-3084(01)00205-5
- Xiong Q, Wilson WK, Pang J. 2007. The Liebermann-Burchard Reaction: Sulfonation, desaturation, and rearrangement of cholesterol in acid. Lipids 42: 87-96. DOI: 10.1007/s11745-006-3013-5
- Yin D, Guan Y, Gu H, Jia Y, Zhang Q. 2017. Polymerized high internal phase emulsion monolithic material: A novel stationary phase of thin layer chromatography. RSC Adv 7 (12): 7303-7309. DOI: 10.1039/C6RA27609A
- Yokota T, Ohnishi T, Shibata K, Asahina M, Nomura T, Fujita T, Ishizaki K, Kohchi T. 2017. Occurrence of brassinosteroids in non-flowering land plants, liverwort, moss, lycophyte and fern. Phytochem 136: 46-55. DOI: 10.1016/j.phytochem.2016.12.020.

Physicochemical composition of some new craft beers consumed in Maroua town from Far North region of Cameroon

M. DIGUIR¹, A. LAYA^{2,3,*}, H. WANGSO¹, J.P. BAYANG², B.B. KOUBALA^{1,3}

¹Department of Chemistry, Faculty of Science, University of Maroua. P.O. Box 814 Maroua, Cameroon

²Department of Biological Sciences, Faculty of Science, University of Maroua. P.O. Box 814 Maroua, Cameroon

³Department of Life and Earth Sciences, Higher Teacher's Training College of Maroua, University of Maroua. P.O. Box 55 Maroua, Cameroon

*email: laya.alphonse@yahoo.fr

Manuscript received: 21 June 2021. Revision accepted: 30 September 2021.

Abstract. Diguir M, Laya A, Wangso H, Bayang JP, Koukala BB. 2021. Physicochemical composition of some new craft beers consumed in Maroua town from Far North region of Cameroon. *Asian J Nat Prod Biochem* 19: 70-80. The work aimed to investigate the physicochemical composition of various craft beers (CB) and their distillates produced in Maroua town. The beers named "Bil-Bil", "Cochette" and "Furdu" were collected both morning and evening time. Dry matter, pH, titratable acid, total dissolved solids, proteins, sugars, carbohydrates, amino acids and alcohol content were evaluated. The results showed that local beers had a pH ranged from 4.06 to 4.22, 4.15 to 3.84 and 3.53 to 3.37 for "Furdu", "Cochette" and "Bil-bil", respectively. The alcohol varied between 5.40 and 6.92%, 1.08 and 5.10%, 1.73 and 2.88 for "Furdu", "Cochette" and "Bil-bil", respectively. The sugars ranged from 11.85 to 19.05 mg/mL for "Furdu", from 2.20 to 9.46 mg/mL for "Cochette" and from 9.92 to 19.78 mg/mL for "Bil-bil". For amino acids, the values varied from 3.40 to 7.37 mg /mL for "Furdu", 3.98 to 6.01 mg / mL for "Cochette" and 2.95 to 3.24 mg/mL for "Bil-Bil". Regarding protein, the values ranged from 0.80 to 0.85 mg/mL for these three CB. The distillates of CB collected evening showed high alcohol in fraction 1. Thus, these CB analysed can be promoted for alternative beer in Maroua town. Furthermore, the unsold CB can be distilled into ethanol.

Keywords: Alcohol, craft beers, distillate, Maroua town, physicochemical

INTRODUCTION

In Africa, cereals are the most widely cultivated and consumed as a staple food in many African households. These cereals are also used to prepare many fermented by-products. A local transformation unit of these cereals has been found in many areas of Cameroon, Republic of Central Africa and Chad (Djanan et al. 2002). The enterprises are mostly performed by women who recruit at least three workers (Maoura et al. 2002). In Cameroon, especially in the Northern part of Cameroon, various cereals such as sorghum, maize, yellow millet, S35, rice and fonio are processed into alcoholic and non-alcoholic beverages called locally beer or ancestor drink. These local beers produced are well appreciated by numerous people of Cameroon and other countries because these are a good source of nutrients such amino acids, carbohydrates, vitamins, minerals, phenolics, etc. (Maoura et al. 1999; Bamforth 2002; Lyumugabe et al. 2012; Cirimina et al. 2018). Because of their probiotics these local beers as known as therapeutic agents. The local beer is a beverage of significant historical and cultural importance. It is gaining popularity of craft beer and is of growing interest in several countries. It is also providing alternatives to mainstream beer production (Einfalt 2021). To date, craft beer is one of the fastest-growing alcoholic beverage industries throughout the world (Gómez-Corona et al. 2016).

These local beers are known as *Bil-Bil Furdu* and *Cochette*, the most widely produced and consumed.

According to the tribes, these beers have different name, "Bil-Bil" beer is known as "Muzum" in Guiziga; "Ouzomm" in Mada, "Zoum" in Mafa; "Balda" in Guidar; "Himi" in Moudang; "Yii" in Toupouri; "Tcheu" in Kapsiki "Mbolo" in Fali; "Mgba" in Laka; "Amgba" in Baya; "Do'di" in Dourou and "Coumouille" in Kera. However, "Furdu" beer is known as "Bazdltah" in Mada, "Pram" or "Bazla Babara" in Guiziga, "Mpedli" in Kapsiki. The "Cochette" beer is known as "Cochette" in almost all tribes but differs by the pronunciation except the name in Mada tribe, "Mohosso". The local name of *Bil-Bil* vary from one country to another as *Bili-Bili* in Cameroon and Chad, *Dolo* in Burkina-Faso, Mali and Senegal; *Tchapalo* in Ivory Coast, *Tchoukoutou* in Benin, *Pito* in Ghana, and *Impéké* in Burundi (Odunfa 1985; N'da et al. 1996; Kayodé et al. 2005). While, *Furdu* and *Cochette* are new local beers which are not documented in the literature. The energy is provided at 70% by alcohol and 30% by carbohydrates (Lariven and Rigal 2017). According to Djanan et al. (2002), one litre of *Bil-Bil* contributes significantly to the recovery of iron magnesium, manganese, phosphorus and calcium dietary allowance per day for an adult. Various studies of *Bil-Bil* beer are investigated, however, these parameters are absent for *Furdu* and *Cochette* beers. Setta et al. (2020) reported that traditional African fermented cereal-based beverages are potential probiotic carriers because of the probiotic *Lactobacillus* spp. and yeasts which are involved in the

fermentation of such products, and can be used as probiotic health benefits to the majority of African populations.

The main concerns of all the local beers is their consumption for one to two days and the next day, these beers become more acidic due to the breakdown of ethanol into acetic acid (oxidation reactions). Finally, the unsold local beers are usually thrown. These unsold beers can be used in the production of quality by-products in order to have added value and solve pollution problem.

Thus, the present study was aimed to determine the physicochemical composition and distil some craft beers named *Bil-Bil*, *Furdu* and *Cochette* in order to valorize the local beers mostly consumed in Maroua town of Cameroon.

MATERIALS AND METHODS

Survey

The survey was conducted in order to find out more about the micro-enterprises in charge of the production of local beer in the town of Maroua, region of the Far North of Cameroon. A preliminary survey was carried out in all streets of the city of Maroua. We identified four streets where they produced mostly local beers such as Pitoare, Pont-vert, Ouro-tchede and Palar. During this survey, several types of local beers are consumed in the town of Maroua. However, on the basis of the most consumed in the city, we have identified three (03) types of local beers: *Furdu*, *Cochette* and *Bil-Bil*. After this first step, we conducted a survey using direct interviews with the producers and consumers of these beers. We interviewed 150 producers and 200 consumers of *Bil-Bil* beer, 20 producers and 35 consumers of *Furdu* beer and 20 producers and 40 consumers of *Cochette* beer.

Samples

Samples of *Cochette*, *Furdu* and *Bil-Bil* beers were purchased through the various producers located in four streets (Pitoare, Pont-vert, Ouro-tchede and Palar). The

collected samples in the morning (start of the sale) and in the evening (end of the sale) from the suppliers and were introduced in suitable and sterile container and then labelled (Figure 1). Samples were transported to the Biochemistry Laboratory and Biological Chemistry of the University of Maroua (LabBBC). All samples were kept at 10°C before physico-chemical analyses and distillation.

Processing flow charity of different beers preparation

All the craft beers were produced by using different cereals shown in Figure 2.

Shortly, the cereal was soaked for 24h and washed prior for germination (2-3days). The germinated cereal was then sun-dried before milling and the malt flour was mixed with water and the mash mixture was decanted for 30min. The supernatant was heated for 2-3hr and its allowed to rest for overnight. Then, the second heating was done for 2-4 to obtain the sweet wort which was cooled for 3hr or more. Finally, the wet yeast was added and fermented overnight to obtain *bil-bil* beer. Figure 2.A shows the *Bil-Bil* beer preparation.

In brief, 7.0 kg of cereal was soaked for 1-2 days before sun-drying for 3hrs and milling in order to obtain flour (malt). Then, mixed with water before boiling at 95°C and the mash obtained was cooled at room temperature for 3hr before adding the wet yeast. The mixture was fermented for 10-12hrs and then filtered throughout the traditional filter tissue. The filtrate was poured in jar and named *Cochette* beer. The Figure 2.B shows different steps to produce *Cochette* beer.

Briefly, 3-7kg of cereal was wetted, cleaned or watered and germinated for three days and sun-dried before milling and the malt flour obtained was mixed with water, heated (95°C) and the mash obtained was cooled at room temperature for 4hr. Then, the mash was fermented for 72-100hr and then filtered throughout the traditional filter mesh. The product obtained was poured in jar and the *furdu* beer was ready for drinking. Figure 2.C is showing the summary methodology to produce *Furdu* beer.



Figure 1. Sample of different craft beers mostly produced and consumed in Maroua town, Cameroon. A. Bil-bil Beer, B. Cochette Beer, C. Furdu Beer




Figure 2. Flow charty preparation of beers in Maroua, Cameroon. A. Bil-bil Beer, B. Cochette Beer, C. Furdu Beer

Reagents and chemicals

Bovine Serum Albumin (BSA), Ninhydrin, Alanine, glucose, and 3,5-Dinitrosalicylic Acid (DNSA). Sulfuric acid, Ethanol, Chloridric acid Acetic acid, Nitrochromicacid, Iodic of potassium blue of coomassie, Sodium of thiosulfate, and all other solvents were obtained from Sigma-Aldrich (Iowa, USA) and Fisher commercial source (New Jersey, USA).

Kinetics of distillation

Sample of beer was collected both in the morning and in the evening from different streets (Pitoare, Pont vert, Ouro tchede and Palar) of Maroua town. The kinetics of distillation was evaluated as affected by the time and temperature. 400 mL of sample was introduced in 500mL rotavapor flash and it was allowed to heat until boiling and at the end two fractions of distillates were collected for analyses.

Physicochemical analyses

Total dissolved solids (TDS), Dry matter content (DM), pH
TDS was determined in triplicate by using TDS instrument and the values were expressed in ppm, however, dry matter content and pH were evaluated as described by AOAC (2000) method.

Titrateable acidity

Titrateable acidity was determined as described by Ranganna (1979). The values were expressed in milligram equivalent oxalic acid per 100 grams of sample beer.

Evaluation of reducing sugars content

Reducing sugars content was determined using the dinitrosalicylic acid as described by Miller (1972). The results were expressed in mg per mL of sample beer.

Evaluation of total carbohydrates content

Total carbohydrates content was determined as described by Dubois et al. (1956). The results were expressed in mg per mL of sample beer.

Determination of total amino acids, proteins and alcohol content

Total amino acids content was determined using ninhydrin (1 %) reagent as described by Michel and Hannequart (1968). The results were expressed in mg of amino acids per mL of sample beer.

Total soluble proteins content was determined using Bradford (1976) method. The results were expressed in mg of BSA per mL of beer sample.

The content of alcohol in sample beer was determined as described by Caputi et al. (1968).

Statistical analyses

SPSS 20.0 Statistical Package for Windows (SPSS Inc., Chicago, IL, USA) was used to perform all statistical analysis. Experiments were done in triplicate and one-way analysis of variance (ANOVA) was performed. Tukey's (HSD) test was used to determine any significant difference between different beers and the significance was accepted at level $p < 0.05$. The results were expressed as means \pm standard deviation.

RESULTS AND DISCUSSION

Sociodemographic profile of producers and consumers of local beers

The results of survey revealed that women aged between 20 and 40 represent the majority of the brewers (Table 1). They are predominantly catholic Christians, followed by Protestants and a few animists for all beers. Most of women are married ($\geq 60\%$), followed by widowed, single or divorced. Their education level varies from primary to secondary school and illiterate as well (Table 1). In Moundou, Tchad, similar results were reported by Djitod (2002) on *Bil-bil* beer who found that the majority of producers are catholic Christians (87%), aged between 26 and 35 years. Also, they are married women (65%), divorced (15%), widows (13%) and single (7%). The producers of local beers in Maroua town were mainly women suggested that men are only the consumers that is consistent with the Northern tradition of Cameroon.

Results also showed that the youngest people aged between 20 and 40 years (75%) are mainly consume *Bil-Bil*

beer (Table 1). Also, it found that *furdu* is more consumed by Protestants (57.7%) than others. However, *Bil-bil* (52%) and *Cochette* (50%) are mostly consumed by catholic Christian (Table 1). Then, other Christians such as Pentecostist, Adventist, Presbyterians and Muslims are consumed all types of beers. The consumers are predominantly married ($\geq 50\%$) for all the beers than widowed, single and divorced. They have a level of education mostly between secondary and higher education and a few proportion of primary and illiterate (Table 1).

Furdu and *Cochette* beers are mostly consumed by people with a primary level of education. Pupils (32%) and students (27.5%) are the consumers of *Bil-Bil* beer. *Furdu* and *Cochette* are mostly consumed by the elder person aged between 40 and 60 years with a percentage of 63% and 45%, respectively, followed by the youngest people who aged between 20 and 39 years (Table 1). Also, according to our survey, low-income men and married people, unemployed and rich people are consumed the local beers in different streets of Maroua town every day from the morning (5 a.m.) to the evening (8 p.m.). The consumption of all these craft beers may be due to their traditional value, interested in tasting (Einfalt 2021), therapeutic value and lower cost than the commercial beers (modern beer). Fact, certain untypical flavors in craft beers even have the potential of being perceived by the consumer as having higher quality compared to commercial beer (Ascher 2012).

Different cereals used for production of local beers in Maroua town

The results of survey revealed that various cereals are used to produce local beers (*Bil-Bil*, *Furdu* and *Cochette*) in Maroua town. Cereals mostly used are maize, rice, fonio and sorghum. Rice and white sorghum were the main cereal for *Cochette* (40%) and for *Furdu* beer, it was white sorghum (85%) (Figure 3). While, *Bil-Bil* was the mixture or all of these different cereals. The mixture of red and yellow sorghum (46%) followed by red and white sorghum (38%), red sorghum, S35 and red-yellow sorghum and S35 were the most used by *Bil-Bil* producers, respectively (Figure 3). The present results are consistent with the results found by Charles et al. (2018), who reported that in the North Cameroon, red sorghum ("djagari") was the main cereal for the production of *Bil-Bil* beer. The results of the survey regarding producers show that *Bil-Bil* beer which produced with the mixture of red sorghum exhibited a better drinking though beer of red sorghum was absent in Maroua town, while, yellow and white sorghum were the best cereal for production of *Bil-Bil* beer (Figure 3).

Therefore, many factors can affect the quality of these local beers such as quality of heat during preparation (33%), type and germination of grain (18%) and other factors (Figure 4). These are consistent with the results reported by Olaniran et al. (2017) who stated that the quality of the final product (beer) is influenced by several variables, such as the quality of the raw material, type of malting, the applied preparation method of the wort, type of hops, type and quality of yeast, fermentation time, maturation, and pasteurization and filtration, among others.

Table 1. Results of survey (%) of the local beers collected from consumers and producers from different street of Maroua town. Values are rated answer concerning the consumption or production of *Bil-Bil*, *Cochette* or *Furdu* beers.

Parameter		<i>Bil-Bil</i>		<i>Cochette</i>		<i>Furdu</i>	
		Consumers	Producers	Consumers	Producers	Consumers	Producers
School level	Primary	22.5	39	50	60	57.1	50
	Secondary	32	41	20	20	22.9	20
	Higher	27.5	0	12.5	0	10	0
	Illiterate	10	20	17.5	20	10	30
Matrimonial status	Married	50	63	60	75	69	50
	Single	40	14	10	15	11	20
	Widow	4	20	17.5	10	20	30
	Divorced	6	3	12.5	0	0	0
Religion	Animist	4	7	12.5	15	22.9	20
	Protestant	33	13	30	5	57.1	0
	Presbyterian	1.5	0	0	0	0	0
	Muslim	3	0	0	0	0	0
	Pentecost	6.5	0	0	0	0	0
	Catholic	52	80	50	80	20	80
	Adventist	0	0	7.5	0	22.9	0
Age	[0-20[10	3	20	5	8.5	10
	[20-40[75	77	35	50	28.5	60
	[40-60[15	20	45	45	63	30




Figure 3. Proportion of the various combination of cereals used according to producers for craft brewery. Ri + WS = Rice + White Sorghum; Ri + F = Rice + Fonio; Ri = Rice, WS = White Sorghum, YS = Yellow Sorghum; M = Maize; RS = Red Sorghum; RS+YS = Red Sorghum + Yellow Sorghum; ReS + WS = Red Sorghum + White Sorghum; ReS+YS+S35 = Red Sorghum + Yellow Sorghum + S35.




Figure 4. Factors influencing the craft brewery.

The influence of time and temperature on the volume of distillate

Figure 5.A-B showed the evolution of distillation of beers collected both morning and evening. The curve (Figure 5.A) showed many stages during the distillation. The curve showed that in the start time, all craft beers had a significant amount of alcohol. According to Anderson (2012) in a binary, tertiary or quaternary non-azeotropic mixture, the most volatile constituents are those whose boiling point is the lowest at constant pressure, which qualified as light at 1atm, and the least volatile constituent is the one with the highest boiling temperature at constant pressure qualified as heavy at 1atm. From the point of boiling temperature of methanol (64°C), ethanol (78°C), propan-2-ol (82°C), propan-1-ol (97°C) of water (100°C) and other products resulting from oxidation such as acetone (56°C). Samples CPiM, BPiM, CPvM, FHoM and BPvM for those collected in the morning (Figure 4.A) and BPvE, FHoE, CPvE, and BPvE for those of the evening (Figure 5.B) and specially at different time intervals lets us believe in their presence. Other samples FPvM (71°C) and BPiM (72°C) for those from the morning and FPvE (75°C) and BPiE (73°C) for those from the evening and their distillation time at a value of maximum boiling temperature strictly lower than that of ethanol is what also leads us to believe in an abundance of methanol, acetone and others.

These differences are due to the raw materials used, to the brewing methodology and the time and period of inoculation of the yeasts to obtain a perfect fermentation of alcohol. Fact, at temperature values strictly below and above the boiling point of ethanol there is no more ethanol to distillate in the local beer but many alcohols and other mixed products. Higher alcohols (2-methyl-1-propanol, 2-methyl-1-butanol and 3-methyl-1-butanol) derived from the catabolism of amino acids and sugars via the Ehrlich pathway are common in fermented and distilled beverages (Hazelwood et al. 2008; Nascimento et al. 2008). Gas chromatography (GC-FID) and GC-MS method for the detection and quantification of these different alcohols and other compounds in mixture has not been carried out in the present study, so we believe that there are many other compounds may be present. Recent works show many volatile compounds namely aldehydes, esters, alcohols and acids have been identified by using different chromatography method in cassava spirit and traditional sorghum beer (Attchelouwa et al. 2020; Coelho et al. 2020) which derived from the fermentative metabolism of yeast and can be further concentrated during the distillation.

Figure 5.C-D showed the evolution of the volume of distillate of various beers collected at different times (morning and evening). We noticed that the increase in volume at different times and temperatures varied in respect with the types of beers. All samples of beers from the morning to the evening had maximum boiling temperature with values greater than or equal to 78°C ($T \geq 78^\circ\text{C}$) showed the highest value in terms of volume, 15.9

mL (BPvM), 15.8mL (CPiE) for low value and 20.85mL (CPiM), 21.3mL (BPvE) for high values. Almost ethanol content of sample beer was collected in different fractions, with distillation efficiency close to 100% (Coelho et al. 2020). On the other hand, we did not collect all the ethanol which was found in some samples, those whose boiling point was strictly below 78°C. We think it might take a longer time than we defined or at least there was not enough ethanol but other alcohols in those initial samples of beers.

Therefore, we obtained the smallest values in terms of volume with the value of 13.65 mL (BPiM) and 15.8 mL (FPvM) for the morning sample (Figure 4.C) and 15.8 mL (CPiE), 16.84 mL (BPiE) and 18.55 mL (FPvE) for those of the evening (Figure 4.D).

In general, the volume of distillate from the evening was significantly greater than those from the morning. This variation in volume between the samples collected in the morning and those from the evening may be due to the influence of the brewing technology, the raw material used (Humia et al. 2019) and the quality of yeast (Einfalt 2021) as well as time of maturation which have an effect on the fermentation of the different beers. This difference proves sufficiently that regardless of the temperature values greater than or equal to, less than or equal to the boiling temperature of ethanol $50 \leq T_b \leq 90$, we do not only collect ethanol and other alcohols mixed but also a significant amount of water. In short, at high temperature of around 100 °C, the distillate to be recovered consist of maximum water content and very low alcohol content.

Variation of some physicochemical characteristics in local beers

The physicochemical characteristics of *Bil-bil*, *Furdu* and *Cochette* are showed in Table 2. Statistical analysis showed a significant ($p < 0.05$) difference among samples (Table 2). The results showed that TDS of all the investigated beers vary among them, however, *Furdu* beer has the highest TDS value (1688-1978 ppm), while the lowest was showed by *Cochette* (513-1409 ppm). The present results are in agreement with those reported by Coulibaly et al. (2020), who found in traditional sorghum beer (*Tchapalo*) in Ivory Coast a value of 15.60 °B which was variable than our values. This difference must be linked to the methodology using and the type of cereal to produce local beer. While, dry matter content of the beers varied between 4.24 to 6.98% (*Bil-Bil*), 2.83 to 6.96 % (*Furdu*), and 5.17 to 8.06 % (*Cochette*). Compared to the result obtained by Amane et al. (2005), the local beers investigated had a similar dry matter (7.87 %) of Sorghum *Tchapalo* produced in Ivory Coast. Similar results were also obtained by Chevauss-Agnes et al. (1976) who found that *Ambga* beer had 7.2 % of dry matter. These differences may be due to practices of production and the variety of cereals used in the processing of these beers.




Figure 5. The distillation temperature and distillate volume according to the time of the local beers collected. A-B. Evolution of the distillation temperature according to the time of local beers collected (A. collected morning, B. collected evening). C-D. Increase in distillate volume according to the time of the local beers collected (C. collected morning, D. collected evening). Note: FHo: *Furdu* beer collected in *Ouro tchéde* street; FPv: *Furdu* beer collected in *Pont vert* street, CPv: *Cochette* beer collected in *Pont vert* street, CPi: *Cochette* beer collected in *Pitoaré* street, BPv: *Bil-Bil* beer collected in *Pont vert* street, BPa: *Bil-Bil* beer collected in *Palar* street, BPi: *Bil-Bil* beer collected in *Pitoaré* street. Those beers were collected in the evening (E).

The pH value of different beers varied among them as shown in Table 2. It varied from 3.53 to 3.37 (*Bil-Bil*), 3.84 to 4.15 for "Cochette" and 4.06 to 4.22 for *furdu* beer. Compared to the values obtained by Roger et al. (2013) in *Amgba* beer (pH 2.5), *Tchapalo* (pH 3.4) by Aka et al. (2008), *Dolo* (pH 3.5) by Abdoul-latif et al. (2013), *Pito* (pH 3.66) by Fadahunsi et al. (2013) and *Burukutu* (pH 3.94) by Eze et al. (2011) and sorghum safari (*Tchapalo*) (pH 3.5 to 4.5) by Coulibaly et al. (2020), all produced in Cameroon, Ivory Coast, Burkina-Faso, Nigeria and Ghana, respectively, our investigated beers had similar values. Also, it showed that all our local beers had similar pH value both in the morning and in the evening, which is in accordance with those reported by Maoura et al. (2006) in Chad, where the pH value of *Bil-Bil* had not changed during the alcoholic fermented process. However, the results revealed that all the sample beers had acidic pH range which can explained the fact that acidic medium is one of the main condition factors for the developing of yeast without any modification of pH (Maoura et al. 2006).

The pH values of our local beers were within the desirable parameters, generally between 3.8 and 4.7, protecting the product against pathogens as reported by Suzuki et al. (2006). The difference in pH value may be attributed to the processing methods, quality of raw material and measure or quantification (Coulibaly et al. 2011). All values of pH of our different local beers produced in Maroua town is within the value of CODEX (2005), which reported that beer with a pH lower than 4.5 could be the good quality for the consumers.

The titratable acid of the local beers varied also significantly among samples ($p < 0.05$) (Table 2). *Bil-Bil* had the highest value (4.89-6.25 mg/mL), however *Furdu* had a value of 3.04-4.58 mg/mL, while *Cochette* had the lowest value (2.31-3.40 mg/mL). The values of titratable acid found in *Bil-Bil* beer are lower to the value (0.06mg/mL) reported by Coulibaly et al. (2020) in sorghum safari (*Tchapalo*). Similarly, these values are lower than those reported by Ourega et al. (2015), who found that plantain beer had a titratable acid content ranged

from 0.08 to 0.9.3 mg/mL suggesting that the raw material had influence on the beers.

For the alcohol content, results summarized in Table 2 revealed that the alcohol value varied significantly among beers ($P < 0.05$). The sample of *Furdu* beer had the contents varied between 5.40 and 6.92 %, while it folds from 1.08 to 5.10 % for *Cochette* and between 1.73 and 2.88 % for *Bil-Bil*. The *frudu* beer had alcoholic value higher than *Cochette* and *Bil-Bil* suggesting high flavour and aroma compounds (Einfalt 2021). However, the alcohol content in *Bil-Bil* beer was significantly higher than the value reported by Muyanja et al. (2010) on *Pito* (3.09 %), however lower than *Tchapalo* (5.22 %) processing with sorghum in Kenya, Nigeria and Ivory Coast, respectively. The rice and white millet used for the processing of *Furdu* (6.16%) and *Cochette* (3.09 %) had higher alcohol content than *Bil-Bil* (Table 2). These differences observed in terms of alcohol content are due to the types or quality of yeasts used to ferment. These are consistent with the results of Einfalt (2021) who found that the alcohol content varied significantly with the type of yeast used for beer fermentation. However, the alcohol levels obtained in our different beers are obviously within the range of the French legal classification of types of beers ranging from table beer to special beers brochures of brewers of France (Hencké 2000).

Reducing sugars content of *Furdu* varied between 11.85 and 19.05 mg/mL, while it fold from 2.20 to 9.46 mg/mL for *Cochette* and 9.92 to 19.78 mg/mL for *Bil-Bil* (Table 3). The present results showed that *Bil-Bil* beers had higher content of reducing sugars than *Tchapalo* (0.4 mg/mL) from Ivory Coast (Amane et al. 2005). This difference may be due to the reducing sugar content of sorghum used to process the local beers. Also, the contents of total carbohydrates varied among these beers (Table 3). The carbohydrates content in *Furdu* beer fold from 16.05 to

46.35 mg/mL, while in *Cochette* beer, it varied between 15.39 mg/mL and 42.53 mg/mL and for *Bil-Bil* the values varied from 12.12 to 168.53 mg/mL. It revealed that *Bil-bil* beer had lower and higher value of carbohydrates than that obtained in *Furdu* and *Cochette*, respectively. Cortacero-Ramírez et al. (2003) stated that beer should contain between 3.3 to 4.4 g/100 of carbohydrates. However, value is higher than that reported by Chevassus-Agnes et al. (1976) in *Ambga* and *Affouk* beers (19 mg/mL) consumed in the North region of Cameroon. The differences in carbohydrate values are linked to the variety of cereals and their technologies used for processing.

The values of soluble amino acids and total soluble proteins was not found varied significantly ($p > 0.05$) among the samples (Table 3). The soluble amino acids amount of the local beers were followed, *Furdu* had 3.40 to 7.37 mg/mL, *Cochette* contains 3.98 to 6.01 mg/mL, while *Bil-Bil* had 2.95 to 3.24 mg/mL. However, *Furdu* beer had the highest amino acid contents than *Cochette* and *Bil-Bil*. Considering the total proteins, local beers had high values ranged from 0.80 to 0.85 mg/mL which was higher than those obtained in *Kaffir* beer (0.05 mg/mL) (Busson et al. 1970), in *Ambga* (0.07 mg/mL) (Chevassus-Agnes et al. 1976) (Table 3). For Ferreira and Guido (2018), the contents of soluble amino acids determine the quality of beer suggesting the good quality of our sample beer analyzed.

The alcohol contents of distillate fractions from the morning and evening samples showed that the alcohol contents varied significantly ($p < 0.05$) among different samples of beers named FHoE, FPvE, CPiE, BPvE, and BPaE, however there is no significant difference among FHoM, FPvM, CPvM, CPiM, BiPvM and BPiE (Figure 6). The values of alcohol content revealed that the first fraction (F1) of the distillate had higher alcohol content than the second fraction (F2) (Figure 6).

Table 2. Total dissolved solids (TDS), pH, dry matter, titratable acid (TA) and alcohol contents of local beers (*Furdu*, *Cochette*, *Bil-Bil*) collected in the morning and evening from various streets in Maroua town.

Sample	TDS (ppm)	pH	Dry matter (g/100g)	T A (mg/mL)	Alcohol contents (g/100 mL)
FHoM	1771 ± 70 ^b	4.06 ± 0.03 ^c	6.96 ± 0.22 ^b	3.04 ± 0.19 ⁱ	5.38 ± 0.13 ^b
FHoE	1688 ± 23 ^c	4.06 ± 0.04 ^c	2.83 ± 0.08 ^h	3.80 ± 0.06 ^g	6.92 ± 0.28 ^a
FPvM	1939 ± 28 ^a	4.04 ± 0.02 ^d	6.96 ± 0.22 ^b	4.58 ± 0.32 ^e	5.47 ± 0.26 ^b
FPvE	1978 ± 20 ^a	4.22 ± 0.03 ^a	6.61 ± 0.32 ^{bc}	4.21 ± 0.06 ^f	5.40 ± 0.15 ^b
CPvM	1409 ± 26 ^f	4.15 ± 0.02 ^b	8.06 ± 0.08 ^a	2.63 ± 0.13 ^j	1.08 ± 0.23 ^h
CPvE	1310 ± 33 ^g	4.13 ± 0.06 ^{bc}	5.84 ± 0.08 ^c	3.40 ± 0.06 ^h	4.71 ± 0.21 ^{cd}
CPiM	625 ± 34 ^h	3.84 ± 0.03 ^c	6.09 ± 0.24 ^d	2.72 ± 0.13 ^j	4.31 ± 0.31 ^d
CPiE	513 ± 28 ⁱ	3.80 ± 0.03 ^e	5.17 ± 0.89 ^f	2.31 ± 0.06 ^k	5.10 ± 0.25 ^e
BPiM	1442 ± 18 ^e	3.46 ± 0.02 ^g	6.98 ± 0.37 ^b	5.07 ± 0.26 ^d	2.55 ± 0.25 ^f
BPiE	1434 ± 11 ^{ef}	3.45 ± 0.05 ^g	5.42 ± 0.44 ^{ef}	5.84 ± 0.32 ^{bc}	2.90 ± 0.23 ^e
BPaM	1495 ± 24 ^d	3.44 ± 0.02 ^g	4.88 ± 0.10 ^f	4.89 ± 0.26 ^{de}	1.73 ± 0.13 ^g
BPaE	1647 ± 29 ^e	3.53 ± 0.03 ^f	6.31 ± 0.07 ^c	5.66 ± 0.06 ^c	1.82 ± 0.26 ^g
BPiM	1483 ± 34 ^{de}	3.42 ± 0.04 ^{gh}	4.24 ± 0.09 ^g	6.25 ± 0.13 ^a	1.92 ± 0.13 ^g
BPiE	1441 ± 11 ^{ef}	3.37 ± 0.02 ^h	5.95 ± 0.38 ^{de}	5.93 ± 0.06 ^b	2.88 ± 0.21 ^{ef}

Values are means ± standard deviation of three replicates ($n = 3$). In the same column, values followed by different superscript letters are significantly different ($p < 0.05$). Note: FHo: *Furdu* beer collected in *Ouro tchédé* street; FPv: *Furdu* beer collected in *Pont vert* street; CPv: *Cochette* beer collected in *Pont vert* street; CPi: *Cochette* beer collected in *Pitoaré* street; BPv: *Bil-Bil* beer collected in *Pont vert* street; BPa: *Bil-Bil* beer collected in *Palar* street; BPi: *Bil-Bil* beer collected in *Pitoaré* street. Those beers were collected in the morning (M) or evening (E).

Table 3: Reducing sugars, total amino acids, total carbohydrates, and total proteins of local beers (*Furdu*, *Cochette*, *Bil-Bil*) collected in the morning and evening from various streets in Maroua town. The values are expressed in mg/mL of sample.

Sample	Reducing sugars	Carbohydrates	Total protein	Amino acids
FHoM	16.35 ± 0.45 ^d	16.05 ± 0.27 ^b	0.80 ± 0.05 ^b	4.29 ± 0.19 ^c
FHoE	17.33 ± 0.63 ^c	37.75 ± 0.99 ^d	0.83 ± 0.03 ^{ab}	3.40 ± 0.13 ^e
FPvM	11.85 ± 1.02 ^f	46.35 ± 1.14 ^b	0.84 ± 0.02 ^a	6.00 ± 0.16 ^a
FPvE	19.05 ± 0.63 ^a	44.71 ± 1.50 ^{bc}	0.83 ± 0.02 ^{ab}	7.37 ± 0.15 ^a
CPvM	8.30 ± 0.54 ^h	35.97 ± 1.17 ^d	0.83 ± 0.06 ^{ab}	5.83 ± 0.17 ^b
CPvE	9.46 ± 0.37 ^{gh}	42.53 ± 0.63 ^c	0.84 ± 0.05 ^a	3.98 ± 0.15 ^d
CPiM	2.20 ± 0.58 ⁱ	28.34 ± 0.89 ^e	0.83 ± 0.04 ^{ab}	6.01 ± 0.07 ^b
CPiE	5.56 ± 0.22 ⁱ	15.39 ± 0.55 ^h	0.83 ± 0.04 ^{ab}	4.44 ± 0.09 ^c
BPvM	13.06 ± 0.65 ^e	19.00 ± 0.80 ^g	0.81 ± 0.02 ^b	3.15 ± 0.17 ^f
BPvE	9.92 ± 0.41 ^g	12.12 ± 0.69 ⁱ	0.81 ± 0.02 ^b	3.03 ± 0.13 ^{gf}
BPaM	17.51 ± 0.84 ^{bc}	18.81 ± 0.43 ^g	0.83 ± 0.04 ^{ab}	3.02 ± 0.16 ^{gf}
BPaE	16.32 ± 0.71 ^d	168.53 ± 1.52 ^a	0.83 ± 0.02 ^{ab}	3.01 ± 0.04 ^g
BPiM	18.02 ± 0.48 ^b	24.49 ± 0.79 ^f	0.85 ± 0.03 ^a	2.95 ± 0.08 ^g
BPiE	19.78 ± 0.58 ^a	27.80 ± 1.04 ^e	0.80 ± 0.02 ^b	3.24 ± 0.10 ^{ef}

Values are means ± standard deviation of three replicates (n= 3). In the same column, values followed by different superscript letters are significantly different (p < 0.05). Note: FHo: *Furdu* beer collected in *Ouro tchéde* street; FPv: *Furdu* beer collected in *Pont vert* street; CPv: *Cochette* beer collected in *Pont vert* street; CPi: *Cochette* beer collected in *Pitoaré* street; BPv: *Bil-Bil* beer collected in *Pont vert* street; BPa: *Bil-Bil* beer collected in *Palar* street; BPi: *Bil-Bil* beer collected in *Pitoaré* street. Those beers were collected in the morning (M) or evening (E).




Figure 5. Alcohol contents (mL/100 mL) of distillate obtained from local beers (*Furdu*, *Cochette*, *Bil-Bil*) collected in the morning and evening from several streets in Maroua town, Cameroon. Note: FHo: *Furdu* beer collected in *Ouro tchéde* street; FPv: *Furdu* beer collected in *Pont vert* street; CPv: *Cochette* beer collected in *Pont vert* street; CPi: *Cochette* beer collected in *Pitoaré* street; BPv: *Bil-Bil* beer collected in *Pont vert* street; BPa: *Bil-Bil* beer collected in *Palar* street; BPi: *Bil-Bil* beer collected in *Pitoaré* street. Those beers were collected in the morning (M) or evening (E).

This result may be due to the fact that at the start of the distillation, a strong alcohol type has been collected with very strong odor with a sour taste and a characteristic aroma which relatively increased the alcohol content (Aka et al. 2008). On the other hand, at the end of the distillation, we observed an increase in temperature and consequently an alteration of the distillate and other volatile compounds which decreased the alcohol content by water interference. In general, ethanol was the main constituent of distillate, however this does not characterize the quality of the distillates. As indicated by Huot and Roy (1999), other flavors from many organic substances such as esters,

aldehydes and other alcoholic compounds must also interfere in the alcoholic degree of the distillates. These compounds may influence the quantity and quality of distilled ethanol. An ethyl acetate which had also been found in cassava spirits indicated that the distilled beverages such as whiskey, rum and Cachaça affected the quality of them (Coelho et al. 2020).

At less significant values, samples of distillates collected in the evening are richer in alcohol than those obtained in the morning, suggesting that this fraction was rich in flavor compounds. Fact, the principal high alcohol in alcoholic beverages include 1-propanol, 2-methyl-1-

propanol (isobutanol), 3-methyl-1-butanol (isoamylol) and 2-phenylethanol (Zhou et al. 2020). Likewise, some sample saw their second fraction more concentrated than the first one (Figure 6). This difference could be due specially to the yeast inoculation time (Einfalt 2021) and the fermentation time because early in the morning some local beers had not reached their fermentation or maturation threshold and can affect the quality of the distillate. Fact, the fermentation process develops with an increase in the alcohol level (Périsse et al. 1959).

In conclusion, the aim of this present study was to investigate the physicochemical characteristics of various local beers produced and consumed in Maroua town as well as the types of cereals used to produce these craft beers. Various cereals (red sorghum, white sorghum, maize, yellow millet, S35, rice and fonio, etc.) are the mainly used to process different local beers. The volume of distillate increased by time and depend on one beers to another. The sample of BPiM and CPiM had the lowest and the highest temperature values during distillation, respectively. *Furdu* beer had the highest TDS value (1688-1978 ppm), pH value (4.06-4.22) and high level of alcohol (5.40-6.92 %). However, *Cochette* had the highest level of dry matter (5.17-8.06 %). *Bil-Bil* had the highest titratable acid value (4.89 mg/mL-6.25 mg/mL). The distillate of the local beers had high alcohol levels and varied from beer to another and one fraction to another, with the values varied between 30 and 58.03%. These results revealed that various cereals would be suitable as raw materials for the production of quality beers such as white beers, beers, and spirits and provide a basis to foster the production of value-added products from these cereals. This study shows that the local beers are rich in nutriment and could offer a greater source of benefit to consumers.

REFERENCES

- Abdoul-Latif FM, Bassolé IH, Dicko MH. 2013. Proximate composition of traditional local sorghum beer "dolo" manufactured in Ouagadougou. *Afr J Biotechnol* 12(13) : 1517-1522.
- Aka S, Djeni NT, N'guessan KF, Yao KC, Dje KM. 2008. Variabilité des propriétés physico-chimiques et dénombrement de la flore fermentaire du tchapalo, une bière traditionnelle de sorgho en Côte d'Ivoire. *Afr Sci* 04 : 2274-286. DOI: 10.4314/afsci.v4i2.61688.
- Amame ND, Assidjo NE, Gbongue MA, Bohoussou K, Cardot P. 2006. Caractérisation physico-chimique d'une bière traditionnelle ouest africaine: Le Tchapalo bière traditionnelle tchadienne. *Agron Afr* 17 (2) : 143-152. DOI: 10.4314/aga.v17i2.1665
- Anderson NG. 2012. Practical Process Research and Development A guide for Organic Chemists. 2nd ed. Springer, New York.
- AOAC. 2000. Official Methods of Analysis of the Association of Official Analytical Chemists International 17th Ed. Published by the Association of Official Analytical Chemists International, Suite 400 2200 Wilson Boulevard, Arlington, Virginia, USA.
- Ascher B. 2012. Global beer-the road to monopoly. *Am Antitrust Instit*, Washington.
- Atchelouwa CK, N'guessan FK, Marcotte S, Amoikon TL, Charmel M, Dje MK. 2020. Characterisation of volatile compounds associated to sensory changes during the storage of traditional sorghum beer by HS-GC/FID and SPME-GC/MS. *J Agric Food Res* DOI: 10.1016/j.jafr.2020.100088
- Bamforth CW. 2002. Nutritional aspects of beer-a review. *Nutr Res* 22: 227-237. DOI : 10.1016/S0271-5317(01)00360-8
- Bradford MM. 1976. A rapid and sensitive Method for the Quantification of microgram quantities of protein utilizing the principe of protein-due binding. *Anal Biochem* 72: 248-254. DOI: 10.1006/abio.1976.9999
- Busson F, Jardin C, Wu Leung WT. 1970. Table de composition des aliments à l'usage de l'Afrique.
- Caputi A, Ueda M, Brown T. 1968. Spectrophotometric determination of ethanol in wine. *Am J Enol Vitic* 68 (2): 188-90.
- Charles T, Jong NE, Robert N. 2018. Production du Bili-Bili, bière traditionnelle de sorgho du Nord Cameroun: Diversité des procédés de production et qualité des produits. *Eur Sci J* 14 (36): 207-224. DOI: 10.19044/esj.v14n36p207
- Chevassus-Agnes JC, Favier, Joseph A. 1976. Traditional technology and the nutritive value of sorghum beer from Cameroon. *Cahiers de Nutrition et de Dietetique* (France).
- Ciriminna R, Albanese L, Di Stefano V, Delisi R, Avellone G, Meneguzzo F, Pagliaro M. 2018. Beer produced via hydrodynamic cavitation retains higher amounts of xanthohumol and other hops prenylflavonoids. *Lebensmittel-Wissenschaft und. Lwt* 91 : 160-167. DOI : 10.1016/j.lwt.2018.01.037
- CODEX Stan 247. 2005. In "Norme générale codex pour les jus et les nectars de fruits".
- Coelho E, Ballesteros LF, Domingues L, Vilanova M, Teixeira JA. 2020. Production of a distilled spirit using cassava flour as raw material: chemical characterization and sensory profile. *Molecules* 25 (14): 3228. DOI: 10.3390/molecules25143228.
- Cortacero-Ramírez S, Hernáinz-Bermúdez de Castro M, Segura-Carretero A, Cruces-Blanco C, Fernández-Gutiérrez A. 2003. Analysis of beer components by capillary electrophoresis methods. *Trends Anal Chem* 22 (7) : 440-445. DOI : 10.1016/S0165-9936(03)00704-0.
- Coulibaly WH, Bouatenin KM, Boli ZB, Alfred KK, Bi YC, N'sa KM, Cot M, Djameh C, Djè KM. 2020. Influence of yeasts on bioactive compounds content of traditional sorghum beer (tchapalo) produced in Côte d'Ivoire. *Cur Res Food Sci* 3 : 195-200. DOI : 10.1016/j.crf.2020.06.001
- Djangan D, Mbayhoudel K, Nandoum M. 2002. Organisation des unités de transformation artisanale en zone de savanes: Cas de la transformation du sorgho en bière locale bili-bili à Moundou au Tchad. In *Savanes africaines: des espaces en mutation, des acteurs face à de nouveaux défis*. Actes du colloque, Garoua, Cameroun. Cirad-Prasac.
- Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. 1956. Colorimetric method for determination of sugars and related substances. *Anal Chem* 28: 350-356. DOI: 10.1021/ac60111a017.
- Einfalt D. 2021. Barley-sorghum craft beer production with *Saccharomyces cerevisiae*, *Torulaspora delbrueckii* and *Metschnikowia pulcherrima* yeast strains. *Eur Food Res Technol* 247: 385-393 DOI : 10.1007/s00217-020-03632-7.
- Eze VC, Eleke OI, Omeh YS. 2011. Microbiological and nutritional qualities of burukutu sold in mammy market Abakpa, Enugu State, Nigeria. *Am J Food Nutr* 1 (3): 141-146. DOI: 10.5251/ajfn.2011.1.3.141.146
- Fadahusi IF, Ogunbanwo ST, Fawole AO. 2013. Microbiological and nutritional assessment of burukutu and pito (indigenously fermented alcoholic beverages in West Africa) during storage. *Nat Sci* 11 (4): 98-103.
- Ferreira IM, Guido LF. 2018. Impact of wort amino acids on beer flavour: A review. *Fermentation* 4 (2): 23. DOI: 10.3390/fermentation402003
- Gómez-Corona C, Escalona-Buendía HB, García M, Cholle S, Valentin D. 2016. Craft vs. industrial: Habits, attitudes and motivations towards beer consumption in Mexico. *Appetite* 96: 358-367. DOI : 10.1016/j.appet.2015.10.002. PMID:26455311
- Hazelwood LA, Daran JM, Van Maris AJA, Pronk JT, Dickinson JR. 2008. The Ehrlich pathway for fusel alcohol production: A century of research on *Saccharomyces cerevisiae* metabolism. *Appl Environ Microbiol* 74: 2259-2266. DOI : 10.1128/AEM.02625-07
- Hencké S. 2002. Utilisation alimentaire de levure, thèse de Doctorat en pharmacie Université Henri Poincaré-Nancy I.
- Humia BV, Santos K, Barbosa A, Sawata M, Mendonca M, Padilha F. 2019. Beer molecules and its sensory and biological properties. *Molecules* 24 (8) : 1568. DOI : 10.3390/molecules24081568
- Huot R, Roy GY. 1999. Livre De Chimie Organique Notions Fondamentales, Ed Carcajou 325-338.
- Kayodé APP, Adégbidi A, Linnemann AR, Nout MJR, Hounhouigan DJ. 2005. Quality of farmer's varieties of sorghum and derived foods as perceived by consumers in Benin. *Ecol Food Nutri* 44 (4): 271-294. DOI : 0.1080/03670240500187302

- Lariven J, Sonia R. 2017. Brasseurs de France la bière dossier de Presse. 2-19.
- Lyumugabe F, Kamaliza G, Bajyana E, Thonart PH. 2010. Microbiological and physico-chemical characteristic of Rwandese traditional beer "Ikigage". *Afr J Biotechnol* 9 (27): 4241-4246.
- Maoura N, Gaillardin MC, Jacques. 2006. Pourquoi Suivi technique, Analytique et Microbiologique de la « bili bili », Bière traditionnelle Tchadienne. *Afr J Biotechnol* 2 (1): 69-82. DOI: 10.4314/afsci.v2i1.61135
- Michel MC, Hannequart G. 1968. Dosage des acides aminés et amine par la ninhydrine. Amélioration pratique. In *Ann Biol Anim Biochim Biophysique* 8 (4): 557-563. DOI: 10.1051/rnd:19680408
- Miller GL. 1972. Use of Dinitrosalicylic Acid Reagent for determination of reducing sugars. *Anal Chem* 31: 426-428. DOI: 10.1021/ac60147a030
- Muyanja CMBK, Narvhus JA, Treimo J, Langsrud T. 2003. Isolation, characterisation and identification of lactic acid bacteria from bushera: A Ugandan traditional fermented beverage. *Int J Food Microbiol* 80 (3): 201-210. DOI : 10.1016/S0168-1605(02)00148-4
- N'da Coulibaly S. 1996. Etude comparative de procédés traditionnels de préparation d'une bière locale appelée tchapalo. 2ème journées annuelles de la SOACHIM, Dakar-Sénégal.
- Nascimento ESP, Cardoso DR, Franco DW. 2008. Quantitative ester analysis in Cachaça and distilled spirits by Gas Chromatography-Mass Spectrometry (GC-MS). *J Agric Food Chem* 56 (14): 5488-5493. DOI: 10.1021/jf800551d
- Ogunfa SA. 1985. African Fermented Foods. *Microbiol Fermented Foods* 2: 155-191. DOI: 10.1080/0142968X.1985.11904309
- Olaniran AO, Hiralal L, Mokoena M P, Pillay N. 2017. Flavour-active volatile compounds in beer: Production, regulation and control. *J Inst Brew* 123 (1): 13-23. DOI : 10.1002/jib.389.
- Ourega L, Ban K, Kouadio FN, Guessan G, Nemlin J, Tano K. 2015. Physico-chemical characteristics of plantain beers produced from waste plantain varieties of Côte d'Ivoire (Orishele, French Horn 2 and 1). *Int J Innov Appl Stud* 10 (2): 666-677.
- Perisse J, Adrian J, Rerat A, Le Berre S. 1959. Nutrient balance in the transformation of sorghum into beer. Preparation, composition and consumption of a Togo beer. In *Annales de la Nutrition et de l'Alimentation* 13: 1-15.
- Ranganna S. 1979. Titratable acidity. Manual of analysis of fruit and vegetable products.
- Roger DD, Venassius L, Justin EN, Franccedil E. 2013. Processing of Amgba: A sorghum-maize based beer, brewed in Cameroon. *J Brew Distilling* 4 (1): 11-18. DOI: 10.5897/JBD12.004
- Setta MC, Matemua A, Mbega ER. 2020. Potential of probiotics from fermented cereal-based beverages in improving health of poor people in Africa. *J Food Sci Technol* 57 (11): 3935-3946. DOI: 10.1007/s13197-020-04432-3.
- Suzuki K, Iijima K, Sakamoto K, Sami M, Yamashita H. 2006. A review of hop resistance in beer spoilage lactic acid bacteria. *J Inst Brew* 112 (2): 173-191. DOI : 10.1002/j.2050-0416.2006.tb00247.x
- Zhou Z, Ni W, Ji Z, Liu S, Han X, Li X, Mao J. 2020. Development of a rapid method for determination of main higher alcohols in fermented alcoholic beverages based on dispersive liquid-liquid microextraction and gas chromatography-mass spectrometry. *Food Anal Methods* 13: 591-600. DOI: 10.1007/s12161-019-01668-4

Comparative determination of total antioxidant effects of ethanol extract of *Phyllanthus amarus* leaves

OJOCHENEMI EJEH YAKUBU, MICHAEL SUNDAY ABU*, JOHN AKIGHIR, JOB ITANYI ONUCHE, ALI ARABI

Department of Biochemistry, Faculty of Pure and Applied Sciences, Federal University Wukari. PMB 1020 Katsina-Ala Road, Wukari, Taraba State, Nigeria. Tel.: +234-803-0000000, *email: abulex77@yahoo.com

Manuscript received: 3 September 2021. Revision accepted: 16 December 2021.

Abstract. Yakubu OE, Abu MS, Akighir J, Onuche JI, Arabi A. 2021. Comparative determination of total antioxidant effects of ethanol extract of *Phyllanthus amarus* leaves. *Asian J Nat Prod Biochem* 19: 81-85. Some medicinal and therapeutic potency have been reported of *Phyllanthus* in traditional settings. Hence, the current study evaluated the total antioxidant potential, phenolics and flavonoids of fractions of ethanol extract of *Phyllanthus amarus* leave obtained using column chromatography. Extraction was done using absolute ethanol; elution was performed using different organic solvents of varying polarities, from n-Hexane, Chloroform, Ethylacetate, Ethanol, Methanol and finally, water. The results obtained from the study revealed that fraction 10b (83 mg/L), has the highest concentration of Total Antioxidant Capacity (TAC), the next being 1a, 5a, 11b (48mg/mL). The reduction in the TAC of the fractions was in the order of 3b>7a>6a>5b>4a> with F7b (13mg/mL) having the lowest TAC. The result for Total Phenolic Content (TPC) shows that fraction 10b (430 mg/ml), has the highest concentration of TPC, just like in antioxidant activity, followed by F4b (428 mg/mL). Fraction 2a (9 mg/ml) has the lowest concentration of TPC. Total Flavonoid Content (TFC) showed that fraction 3b (201 mg/mL) has the most concentration of total flavonoids, followed by 4b (120mg/mL). Meanwhile, fractions 1a, 2b and 9a have the same TFC. While fraction 2a showed the lowest concentration (25mg/mL) of flavonoids. The plant extract fractions were found to contain more both phenolics and flavonoids which may have contributed to the antioxidant activity observed.

Keywords: Antioxidant, ethanol extract, flavonoids, phenolic, *Phyllanthus amarus*

INTRODUCTION

Phyllanthus amarus is of the Euphorbiaceae family, while the genus is *Phyllanthus*. The species is in *amarus* and can be readily found in tropical and subtropical countries of the world (Trease and Evans 2001). It is known in Nigeria among Yoruba as “eyin olobe,” Hausa as “geerontsuntsaaayee,” and Igbo as “Ite knwonwa nazu,” and in English as leaf flower or stone breaker. The plant is usually prepared as an herbal mixture and used by Nigerians for health purposes and is considered to have great economic importance (Umoh et al. 2013). Previously, the whole-plant extract has been used for the treatment of urinary disorders, liver disease, dyspepsia, anorexia, constipation, and dysentery (Samy et al. 2008), while many female anomalies like leucorrhoea, menorrhagia, and mammary abscess have been managed by this plant exudate (Samuel and Andrews 2010). Again, fresh leaf paste prepared from *phyllanthus amarus* has the potential to alleviate white spots on the skin, manage diabetes, and reduce the incidence of jaundice (Ignacimuthu et al. 2008). Similarly, malaria has been successfully managed by *phyllanthus amarus* whole plant extract (Kuppusamy and Murugan 2010).

Methanolic extract of *Phyllanthus amarus* was found to have an antioxidant effect that inhibited lipid oxidation. It masked up reactive oxygen species in vitro (Meena et al.

2018), while ethanolic extract of *Phyllanthus amarus* hairy roots showed antiproliferative activity against apoptosis initiated by increased intracellular reactive oxygen species (Abhyankar et al. 2010). The effect of nor-securinine, an alkaloid extracted from *Phyllanthus amarus*, was examined against spore growth of some fungi as well as powdery pea mildew (*E. pisi*) and was found to be potent (Meena et al., 2018). *Phyllanthus amarus* ethanol, aqueous, and hexane extracts showed restriction of LPS-induced production of NO and PGE2 and decreased the LPS-induced secretion of Tumor necrosis (Kiemer et al. 2003). The combination of *Phyllanthus amarus* and *Andrographis paniculata* plant extracts demonstrated active snake anti-venom potential that could be useful against snakebites (Meena et al. 2018). Similarly, Sornakumar et al. (2014) reported that about 0.24 mg of di-herbal plant extracts comprising *Phyllanthus amarus* effectively neutralized the cobra venom-induced toxicity.

Certainly, *Phyllanthus amarus* has been reported to have medicinal and therapeutic potency in traditional settings. These medicinal properties are mostly implicated in the presence of phytochemicals which are believed to exert some specific pharmacological effects on the body. Hence, the current study evaluated the total antioxidant capacity, phenolics, and flavonoids of GC-MS partially fractionated ethanol extract of *Phyllanthus amarus* leaves.

MATERIALS AND METHOD

Plant material

Fresh, healthy-looking leaves of *Phyllanthus amarus* were collected within the premises of Federal University Wukari, Taraba State. *Phyllanthus amarus* was identified and authenticated by Dr. Yakubu J.O.E, a Pharmacologist in the Department of Biochemistry, Faculty of Pure and Applied Science Federal University Wukari, Taraba State. The leaves were air-dried in the laboratory at 25°C for one week, pulverized manually using laboratory mortar and pestle into a smooth particle, and stored in an air-tight container for further use.

Solvent extraction

About 500 g of the pulverized plant leaves were soaked in 1.3 L of 97 % ethanol for 48 hours at 25°C with periodic shaking to ensure maximum extract yield, according to Yakubu et al. (2014). The crude extract was filtered using clean white sieving mesh and Whatman number 1 filter paper. The filtrate was evaporated to dryness with a rotary evaporator at low temperature under reduced pressure to obtain green crude extract.

Fractionation of ethanol extract by column chromatography

Column chromatography to separate the extract into its component fractions. Silica gel was used as the stationary phase, while different solvents of increasing polarity were used as the mobile phase. According to Yakubu et al. (2014), the bottom part of the glass column was packed with glass wool with the help of a glass rod. Exactly 235 g of silica gel of mesh size 60-200 was dissolved in 225 mL absolute n-Hexane to make the slurry (activating silica gel). The chromatographic column (30 mm diameter by 40 mm height) was packed with silica gel, and the solvent was allowed to flow into a conical flask below freely. At the end of the packing process, all taps were locked, the setup was allowed to stand for 24 hours, and the clear solvent at the top of the silica gel was allowed to drain down the meniscus.

Elution

The Yakubu et al. (2014) method was adopted for the elution. Two grams of the ethanol extract of *Phyllanthus amarus* leaves was dissolved in 15 % absolute ethanol, and the solution was applied into the chromatography column (30 mm in diameter and 400 mm in height), followed by elution of the extract using n-hexane, chloroform, ethyl acetate, ethanol, methanol, and finally distilled water. The following ratios (v/v) in mL of solvent combinations were frequently used in the elution process: n-Hexane: chloroform 100:0, 50:50; Chloroform: ethylacetate 100:0, 50:50; Ethylacetate: ethanol 100:0, 50:50; Ethanol: methanol 100:0 50:50; Methanol: distilled water 100:0 50:50; Distilled water:100. A measured volume (200 mL) of each solvent combination was poured into the column using a separator funnel. The eluted fractions were collected in aliquots of 100 mL in conical flasks.

Determination of total antioxidant capacity (TAC) (Singleton et al. 1999)

The absorbance was measured in triplicate for each fraction. Total antioxidant capacity (TAC) was calculated as mg/mL of Trolox equivalent (TE) using the regression equation from the calibration curve. Exactly 0.05g of the extract was dissolved in 250 mL of methanol to prepare 0.2mm. One mL of methanol extract was added to 2mL of DPPH in each test tube, followed by 100 µL of the fraction to each test tube. The mixture was shaken vigorously and left in the dark at room temperature for 30 minutes. The absorbance of the solution was measured immediately at 517 nm in a UV-Visible spectrophotometer. The experiment was performed in duplicate.

Determination of total flavonoids content (TFC) (Chang et al. 2002)

Quercetin standard was used for derivation of the calibration curve. The total flavonoids were expressed as mg/mL quercetin equivalent (QE). A 10% aluminium chloride solution was prepared by dissolving 10 g of aluminium chloride (ALCL₃) in 100mL ethanol. Exactly 100 µL of diluted sample was added to the test tube containing 1.5 mL methanol followed by 100 µL of aluminium chloride (ALCL₃) and 100 µL potassium acetate solutions. The mixture was incubated at room temperature for 30 minutes. The absorbance of the reaction mixture was measured at 765 nm wavelength using a UV-Visible spectrophotometer using distilled water as the blank.

Determination of total phenolic content (TPC) (Yakubu et al. 2014)

The Folin ciocalteu reagent was diluted (1:10 using distilled water). A volume of 100 µL of the crude sample was added to each test tube, and 2.5mL of Folin ciocalteu reagent was added to each test tube, followed by 2mL of Na₂CO₃ solution. The solution was incubated for 15 minutes at 45°C. The absorbance was measured at 415 nm in a UV-Visible spectrophotometer using distilled water as the blank.

RESULTS AND DISCUSSION

Total antioxidant capacity (TAC)

Fractions obtained from ethanolic extract of *Phyllanthus amarus* leaves. Fraction 10b (83 mg/mL) has the highest concentration of total antioxidant capacity, followed by F1a, F5a, and F11b, which all have similar TAC values (48 mg/mL). Other fractions, F4b, F9b, F10a are also of the same value (44 mg/mL) while fractions, 1b, 2a, 2b, 3a, 6a, 8b, and 9a (23 mg/mL) possess the same TAC. The decrease observed in the TAC of the fractions is in the order of F3b>F7a>F6a>F5b>F4a>, with F7b (13mg/mL) showing the lowest total antioxidant capacity.

Total phenolic content (TPC)

The result for total phenolic content of ethanol extract of *Phyllanthus amarus* leaves shows that fraction 10b (430 mg/mL) has the highest concentration of total phenolic content as it was in the case of antioxidant activity. This was followed by F4b (428 mg/mL), while fraction 2a (9 mg/mL) has the lowest concentration of TPC, as shown in Figure 2.

Total flavonoid content (TFC)

The total flavonoid content (TFC) of ethanol extract of *Phyllanthus amarus* leaves revealed that fraction 3b (201 mg/mL) possesses the highest concentration of total flavonoid content, as depicted in Figure 3, followed by 4b (120 mg/mL). Fractions 1a, 2b, and 9a have the same TFC. The decrease observed in the TFC of the fractions is in the order of 8b>5a>5b>7a>11a>7b. Meanwhile, F2a has the lowest concentration (25 mg/mL).




Figure 1. Total antioxidant capacity (TAC). F1: n-Hexane: (100:0), F2: n-Hexane/chloroform: (50:50), F3: Chloroform: (100:0), F4: Chloroform/ethyl acetate (50:50), F5: ethyl acetate: (100:0), F6: Ethyl acetate/ethanol (50:50), F7: Ethanol (100:0), F8: Ethanol/Methanol: (50:50), F9: Methanol: (100:0), F10: Methanol/water (50:50), F11: Distilled water: (100:0)




Figure 2. Total phenolic content (TPC). F1: n-Hexane: (100:0), F2: n-Hexane/chloroform: (50:50), F3: Chloroform: (100:0), F4: Chloroform/ethyl acetate (50:50), F5: Ethyl acetate: (100:0), F6: Ethyl acetate/ethanol (50:50), F7: Ethanol (100:0), F8: Ethanol/Methanol: (50:50), F9: Methanol: (100:0), F10: Methanol/water (50:50), F11: Distilled water: (100:0)




Figure 3. Total flavonoid content (TFC). F1: n-Hexane: (100:0), F2: n-Hexane/chloroform: (50:50), F3: Chloroform: (100:0), F4: Chloroform/ethyl acetate (50:50), F5: Ethyl acetate: (100:0), F6: Ethyl acetate/ethanol (50:50), F7: Ethanol (100:0), F8: Ethanol/Methanol: (50:50), F9: Methanol: (100:0), F10: Methanol/water (50:50), F11: Distilled water: (100:0)

Discussion

Phyllanthus amarus extract has attracted several researchers over the decades because of its potent pharmaceutical uses. Every country has its traditional use of *Phyllanthus amarus*, but the way of curing disease is almost common everywhere (Meena et al. 2018). Antioxidant compounds usually deactivate free radicals and inhibit the process of lipid peroxidation, which is one of the major reasons for the deterioration of food and biological cells (Halliwell 2010).

In this study, a considerable high level of antioxidant activity was observed in the ethanol extract of *Phyllanthus amarus* leaves. The TAC revealed that fractions 10b (84 mg/mL), 1a, and 5a (48 mg/mL), possessed higher antioxidant capacity, implying that the fractions have higher antioxidant values when compared with the other fractions. This observation was substantiated by the positive correlation that was seen between the TAC and the concentration of phenolic compounds present in the fractions; since it has been demonstrated by several studies that the higher the phenolic content, the higher the TAC. The result also indicated that methanol-water (50:50) is the most efficient solvent for elution of antioxidant activity of ethanolic extract of *Phyllanthus amarus* leaves, as reported by Yakubu et al. (2018). However, the low antioxidant capacity observed in fractions 4a (17mg/mL) and 7b (13mg/mL) could be attributed to the fact that antioxidant activity depends on the type, polarity of the extracting solvent, and purity of active compounds (Demiray et al. 2009).

Plants have diverse phenolic compounds, including simple phenolics, phenolic acids, anthocyanins, hydroxycinnamic acid derivatives, and flavonoids, which have radical scavenging power (Olamide et al., 2017). The result obtained for the TPC shows that fraction 10b (430 mg/mL) has the highest concentration of total phenolic content. Yakubu et al. (2019b) stated that phenolic compounds are plants' most effective and abundant natural

antioxidants. They are very important plant constituents because their hydroxyl groups confer scavenging ability (Motawi et al., 2011). Thus, it can be inferred with certainty that these phenolics are responsible for the marked antioxidant activity of these fractions, which is compatible with several studies reported on the relationships between phenolic content and antioxidant activity (Yakubu et al., 2018).

In conclusion, different methods employed in the study revealed that the ethanolic extract of *Phyllanthus amarus* leaves is rich in phenolic and flavonoid compounds, demonstrated good antioxidant activity, and can be useful in masking up free radicals as assayed through various in vitro models in this study. Thus, preventing free-radical related disorders respectively. There appears to be no correlation between antioxidant activity and flavonoid content, implying that the plant extract contains several phytochemicals other than flavonoids which may contribute to the antioxidant capacity. Hence, supporting the use of this plant as a good source of natural antioxidants should be encouraged.

REFERENCES

- Abhyankar G, Suprasanna P, Pandey BN, Mishra KP, Rao KV, Reddy VD. 2010. Hairy root extract of *Phyllanthus amarus* induces apoptotic cell death in human breast cancer cells. *Innov Food Sci Emerg Technol* 11: 526-532. DOI: 10.1016/j.ifset.2010.02.005.
- Ankute DK, Eziah VY, Afreh-Nuamah K. 2021. Insecticidal activities of indigenous plants from Volta Region, Ghana in managing *Sitophilus zeamais* and *Prostephanus truncatus* stored maize. *Intl J Trop Drylands* 5: 48-60. DOI: 10.13057/tropdrylands/t050202
- Chang CC, Yang MH, Wen HM, Chern JC. 2002. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal* 10: 178-182. DOI: 10.38212/2224-6614.2748.
- Demiray S, Pintado ME, Castro PML. 2009. Evaluation of phenolic profiles and antioxidant activities of Turkish medicinal plants: *Tilia argentea*, *Crataegi folium* leaves and *Polygonum bistorta* roots. *World Acad Sci Eng Technol* 54: 213-217

- Halliwell B. 2010. Free radical and antioxidant-Quo Vadis? Trends Pharmacol Sci 32: 125-130. DOI: 10.1016/j.tips.2010.12.002.
- Ignacimuthu S, Ayyanar M, Sankarasivaraman K. 2008. Ethnobotanical study of medicinal plants used by Paliyar tribals in Theni District of Tamil Nadu, India. Fitoterapia 79: 562-568. DOI: 10.1016/j.fitote.2008.06.003.
- Kiemer AK, Hartung T, Huber C, Vollmar AM. 2003. *Phyllanthus amarus* has anti-inflammatory potential by inhibition of iNOS, COX-2, and cytokines via the NF- κ B pathway. J Hepatol 38: 289-297. DOI: 10.1016/S0168-8278(02)00417-8.
- Kuppusamy C, Murugan K. 2010. In-vitro antimalarial activity of traditionally used western ghats plants from India and their interactions with chloroquine against chloroquine resistant *Plasmodium falciparum*. Parasitol Res 107: 1351-1364. DOI: 10.1007/s00436-010-2005-9.
- Meena J, Sharma RA, Rashmi R. 2018. A review on phytochemical and pharmacological properties of *phyllanthus amarus* Schum and Thonn. Intl J Pharm Sci Res 9 (4): 1377-1386.
- Motawi TK, Hamed MA, Shabana MH, Hashem RM, Aboul AF. 2011. *Zingiber officinale* acts as aneutral agent against liver fibrosis. Nutr Metab 40: 7075-7078. DOI: 10.1186/1743-7075-8-40.
- Olamide EA, Funsho OO, Tan NH, Zeng GZ. 2017. In vitro antioxidant activity, total phenolic and flavonoid contents of ethanol extract of stem and leaf of *Grewia carpinifolia*. Beni-Suef Univ J Basic Appl Sci 6: 10-14. DOI: 10.1016/j.bjbas.2016.12.003.
- Samuel JK, Andrews B. 2010. Traditional medicinal plant wealth of Pachalur and Periyur Hamlets, Dindigul District, Tamil Nadu. Indian J Tradit Knowl 9: 264-270. DOI: 10.6026/97320630003100.
- Samy RP, Pushparaj PN, Gopalakrishnakone P. 2008. A compilation of bioactive compounds from Ayurveda. Bioinformation 3: 100-110.
- Singleton VL, Orthofer RM, Ramuela-Raventos RM. 1999. Analysis of total phenols and antioxidants and other substrates by means of Folin-Ciocalteu reagent. Methods Enzymol 299: 152-178. DOI: 10.1016/S0076-6879(99)99017-1.
- Sornakumar RSA, Kunthavai PC, Gnaniah S. 2014. Isolation, purification and characterization of active compound from *Andrographis paniculata* L. and *Phyllanthus amarus* Linn. and testing the antivenom activity of the di-herbal extract by in-vitro and in-vivo studies. Am J Clin Nutr 84: 95-135.
- Trease GE, Evans WC. 2001. Text Book of Pharmacognosy. 14th Ed. WB Saunders Company Limited, King's College London, UK.
- Umoh ED, Ukana DA, Ito EU. 2013. Phytochemical screening and nutrient analysis of *Phyllanthus amarus*. Asian J Plant Sci Res 3 (4): 116-122.
- Yakubu OE, Kayode AA, Christopher S, Terungwa JI, Kenneth C. 2019. Chemical composition of baobab leaves and fractionation of its ethanolic extract using column chromatography. Intl J Sci Res 8 (7): 812-821. DOI: 10.21275/ART20199083.
- Yakubu OE, Olawale O, Edobor PKI, Silas VT, Bilyaminu H. 2018. Fractionation and determination of antioxidant activities, of the leaves of *Senna Occidentalis* ethanol extract. J Pharm Pharmacol 6 (3): 26-30.
- Yakubu OE, Nwode OFC, Joshua PE, Ugwu MN, Odu AD, Okwo F. 2014. Fractionation and determination of total antioxidant capacity, total phenolics and flavonoids content of aqueous, ethanol and n-hexane extract of *Vitexdonia leaves*. Afr J Biotechnol 13 (5): 693-698. DOI: 10.5897/AJB2013.13225.